



Synergistic Effects of Dietary Vitamin C and Selenium on Induced Methylmercury Toxicity in Juvenile Olive Flounder *Paralichthys olivaceus*

Seunghan Lee¹, Gunhyun Park¹, Mohammad Moniruzzaman¹, Jinho Bae¹, Yujin Song¹, Kang-Woong Kim², Sungchul C. Bai¹*

¹Pukyong National University, Department of Marine Bio-materials & Aquaculture/Feeds & Foods Nutrition Research Center, Busan, Korea.

²National Institute of Fisheries Science (NIFS), Aquafeed Research Center, Pohang, 791-923, Korea.

* Corresponding Author: Tel.: +82.51 6297922; Fax: +82.51 6286873;
E-mail: scbai@pknu.ac.kr

Received 01 September 2016
Accepted 24 November 2016

Abstract

The synergistic effects of dietary vitamin C and selenium (Se) on induced methylmercury (MeHg) toxicity in juvenile olive flounder *Paralichthys olivaceus* were examined in this study. Nine diets containing 3 different vitamin C levels (0, 200 and 400 mg kg⁻¹ diet in the form of L-ascorbyl-2-monophosphate), 3 different Se levels (0, 2 and 4 mg kg⁻¹ diet in the form of selenomethionine) at a constant level of MeHg (20 mg kg⁻¹ diet in the form of MeHg) were formulated and fed to triplicate groups of juvenile olive flounder with mean weight of 2.00±0.04 g (mean±SD) in semi recirculation system using 3² factorial design. Growth performance and tissue Hg burden were determined after 8 weeks of feeding. Fish fed diets containing 400 mg kg⁻¹ vitamin C together with 2 and 4 mg kg⁻¹ Se (C₄₀₀Se₂ and C₄₀₀Se₄) showed significantly (P<0.05) higher weight gain (WG), specific growth rate (SGR) and feed efficiency (FE). Whereas fish which were under the C₄₀₀Se₄ diet exhibited significantly (P<0.05) higher and protein efficiency ratio (PER) than other feeding groups. Tissue Hg burden in muscle, liver and kidney showed a tendency of increasing with decreasing the levels of vitamin C and Se. However, significantly low tissue Hg burden was observed from fish fed diets containing 400 mg kg⁻¹ vitamin C together with 2 and 4 mg kg⁻¹ Se (C₄₀₀Se₂ and C₄₀₀Se₄). The results suggested that tissue Hg burden could be reduced and MeHg mediated growth problems could be ameliorated by supplementing dietary vitamin C and Se in juvenile olive flounder.

Keywords: Mercury, bioaccumulation, tissue burden, growth.

Introduction

Fish is a good source of two long chain omega 3 polyunsaturated fatty acids (n-3 lcPUFAs) such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA). Fish consumption is also known to reduce the risk of death due to coronary heart disease by 36% (Mozaffarian & Rimm, 2006). Nevertheless, organic compounds such as polychlorinated biphenyls (PCBs), dioxins and insecticides together with inorganic chemicals such as arsenic, cadmium, lead, mercury, copper, zinc and iron can contaminate fish and seafood in general (FAO, 2005). As a result, concerns have been raised in recent years about contaminants found in fish originated from environmental pollution.

Among all of these contaminants, mercury (Hg) is the major contaminant and ubiquitous environmental toxin (Moniruzzaman *et al.*, 2015) which is appearing to be a threat to human health. In fact, both inorganic and organic Hg may be found in fish (Lee *et al.*, 2016). However, methylmercury (MeHg) is the predominant form of Hg in fish (Hoffman, Rattner, Burton, & Cairns, 2002). Its

chemical properties allow it to rapidly diffuse and tightly bind to proteins in aquatic biota, including the proteins in muscle tissue of fish. This leads to bioaccumulation in the fish, with the Hg level increasing with age of the fish. In turn, biomagnification along the food chain leads to higher Hg levels in piscivorous fish that are higher in the food chain. The ability of MeHg to biomagnify in aquatic food chains was responsible for past epidemics in human populations and is a continuing concern for both human and environmental health (Ausili *et al.*, 2008; Driscoll, Mason, Chan, Jacob, & Pirrone, 2013).

Commercial fish diets are composed of various ingredients and Hg contamination in commercial fish feeds is mainly due to the high metal levels in the raw materials (Wang, Onsanit, & Dang, 2012). For this reason dietary exposure is one of the main routes of Hg contamination in fish (Choi & Cech, 1998). As the result, Hg has been regarded as undesirable substance in animal feed (EFSA, 2008). Fish consumption is one of the main paths through which human exposure to MeHg occurs (Passos, Mergler, Lemire, Fillion, & Guimaraes, 2007). Seafood contamination by Hg is a

public health concern particularly in countries with high rate of fish consumption such as Korea. Daily seafood consumption in Korea has reached 50.6 g, which accounted for 3.8% of the total food ingested (Moon, Kim, Choi, Yu, & Choi, 2009; Choi, Moon, & Choi, 2012). Consequently, blood Hg level in a representative sample taken from Korean adult population found to be associated with fish consumption (Kim & Lee, 2010). Moon *et al.* (2011) suggested that implementation of systematic monitoring programs for seafood contaminations by Hg are necessary in Korea. The various toxic effects induced by Hg in biological systems are often due to alterations in the antioxidant defense system (Sheweita, 1998; Berntssen, Waagbø, Toften, & Lundebye, 2003; Alves, Rosa, & Santana, 2007; Berg, Puntervoll, Valdernesnes, & Goksøyr, 2010).

Vitamin C is a potent antioxidant and enzyme cofactor which can protect indispensable molecules in the body, such as proteins, lipids, carbohydrates, and nucleic acids (DNA and RNA), from damage by free radicals and reactive oxygen species (ROS) that are generated during normal metabolism, by active immune cells, and through exposure to toxins (Combs Jr., 2012). Vitamin C also participates in redox recycling of other important antioxidants; for example, vitamin C is known to regenerate vitamin E from its oxidized form (Bruno *et al.*, 2006).

Selenium (Se) as an essential element required in small amounts to maintain good health, plays a role in antioxidant defenses and is a cofactor for the antioxidant enzyme glutathione peroxidase. Selenium interacts with the accumulation and toxicity of Hg in aquatic organisms in various ways. Because of its biological importance and nutraceutical component Se has attracted various researchers. It is involved in numerous biological functions including, preventing oxidative damage, maintaining homeostasis of thyroid

hormone, enhancing immune functions (Hoffmann & Berry, 2008). Fish and crayfish treated with Se showed decreased Hg content, and fish in Se contaminated areas have been shown to contain reduced amounts of Hg (Southworth, Peterson, & Turner, 1994; Southworth, Peterson, & Ryon, 2000).

Various antioxidants including vitamin C, vitamin E, and Se are known to decrease Hg toxicity in Japanese quail (Kung, Soares, & Haltman, 1987) and in various other organisms (Chapman & Chan, 2000). Vijayalakshmi, Bapu and Sood (1992) and Bapu, Vijayalakshmi and Sood (1994) also examined the effects of vitamin C treatment after subcutaneous injections of methylmercuric chloride (MeHgCl) for 7 days in mice and found improvements in recoveries of enzymes activities. Chen *et al.* (2006) demonstrated that selenoproteins help eliminate ROS induced changes by metals because of their antioxidant properties. In this study, we evaluated effects of dietary vitamin C and Se levels on induced Hg accumulation in juvenile olive flounder *Paralichthys olivaceus*, a most commercially important marine aquaculture fish species in Republic of Korea (Lee *et al.*, 2016).

Materials and Methods

Experimental Diets

Composition of the semi-purified basal diet is shown in Table 1. Nine diets contain three different vitamin C levels (0, 200 and 400 mg kg⁻¹ diet in the form of L-ascorbyl-2-monophosphate) and three different Se levels (0, 2 and 4 mg kg⁻¹ diet in the form of selenomethionine) with similar Hg toxicity levels (20 mg Hg kg⁻¹ diet in the form of MeHg) were formulated. The 20 mg MeHg level kg⁻¹ diet was chosen based on previous study from our lab. In diets

Table 1. Composition of the experimental diets (% dry matter basis)

| Ingredients | Diets | | | | | | | | |
|--|--------------------------------|--------------------------------|--------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
| | C ₀ Se ₀ | C ₀ Se ₂ | C ₀ Se ₄ | C ₂₀₀ Se ₀ | C ₂₀₀ Se ₂ | C ₂₀₀ Se ₄ | C ₄₀₀ Se ₀ | C ₄₀₀ Se ₂ | C ₄₀₀ Se ₄ |
| Casein ¹ | 32 | 32 | 32 | 32 | 32 | 32 | 32 | 32 | 32 |
| Defatted fish meal ² | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 | 25 |
| Wheat flour ³ | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 | 18 |
| Corn starch ³ | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| Fish oil ⁴ | 9.6 | 9.6 | 9.6 | 9.6 | 9.6 | 9.6 | 9.6 | 9.6 | 9.6 |
| Vitamin premix (C free) ⁵ | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| Minerals premix (Se free) ⁶ | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.0 |
| Hg-premix | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 |
| Cellulose | 6.0 | 5.0 | 4.0 | 4.0 | 3.0 | 2.0 | 2.0 | 1.0 | 0.0 |
| Vitamin C premix | 0.0 | 0.0 | 0.0 | 2.0 | 2.0 | 2.0 | 4.0 | 4.0 | 4.0 |
| Se-premix | 0.0 | 1.0 | 2.0 | 0.0 | 1.0 | 2.0 | 0.0 | 1.0 | 2.0 |

¹ United States Biochemical (Cleveland, OH) 44122.

² Suhyup Feed Co. Ltd.

³ Young Nam Flour Mills Co., Pusan, Korea.

⁴ E-Wha oil Co., Ltd., Busan Korea.

⁵ Contains (as mg kg⁻¹ diet): dl-calcium pantothenate, 150; choline bitartrate, 3,000; inositol, 150; menadione, 6; niacin, 150; pyridoxine-HCl, 15; riboflavin, 30; thiamine mononitrate, 15; retinyl acetate, 6; biotin, 1.5; folic acid, 5.4; B12, 0.06; cholecalciferol, 2.4

⁶ Contains (as mg kg⁻¹ diet): Al, 1.2; Ca, 5000; Cl, 100; Cu, 5.1; Co, 9.9; Na, 1280; Mg, 520; P, 5000; K, 4300; Zn, 27; Fe, 40; I, 4.6; Mn, 9.1.

supplemented with a MeHg and ascorbic acid sources, an equivalent amount of cellulose was removed. In a 3² factorial design 9 experimental diets (C₀Se₀, C₀Se₂, C₀Se₄, C₂₀₀Se₀, C₂₀₀Se₂, C₂₀₀Se₄, C₄₀₀Se₀, C₄₀₀Se₂ and C₄₀₀Se₄) were formulated to be isonitrogenous and isoenergetic, containing 50% crude protein (CP) and 16.7 kJ available energy g⁻¹ diet (16.7, 16.7 and 37.7 kJ g⁻¹ for protein, carbohydrate and lipid respectively). Vitamin free casein was used as the main protein sources. All the ingredients were mixed completely and then pelleted by using 1-mm- and 2-mm-diameter dies (Bai & Lee, 1998). After processing, all the diets were packed into small bags and kept at -20°C until use.

Experimental Fish and Feeding Trials

Juvenile olive flounder, *Paralichthys olivaceus* were obtained from Tong-Yeong, Korea. Prior to the start of feeding trial, fish were fed the basal diet for 10 days to adjust to the semi-purified diet and to deplete possible body reserves of vitamin C. The feeding trial was conducted in a semi-circulated system with 30 L aquariums receiving filtered sea water at a rate of 2 L min⁻¹. Supplemental aeration was provided to maintain dissolved oxygen near saturation and water temperature was kept at 20±1°C. Experimental fish averaging 2.00±0.04 g (mean±SD) were randomly distributed into each aquarium as a group of 20 fish. Each diet was fed to triplicate groups to satiation level three times a day at a feeding rate of 2.0 to 3.5% of wet body weight. Total fish weight in each aquarium was determined every three weeks and the amount of diet fed to fish was adjusted accordingly. Aquariums were kept clean during the experiment time to minimize algae and fungal growth which could provide a source of vitamin C to cultured fish.

Sample Collection and Analysis

Weight gain (WG), specific growth rate (SGR), feed efficiency (FE), protein efficiency ratio (PER), and survival rate were measured and calculated after weight checking. After the final weighing, 3 fish was randomly removed from each aquarium for analysis.

Vitamin C Analysis

Ascorbic acid concentration was determined by High Performance Liquid Chromatography (HPLC; Dionex Softron, Sunnyvale, CA, USA). The ultraviolet detector was set at 254 nm and the mobile phase was 0.05 M KH₂PO₄ with flow rate of 1.0 mL min⁻¹. Weighed samples were homogenized in 10% cold metaphosphoric acid. Homogenates were centrifuged at 3000×g for 20 minutes and supernatants were analyzed after filtered through a 0.45 µm pore size syringe filter.

Selenium Analysis

Diet and tissue Se concentrations were assessed by the digestion of samples in nitric acid. Weighed samples were put into a 250 mL Kjeldahl flask, and 50 mL of HNO₃ was added to the flask. Then, the flask was heated in a heating mantle until the sample was fully digested. Approximately 5 mL of H₂O₂ was added to make sure that the sample was totally digested, and the digested sample was diluted with H₂O. The concentration of Se in the diluted digest solution was determined using a Perkin-Elmer 3300 Inductively Coupled Plasma Mass Spectrometer (ICP-MS, Perkin-Elmer, Waltham, MA, USA).

Mercury Analysis

Over 90% of Hg present in fish is MeHg. For this reason, direct total concentration of Hg was measured instead of MeHg (Bloom, 1992; Amlund, Lundebye, & Berntssen, 2007). Hg analyzer (DMA-80, Milestone, Inc., Shelton, CT) was used to determine tissue Hg concentration following the method similar to the one used in Lee et al. (2011). A certified reference material (DORM-2 dogfish liver, National Research Council, Canada) was used simultaneously during the analyses.

Statistical Analysis

Data were analyzed by two-way ANOVA to test for the effect of dietary treatments using SPSS version 16.0 (SPSS Inc., Chicago, IL, USA). Least Significant Difference (LSD) was used to compare means when significant difference between treatments was observed and P-values of 0.05 or less (P≥0) were considered to be statistically significant.

Results

Growth Performance

Growth performance of fish is summarized in Table 2. Fish fed 400 mg kg⁻¹ vitamin C comprising diets (C₄₀₀Se₀, C₄₀₀Se₂ and C₄₀₀Se₄) showed significantly (P<0.05) higher WG at all Se levels, than other feeding groups. Even though no significant difference was observed between feeding groups which were under 200 mg kg⁻¹ vitamin C containing diets at all Se levels, all of them showed significantly (P<0.05) higher WG than those groups which were not supplemented with vitamin C. Specific growth rate of fish which supplemented 400 mg kg⁻¹ vitamin C with 2 and 4 mg kg⁻¹ Se (C₄₀₀Se₂ and C₄₀₀Se₄) appeared to be significantly (P<0.05) higher than the other feeding groups. In general, feeding groups which were under 400 mg kg⁻¹ vitamin C categories (C₄₀₀Se₀, C₄₀₀Se₂ and C₄₀₀Se₄) exhibited significantly higher SGR than groups which were given 0 and 200 mg kg⁻¹ vitamin C at all Se levels. No significant

Table 2. Growth performance of juvenile olive flounder fed the experimental diet for 8 weeks¹

| Diets | WG (%) ² | SGR (% day ⁻¹) ³ | FE (%) ⁴ | PER ⁵ |
|----------------------------------|---------------------|---|---------------------|-------------------|
| C ₀ Se ₀ | 123 ^l | 1.66 ^e | 39.0 ^e | 0.76 ^e |
| C ₀ Se ₂ | 127 ^l | 2.16 ^l | 64.0 ^d | 1.27 ^d |
| C ₀ Se ₄ | 214 ^{de} | 2.38 ^d | 64.2 ^d | 1.26 ^d |
| C ₂₀₀ Se ₀ | 166 ^{cd} | 2.04 ^e | 43.6 ^e | 0.86 ^e |
| C ₂₀₀ Se ₂ | 242 ^{cd} | 2.56 ^{cd} | 67.0 ^d | 1.30 ^d |
| C ₂₀₀ Se ₄ | 267 ^{bcd} | 2.71 ^c | 78.0 ^c | 1.53 ^c |
| C ₄₀₀ Se ₀ | 310 ^{abc} | 2.94 ^b | 91.6 ^b | 1.81 ^b |
| C ₄₀₀ Se ₂ | 334 ^{ab} | 3.05 ^{ab} | 93.6 ^{ab} | 1.83 ^b |
| C ₄₀₀ Se ₄ | 374 ^a | 3.24 ^a | 102 ^a | 2.06 ^a |
| Pooled SEM ⁶ | 16.1 | 0.09 | 3.95 | 0.08 |
| Two-way ANOVA | | | | |
| Vitamin C | 0.0001 | 0.0001 | 0.0001 | 0.0001 |
| Selenium | 0.0001 | 0.0001 | 0.0001 | 0.0001 |
| Vitamin C × Selenium | 0.0100 | 0.0001 | 0.0064 | 0.0086 |

¹ Values are means from groups (n=3) of fish where the values in each row with different superscripts are significantly different (P<0.05).

² Weight gain (%) = (final weight - initial weight) × 100 / initial weight

³ Specific growth rate (%) = 100 × (Ln final wt. - Ln initial wt.) / days

⁴ Feed efficiency (%) = (wet weight gain / dry feed intake) × 100

⁵ Protein efficiency ratio = (wet weight gain / protein intake)

⁶ Pooled standard error of means: SD/√n

difference in SGR observed between fish fed 200 mg kg⁻¹ vitamin C at 2 and 4 mg kg⁻¹ Se levels (C₂₀₀Se₂ and C₂₀₀Se₄). Similarly, significantly (P<0.05) higher FE observed in (C₄₀₀Se₄) feeding group. However, no significant differences were observed between fish fed C₀Se₂, C₂₀₀Se₂, or C₀Se₄ and between fish fed C₂₀₀Se₀ or C₀Se₀ diets. Protein efficiency ratio followed quite similar pattern like FE where, fish fed C₄₀₀Se₄ diet exhibited significantly (P<0.05) higher PER than other feeding groups.

Tissues Mercury Burden

Muscle of fish fed diets containing 400 mg kg⁻¹ vitamin C along with Se (C₄₀₀Se₂ and C₄₀₀Se₄) showed significantly (P<0.05) lower Hg burden than other feeding groups. Whereas the control group (C₀Se₀) deposited significantly (P<0.05) higher Hg in their muscle (Table 3). Tissue Hg burden showed a tendency of increasing with decreasing levels of vitamin C and Se. Similarly, liver Hg burden appeared to be significantly lower for fish which fed 400 mg kg⁻¹ vitamin C diets at 2 and 4 mg kg⁻¹ Se levels. However, no significant difference in liver Hg burden was observed between fish fed C₀Se₀, C₀Se₂ or C₀Se₄ diets and between fish fed C₀Se₂, C₀Se₄ or C₂₀₀Se₀ diets. Kidney Hg deposition followed similar pattern like muscle where fish fed C₄₀₀Se₂ and C₄₀₀Se₄ diets accumulated significantly lower Hg than other feeding groups.

Discussion

In the present study, olive flounder fed MeHg containing diet (C₀Se₀) demonstrated clear growth depression possibly due to the higher accumulation of Hg in fish. Similar results have been found from the

one of our recent study dealt with the effects of dietary vitamin C on inorganic mercury (HgCl₂) toxicity in juvenile olive flounder (Lee et al., 2016). Moreover, in the present study, the poor growth performance observed in olive flounder fed diets which do not contain either vitamin C and/or Se might have occurred due to decreased enzyme activity, altered structural functionality and transport process problems caused by Hg accumulation (Zalups & Lash, 1994). However, when the diets were supplemented with higher level of Se (C₀Se₄), significant growth improvement was observed in fish which may be due to the antioxidative nature of selenium. Interestingly, the diet supplied with higher level of vitamin C (C₄₀₀Se₀) showed more pronounced effect in terms of improvement of WG, SGR, FE and PER in fish compared with the high level of Se (C₀Se₄) containing diet. The results suggest that vitamin C might have more protective effect than Se in terms of growth improvement in fish. In addition, when the diets were supplied with higher level of vitamin C (400 mg/kg diet) with irrespective level of Se (2 or 4 mg/kg diet) showed significant growth improvement in fish compared to rest of the diets. Lee et al. (2016) reported that dietary vitamin C (100 or 200 mg kg⁻¹ diet) could significantly improve the growth performance of olive flounder on inorganic mercury (HgCl₂) induced toxicity which is in agreement of the present study. In this study, the two-way ANOVA also demonstrated that there is a significant interactive effect between dietary vitamin C and Se on growth performance in terms of WG, SGR, FE and PER of fish on induced methylmercury toxicity. These results suggested that dietary vitamin C and Se had synergistic effects on reduction of methylmercury toxicity in this fish species.

In this study, dietary methylmercury

Table 3. Total tissue mercury concentrations ($\mu\text{g g}^{-1}$ of wet matter basis) juvenile olive flounder fed the experimental diets for 8 weeks¹

| Diets | Muscle | Liver | Kidney |
|----------------------------------|--------------------|--------------------|-------------------|
| C ₀ Se ₀ | 15.2 ^a | 18.4 ^a | 23.7 ^a |
| C ₀ Se ₂ | 13.6 ^b | 17.6 ^{ab} | 21.3 ^b |
| C ₀ Se ₄ | 12.1 ^c | 17.6 ^{ab} | 18.2 ^c |
| C ₂₀₀ Se ₀ | 12.0 ^c | 16.6 ^b | 18.5 ^c |
| C ₂₀₀ Se ₂ | 10.7 ^d | 15.2 ^c | 16.6 ^d |
| C ₂₀₀ Se ₄ | 9.7 ^{ef} | 13.2 ^d | 14.8 ^e |
| C ₄₀₀ Se ₀ | 10.1 ^{de} | 13.0 ^d | 15.0 ^e |
| C ₄₀₀ Se ₂ | 9.0 ^g | 11.6 ^e | 12.5 ^f |
| C ₄₀₀ Se ₄ | 9.3 ^{fg} | 11.4 ^e | 12.1 ^f |
| Pooled SEM ² | 0.37 | 0.49 | 0.69 |
| Two-way ANOVA | | | |
| Vitamin C | 0.0001 | 0.0001 | 0.0001 |
| Selenium | 0.0001 | 0.0001 | 0.0001 |
| Vitamin C × Selenium | 0.0031 | 0.0154 | 0.0128 |

¹ Values are means from groups (n=3) of fish where the values in each row with different superscripts are significantly different (P<0.05).

² Pooled standard error of means: SD/ \sqrt{n}

accumulated in the tissues, in increasing order, muscle<liver<kidney. The highest deposition of Hg was found in kidney tissue and the lowest in muscle tissue of fish which is in agreement with Lee *et al.* (2016). The results of the present study suggest that accumulation of organic mercury (methylmercury) is higher than inorganic mercury accumulation in juvenile olive flounder which was reported by Lee *et al.* (2016). Our results also showed that higher amounts of Hg accumulated in muscle tissue of fish compared to that of reported by Lee *et al.* (2016). It has been reported that a substantial amount of organic mercury can accumulated in fish muscle (NRC, 2005). In this study, the amount of Hg deposited in fish muscle is very close to the Hg contents in liver and kidney. However, as the dietary vitamin C and Se levels increases in the diets, Hg contents in the muscle, liver and kidney tissues of fish also decreases. Significantly lowest amount of Hg was found in fish fed the higher level of vitamin C with lower/higher levels of Se in diets which supports the growth performance data of the present study. In present study, both of the dietary vitamin C and Se have shown their strong effects in reducing mercury contents in tissue levels of fish. In addition, dietary vitamin C and Se exhibited their interaction effect in reducing Hg contents in muscle, liver and kidney tissue of fish. However, Lee *et al.* (2016) found interaction effect between dietary vitamin C and Hg in kidney tissue only on induced HgCl₂ toxicity in olive flounder. Therefore, in the present study, the results clearly demonstrated the synergistic effects of dietary vitamin C and Se on induced MeHg toxicity in terms of tissue mercury reduction.

In the present study, the better growth performance and lower tissue Hg burden observed in fish fed diets supplemented with dietary vitamin C (200 and 400 mg kg⁻¹) together with Se (2 and 4 mg kg⁻¹) might, presumably, attribute to the protective role of vitamin C and Se against MeHg induced immune suppression and their ability to maintain/or

elevate the activities of several key antioxidants. Consequently, effective disposal of Hg from SH groups by vitamin C along with its ability to inhibit, minimize and remove free radicals might have reduced tissue Hg burden and helped to improve the poor growth performance observed in this study which is in agreement with Durak, Kalender, Uzun, Demir and Kalender (2010).

Conclusion

In this study, the results showed that the supplemental Se might have ensured adequate levels of Se and replaced the amount of Se lost to Hg sequestration. This in turn helped the normal selenoprotein synthesis to continue. On the other hand, vitamin C likely inhibited free radical formation and lipid peroxidation that could have been caused by MeHg. Therefore, we may conclude that poor growth performance resulted from induced MeHg could be enhanced and Hg burden on muscle, liver and kidney could be reduced by supplementing dietary vitamin C (400 mg kg⁻¹ diet) together with Se (2 or 4 mg kg⁻¹ diet) in juvenile olive flounder.

Acknowledgments

This research was supported by National Research Foundation (NRF), Republic of Korea with grant number NRF-2011-0016221. We also would like to thank the members of the Feeds and Foods Nutrition Research Center (FFNRC), Pukyong National University, Busan, Rep. of Korea.

References

- Alves, R.R.N., Rosa, I.L., & Santana, G.G. (2007). The role of animal-derived remedies as complementary medicine in Brazil. *BioScience*, 57(11), 949-955. <http://dx.doi.org/10.1641/B571107>
- Amlund, H., Lundebye, A-K., & Berntssen, M.H.G. (2007).

- Accumulation and elimination of methylmercury in Atlantic cod (*Gadus morhua* L.) following dietary exposure. *Aquatic Toxicology*, 83(4), 323-330. <http://dx.doi.org/10.1016/j.aquatox.2007.05.008>
- Ausili, A., Torrecillas, A., Aranda, F.J., Mollinedo, F., Gajate, C., Corbalán-García, S., de Godos, A., & Gómez-Fernández, J.C. (2008). Edelfosine Is Incorporated into Rafts and Alters Their Organization. *The Journal of Physical Chemistry B*, 112(37), 11643-11654. <http://dx.doi.org/10.1021/jp802165n>
- Bai, S.C., & Lee, K.-J. (1998). Different levels of dietary dl- α -tocopheryl acetate affect the vitamin E status of juvenile Korean rockfish, *Sebastes schlegelii*. *Aquaculture*, 161(1-4), 405-414. [http://dx.doi.org/10.1016/S0044-8486\(97\)00288-3](http://dx.doi.org/10.1016/S0044-8486(97)00288-3)
- Bapu, C., Vijayalakshmi, K., & Sood, P.P. (1994). Comparison of monothiols and vitamin therapy administered alone or in combinations during methylmercury poisoning. *Bulletin of Environmental Contamination and Toxicology*, 52(2), 182-189. <http://dx.doi.org/10.1007/BF00198486>
- Berg, K., Puntervoll, P., Valdernes, S., & Goksøyr, A. (2010). Responses in the brain proteome of Atlantic cod (*Gadus morhua*) exposed to methylmercury. *Aquatic Toxicology*, 100(1), 51-65. <http://dx.doi.org/10.1016/j.aquatox.2010.07.008>
- Berntssen, M.H.G., Waagbø, R., Toften, H., & Lundebye, A.K. (2003). Effects of dietary cadmium on calcium homeostasis, Ca mobilization and bone deformities in Atlantic salmon (*Salmo salar* L.) parr. *Aquaculture Nutrition*, 9(3), 175-183. <http://dx.doi.org/10.1046/j.1365-2095.2003.00245.x>
- Bloom, N.S. (1992). On the chemical form of mercury in edible fish and marine invertebrate tissue. *Canadian Journal Fisheries and Aquatic Science*, 49(5), 1010-1017. <http://dx.doi.org/10.1139/f92-113>
- Bruno, R.S., Leonard, S.W., Atkinson, J., Montine, T.J., Ramakrishnan, R., Bray, T.M., & Traber, M.G. (2006). Faster plasma vitamin E disappearance in smokers is normalized by vitamin C supplementation. *Free Radical Biology & Medicine*, 40(4), 689-697. <http://dx.doi.org/10.1016/j.freeradbiomed.2005.10.051>
- Chapman, L., & Chan, H.M. (2000). The influence of nutrition on methylmercury intoxication. *Environmental Health Perspectives*, 108 (Suppl 1), 29-56. <http://dx.doi.org/10.1289/ehp.00108s129>
- Chen, C., Yu, H., Zhao, J., Li, B., Qu, L., Liu, S., Zhang, P., & Chai, Z. (2006). The roles of serum selenium and selenoproteins on mercury toxicity in environmental and occupational exposure. *Environmental Health Perspectives*, 114(2), 297-301. <http://dx.doi.org/10.1289/ehp.7861>
- Choi, M., Moon, H.B., & Choi, H.G. (2012). Intake and potential health risk of butyltin compounds from seafood consumption in Korea. *Archives of Environmental Contamination and Toxicology*, 62(2), 333-340. <http://dx.doi.org/10.1007/s00244-011-9688-5>
- Choi, M.H., & Cech, J.J. (1998). Unexpectedly high mercury level in pelleted commercial fish feed. *Environmental Toxicology and Chemistry*, 17(10), 1979-1981. <http://dx.doi.org/10.1002/etc.5620171013>
- Combs, Jr. G.F. (2012). Chapter 9 - Vitamin C. In: COMBS, GERALD F. (ed.) *The Vitamins (Fourth Edition)*. Academic Press, San Diego, USA.
- Driscoll, C.T., Mason, R.P., Chan, H.M., Jacob, D.J., & Pirrone, N. (2013). Mercury as a Global Pollutant: Sources, Pathways, and Effects. *Environmental Science and Technology*, 47(10), 4967-4983. <http://dx.doi.org/10.1021/es305071v>
- Durak, D., Kalender, S., Uzun, F.G., Demir, F., & Kalender, Y. (2010). Mercury chloride-induced oxidative stress in human erythrocytes and the effect of vitamins C and E in vitro. *African Journal of Biotechnology*, 9(4), 488-495. <http://dx.doi.org/10.5897/AJB09.1314>
- EFSA, (European Food Safety Authority). (2008). Opinion of the scientific panel on contaminants in the food chain on a request from the European Commission on mercury as undesirable substance in feed. *The European Food Safety Authority Journal*, 654(2), 1-76. <http://dx.doi.org/10.2903/j.efsa.2008.654>
- FAO, Food and Agriculture Organization of the United Nations. (2005). *Fisheries and Aquaculture Topics: Fish contaminants. Topics Fact Sheets* [Online]. Fisheries and Aquaculture Department, Food and Agriculture Organization of the United Nations (FAO). <http://www.fao.org/fishery/topic/14815/en> [Accessed 26 September 2014].
- Hoffman, D.J., Rattner, B.A., Burton, G.A., & Cairns, J. (2002). *Handbook of Ecotoxicology. Second Edition*, ed., Taylor & Francis. ISBN 9781566705462
- Hoffmann, P.R., & Berry, M.J. (2008). The influence of selenium on immune responses. *Molecular Nutrition & Food Research*, 52(11), 1273-1280. <http://dx.doi.org/10.1002/mnfr.200700330>
- Kim, N.-S., & Lee, B.-K. (2010). Blood total mercury and fish consumption in the Korean general population in KNHANES III, 2005. *Science Total Environment*, 408(20), 4841-4847. <http://dx.doi.org/10.1016/j.scitotenv.2010.06.026>
- Kung, L.J., Soares, J.H. Jr., & Haltman, W.A. (1987). Effect of vitamin E and synthetic antioxidants on the survival rate of mercury-poisoned Japanese quail. *Poultry Science*, 66(2), 325-331. <http://dx.doi.org/10.3382/ps.0660325>
- Lee, J.H., Moniruzzaman, M., Yun, H., Lee, S., Park, Y., & Bai, S.C. (2016). Dietary vitamin C reduced mercury contents in the tissues of juvenile olive flounder (*Paralichthys olivaceus*) exposed with and without mercury. *Environmental Toxicology and Pharmacology*, 45(5), 8-14. <http://dx.doi.org/10.1016/j.etap.2016.05.009>
- Lee, J.W., Riu, N.D., Lee, S., Bai, S.C., Moniello, G., & Hung, S.S.O. (2011). Effects of dietary methylmercury on growth performance and tissue burden in juvenile green (*Acipenser medirostris*) and white sturgeon (*A. transmontanus*). *Aquatic Toxicology*, 105(3-4), 227-234. <http://dx.doi.org/10.1016/j.aquatox.2011.06.013>
- Lee, S., Moniruzzaman, M., Bae, J., Seong, M., Song, Y., Dosanjh, B., and Bai, S.C. (2016). Effects of extruded pellet and moist pellet on growth performance, body composition, and hematology of juvenile olive flounder, *Paralichthys olivaceus*. *Fisheries and Aquatic Sciences*, 19, 32. <http://dx.doi.org/10.1186/s41240-016-0032-x>
- Moon, H.B., Kim, H.S., Choi, M., Yu, J., & Choi, H.G. (2009). Human health risk of polychlorinated biphenyls and organochlorine pesticides resulting from seafood consumption in South Korea, 2005-2007. *Food and Chemical Toxicology*, 47(8), 1819-1825. <http://dx.doi.org/10.1016/j.fct.2009.04.028>
- Moon, H.B., Kim, S.J., Park, H., Jung, Y.S., Lee, S., Kim,

- Y.H., & Choi, M. (2011). Exposure assessment for methyl and total mercury from seafood consumption in Korea, 2005 to 2008. *Journal of Environmental Monitoring*, 13(9), 2400-2405. <http://dx.doi.org/10.1039/C1EM10504C>
- Moniruzzaman, M., Park, G., Yun, H., Lee, S., Park, Y., & Bai, S.C. (2015). Synergistic effects of dietary vitamin E and selenomethionine on growth performance and tissue methylmercury accumulation on mercury-induced toxicity in juvenile olive flounder, *Paralichthys olivaceus* (Temminck et Schlegel). *Aquaculture Research*, 1-11. <http://dx.doi.org/10.1111/are.12904>
- Mozaffarian, D., & Rimm, E.B. (2006). Fish intake, contaminants, and human health: Evaluating the risks and the benefits. *The Journal of the American Medical Association*, 296(15), 1885-1899. <http://dx.doi.org/10.1001/jama.296.15.1885>
- NRC, (National Research Council). (2005). *Mineral tolerance of animals, Second Revised* ed. Washington, D.C., National Academies Press, USA.
- Passos, C.J., Mergler, D., Lemire, M., Fillion, M., & Guimaraes, J.R. (2007). Fish consumption and bioindicators of inorganic mercury exposure. *Science Total Environment*, 373(1), 68-76. <http://dx.doi.org/10.1016/j.scitotenv.2006.11.015>
- Sheweita, S.A. (1998). Heavy metal-induced changes in the Glutathione levels and Glutathione Reductase/Glutathione S-Transferase activities in the liver of male mice. *International Journal of Toxicology*, 17(4), 383-392. <http://dx.doi.org/10.1080/109158198226224>
- Southworth, G.R., Peterson, M.J., & Ryon, M.G. (2000). Long-term increased bioaccumulation of mercury in largemouth bass follows reduction of waterborne selenium. *Chemosphere*, 41(7), 1101-1105. [http://dx.doi.org/10.1016/S0045-6535\(99\)00562-7](http://dx.doi.org/10.1016/S0045-6535(99)00562-7)
- Southworth, G.R., Peterson, M.J., & Turner, R.R. (1994). Changes in concentrations of selenium and mercury in largemouth bass following elimination of fly ash discharge to a quarry. *Chemosphere*, 29(1), 71-79. [http://dx.doi.org/10.1016/0045-6535\(94\)90091-4](http://dx.doi.org/10.1016/0045-6535(94)90091-4)
- Vijayalakshmi, K., Bapu, C., & Sood, P.P. (1992). Differential effects of methylmercury, thiols, and vitamins on galactosidases of nervous and non-nervous tissues. *Bulletin of Environmental Contamination and Toxicology*, 49(1), 71-77. <http://dx.doi.org/10.1007/BF00193343>
- Wang, W-X., Onsanit, S., & Dang, F. (2012). Dietary bioavailability of cadmium, inorganic mercury, and zinc to a marine fish: Effects of food composition and type. *Aquaculture*, 356-357(8), 98-104. <http://dx.doi.org/10.1016/j.aquaculture.2012.05.031>
- Zalups, R., & Lash, L. (1994). Recent advances in understanding the renal transport and toxicity of mercury. *Journal of Toxicology and Environmental Health*, 42(1), 1-44. <http://dx.doi.org/10.1080/15287399409531861>