



## Phylogenetic Relations and Electrophoretic Identification of Allozyme in Four Species of Snappers

Govinda Rao, V<sup>1,\*</sup>, N. Muddula Krishna<sup>1</sup>, K. Sujatha<sup>1</sup>

<sup>1</sup> Department of Marine Living Resources, Andhra University, Visakhapatnam, Andhra Pradesh, India- 530003.

\* Corresponding Author: Tel.: +9440395067;  
E-mail: govindarao.mlrau@gmail.com

Received 20 September 2016  
Accepted 22 March 2017

### Abstract

In the present study, four species of snappers, *Lutjanus fulvus*, *lemniscatus*, *lutjanus* and *madras* were embodying the catches of Visakhapatnam coast, analyzed for allozyme variations. The four enzyme systems screened five loci was 12 in *fulvus*, 3 in *lemniscatus*, 29 in *lutjanus* and 26 in *madras* were found to be polymorphic at  $p = 0.95$  level. Diagnostic alleles help in clear distinctive species had been identified. An unweighted pair-group method with arithmetic mean (UPGMA) method was used to construct the dendrogram to know the genetic relationship between the four species of snappers *Lutjanus fulvus*, *lemniscatus*, *lutjanus* and *madras* based on genetic distance revealed the presence of three clusters showing the percentage similarity between *lutjanus* and *madras* 0.961%, *lemniscatus* showed 0.918% similarity with above two species and *fulvus* showed 0.890% similarity with above three species.

**Keywords:** Allozyme, snapper, phylogenesis, allele.

### Introduction

*Lutjanidae* family is one of the prime in the order Perciformes and comprises 4 subfamilies, 17 genera and 110 species make the *lutjanids* are the most specious of the families Froese and Pauly (2016) and (Nelson, 2006). Snappers have been the focused of several allozyme studies. However, given the number of species in the family *lutjanids* have been under represented in such studies to date. Taxonomy is basic thing to conserve the efforts of marine fishery samples and the units on which conservation is based are determined ultimately by species. Snappers are well-known by their coloring patterns or a suite of morphologic and characteristics like body configuration, size and by meristic characters. Identification was based on morphological features leads to confusion in closely resembling species. Possibility of misidentification poses problems for management of fishery and stock structure analysis (Ward *et al.*, 1995). Species identification Genetic levels of snappers also extremely valuable since morphological discrimination especially of early juveniles can be difficult. Therefore the combined efficacy of morphological analyses and biochemical phylogenetic methods have been followed here for providing a meaningful approach to solve taxonomic uncertainties concerning the status of species (Chow *et al.*, 1993). Electrophoretic data in allozyme and

isozyme form can be applied in systematic and taxonomic investigations. Under such circumstances when morphological characters are unreliable, biochemical genetic methods have long been used to corroborate the species identity (Avisé, 1975; Utter, 1991 and Chow *et al.*, 1993). In fisheries, these methods have been used to disclose cryptic species (Shaklee and Tamaru, 1981; Smith *et al.*, 1991; Grant and Utter, 1984) to resolve taxonomic problems (Smith and Robertson, 1981; Graves *et al.*, 1988). Nevertheless, *lutjanids* have contributed to our understanding of genetic variation in natural populations and an assortment of evolutionary process. Virtually that all previous applications to snappers systematics study should be reevaluated. Evaluation of these applications and recommendations for standard methods and future research are also discussed in present paper. Previously such studies were carried out by Ovenden *et al.*, (2004) studied the genetic diversity of *Pristipomoides multidens* from East Indies triangle waters. Vasconcellos *et al.*, (2008) studied nuclear and mitochondrial DNA sequence differences in between Brazilian and Caribbean population of yellow snapper, *Ocyurus chrysurus*. Sulaiman *et al.*, (2003) investigate the genetic diversity of red snapper *Lutjanus malabaricus* in Brunei and Sabah using allozyme electrophoresis. So far no attempt has been made to analyse the genetic structure of *lutjanids* off

Visakhapatnam coast. Understanding the present genetic makeup of wild *lutjanid* populations has a significant practical value particularly for fishery management and conservation of stocks. Hence in the present study, general proteins and allozyme based survey of genetic variation in the four species of snappers *Lutjanus fulvus*, *L. lemniscatus*, *L. lutjanus* and *L. madras* represented in the catches of Visakhapatnam, east coast of India. Govinda Rao et al., 2014 and Muddula Krishna et al., 2015 studied length weight relationship and length groups of two species of snappers, diversity of intertidal fish species works done from Visakhapatnam coastal waters, east coast of India.

### Material and Methods

A total of 70 fresh specimens of all size groups belonging to four species of snappers of genus *Lutjanus fulvus*, *lemniscatus*, *lutjanus* and *madras* that were collected from traditional fish landing centers and Visakhapatnam fishing harbour, east coast of India (Figure 1) during 2010-2012 in the catches. After species identification, total length, weight and masculinity of each specimen were noted and immediately muscle tissue was taken and stored at -20°C for further analysis. Muscle tissue crushed and homogenized with extraction buffer (0.2 M Tris EDTA buffer pH 7.0). Homogenates were centrifuged for 1 hour at 10,000 rpm at 4°C and supernatant was recentrifuged for 30 minutes. The supernatant

collected was used for further analysis. Vertical polyacrylamide gel electrophoresis was used for separation of allozymes at different enzyme loci. Gels of 10\*8 cm size were used. Electrophoresis was carried out at 100 V in cooling chamber at 4°C. The bands of each of four enzymes were visualized using specific histochemical staining methods (Whitmore, 1990), until sharp bands were visualized. Enzyme systems studied in allozyme analysis nomenclature of loci and alleles was followed as recommended by Shaklee et al., (1990). At all the loci, most common allele was designated as 100. Alternate alleles designated as per their mobility in relation to the most common allele. Calculation of allele frequencies and tests for conformity to Hardy – Weinberg expectations (probability test) were undertaken using GENEPOP version 3.4 software (Raymond and Rousset, 1995a).

### Results

Allozyme banding pattern of 70 specimens belonging to four species of snappers of genus *Lutjanus*: *Lutjanus fulvus* (Figure 2); *L. lemniscatus* (Figure 3); *L. lutjanus* (Figure 4); *L. madras* (Figure 5) that were represented in the catches of Visakhapatnam, east coast of India was studied. The four enzyme systems were Alcohol Dehydrogenase (Figure 7), Lactate dehydrogenase (Figure 8), Phosphoglucumutase (Figure 9) and Super Oxide Dismutase (Figure 10) in four species genus *Lutjanus*:




Figure 1. Outline map of India showing sampling station.

*fulvus*, *lemniscatus*, *lutjanus* and *madras* were studied. Of the four systems studied, eight loci were detected in four species of lutjanids. In the present study, single locus for the allozymes Alcohol Dehydrogenase (ADH\*) and Super Oxide Dismutase (SOD\*) and three loci each for Lactate Dehydrogenase (LDH-1\*, LDH-2\*, LDH-3\*) and Phosphoglucomutase (PGM-1\*, PGM-2\*, PGM3\*) were detected. A total of 21 alleles were detected and their frequencies were given in Table 1. The locus ADH\* was expressed by four alleles, loci LDH-1\* and SOD\* PGM-1\*, PGM-2\*, PGM3\* were expressed by three alleles and remaining all loci by two alleles each.

## Discussion

Diagnostic locus where no alleles shared with any other species (Ward et al., 1995) for *lemniscatus* was observed at LDH-2\* and for *lutjanus* at LDH-3\*. Among the four species between one and four

diagnostic alleles were detected (Table 2). In ADH\* alleles *a* and *d* observed to be commonly found in all four species. In LDH-1\* allele observed to be commonly shared by the four species. Locus LDH-2\* is expressed only in *lemniscatus* and LDH-3\* only in *lutjanus*. Thus they can be referred to as diagnostic loci.


From Indian waters, Menezes (1993, 1994) studied genetic relationship among three species of genus *Sardinella* and four Sciaenid species from Arabian Sea, genetic and morphometric differences between yellowtail snapper populations of tropical West Atlantic were studied by Vasconcellos (2008) using allozymes and mitochondrial DNA was observed that morphometric differences were not incompatible with the existence of discreet fish stocks. Miller and Cribb (2007a) studied mitochondrial DNA based phylogenetic relationships among Indo-Pacific snappers. In this study, the precise cause to the divergence could perhaps given by carrying out further studies with sample from other



Figure 2. *Lutjanus fulvus*, 164 mm, TL.



Figure 3. *Lutjanus lemniscatus*, 296 mm TL.



**Figure 4.** *Lutjanus lutjanus* , 203 mm, TL.



**Figure 5.** *Lutjanus madras*, 211 mm, TL.

areas of this region. Estimates of genetic identification and genetic distance between pairs of four species of snappers given in Table 2. The present study based on allozyme, UPGMA dendrogram (Figure 6) has been constructed using POPGENE software version 3.4 which illustrated that *lutjanidae* species were closely related. This study assists us to evaluate the variation of phylogenetic existing among species variation and also derive diagnostic loci and alleles that are useful for evaluation of closely related species of this region.

The present study is an attempt to indicate the genetic structure of these four species using general proteins and allozymes as there were no previous studies on these species. In recent years there has been a worldwide diminishing the fishing stocks (Garcia and Grainger, 2005) overexploited, fully exploited and exhausted fisheries that were 69% in 1995 increased to 75% in 2002, only 1% of the stocks were in a state of retrieval (Ormerod, 2003). For this reason competent management policies based on unambiguous scientific data are necessary both to

defend the fishing stocks and to maximize their exploitation without compromising their integrity. The correct stock as assessment and identification of species is essential to ascertain the maximum sustainable effort of a given marine resource (Ryman and Utter, 1987; Ryman, 1991). Many stock concepts can be found in the literature (Booke, 1981; Ovenden, 1990; Carvalho and Hauser, 1995), but one of the most accepted and used is “a stockist an intraspecific group of randomly mating individuals with temporal and spatial integrity” (Ihssen *et al.*, 1981) which swathe most of the definitions given by other authors. The fish stock identification can be made efficiently by the use of highly polymorphic molecular markers (Blaber *et al.*, 2005; Caddy and Seijo, 2005).

Several species of snappers are morphologically similar and even the meristic counts are almost identical for these species. Identification based purely on morphological features leads to confusion in closely resembling species. Possibility of misidentification poses problems for management of fishery and stock structure analysis. As an aid to




Fig. 7. Alcohol dehydrogenase




Fig. 8. Lactate dehydrogenase




Fig. 9. Phosphoglucumutase




Fig.10. Superoxide Dismutase

**Figure 7-10.** Allozyme banding pattern in four species of genus *Lutjanus*. 1,2- *fulvus*; 3,4- *lemniscatus*; 5,6- *lutjanus*; 7 to 9- *madras*; 10- tracking dye.

traditional taxonomic characters, biochemical genetic methods have been used in systematics to solve taxonomic ambiguities. Proteins are frequently used as biochemical genetic markers and the projected genetic variations are used to distinguish closely related species and natural populations of a given species. However, recent advancements in the field of molecular biology proved that allozymes are powerful biochemical genetic markers compared to proteins as they generate more reliable data. So far no attempt has been made to analyse the genetic structure of snappers of Indian waters as until recently they have

not received due attention from geneticists given their commercial significance and need for information on stock structure to ensure sustainable management. As understanding the present genetic makeup of wild snappers population has a significant practical value particularly for fishery management and conservation of stocks.

#### Acknowledgement


Authors are grateful to the Head, Department of Marine Living Resources, College of Science and

**Table 1.** Allele frequencies at nine loci in the four species of genus *Lutjanus* represented in the catches of Visakhapatnam

Locus	Allele	<i>fulvus</i>	<i>lemniscatus</i>	<i>lutjanus</i>	<i>madras</i>
ADH*	100(a)	0.75	0.75	0.846	0.75
	102(b)	-	0.20	-	-
	104(c)	-	0.05	-	0.167
	106(d)	0.25	-	0.154	0.083
	<i>n</i>	12	3	29	26
LDH-1*	100(a)	0.927	0.75	0.769	0.875
	102(b)	0.073	-	0.231	0.125
	104(c)	-	0.25	-	-
	<i>n</i>	12	3	29	26
LDH-2*	100(a)	-	0.964	-	-
	102(b)	-	0.036	-	-
	<i>n</i>	12	3	29	26
LDH-3*	100 (a)	-	-	0.973	-
	102 (b)	-	-	0.027	-
	<i>n</i>	12	3	29	26
PGM-1*	100(a)	0.968	0.875	0.95	0.972
	102(b)	0.032	0.125	0.05	0.28
	<i>n</i>	12	3	29	26
PGM -2*	98(a)	0.075	0.025	0.168	0.005
	100(b)	0.925	0.975	0.832	0.930
	102(c)	-	-	-	0.065
	<i>n</i>	12	3	29	26
PGM -3*	98(a)	-	0.925	0.90	-
	100(b)	-	0.075	0.10	-
	<i>n</i>	12	3	29	26
SOD*	100(a)	0.908	0.875	0.925	0.918
	102(b)	-	0.024	0.075	0.082
	104(c)	0.092	0.101	-	-
	<i>n</i>	12	3	29	26

**Table 2.** Alleles present at each locus, given as letters in alphabetic order according to their anodal mobility 'a' representing the fastest migrating. Diagnostic alleles are underlined

Locus	<i>fulvus</i>	<i>lemniscatus</i>	<i>lutjanus</i>	<i>madras</i>
ADH*	<u>ad</u>	<u>abc</u>	<u>ad</u>	<u>acd</u>
LDH-1*	<u>ab</u>	<u>a<math>\bar{c}</math></u>	<u>ab</u>	<u>ab</u>
LDH-2*	-	<u>ab</u>	-	-
LDH-3*	-	-	<u>ab</u>	-
PGM -1*	<u>ab</u>	<u>ab</u>	<u>ab</u>	<u>ab</u>
PGM -2*	<u>ab</u>	<u>ab</u>	<u>ab</u>	<u>ab<math>\bar{c}</math></u>
PGM -3*	-	<u>ab</u>	<u>ab</u>	-
SOD*	<u>ac</u>	<u>abc</u>	<u>ab</u>	<u>ab</u>

**Figure 6.** UPGMA dendrogram showing hierarchic relationship between four species of genus *Lutjanus*.

Technology, Andhra University for providing facilities to carry out this work.

## References

- Avise, J.C. (1975). Systematic value of electrophoretic data. *Systematic Zoology*, 23, 465-81.  
<http://dx.doi.org/10.1086/400132>
- Blaber, S.J.M., Dichmont, C.M., Buckworth, R.C., Badrudin, S.B., Nurhakim, S., Iskandar, B., Fegan, B., Ramm, D.C., & Salini, J.P. (2005). Shared stocks of snappers (Lutjanidae) in Australia and Indonesia: Integrating biology, population dynamics and socio-economics to examine management scenarios. *Review of Fish Biology and Fisheries*, 15, 111-127.  
<http://dx.doi.org/10.1007/s11160-005-3887-y>
- Booke, H. E. (1981). The conundrum of the stock concept: are nature and nature definable in fisheries science. *Canadian Journal of Fisheries and Aquatic Sciences*, 38, 1479-1480.  
<http://dx.doi.org/10.1139/f81-200>
- Caddy, J. F., & Seijo, J.C. (2005). This is more difficult than we thought. The responsibility of scientists, managers and stakeholders to mitigate the unsustainability of marine fisheries. *Philosophical Transactions of the Royal Society B*, 360, 59-75.  
<http://dx.doi.org/10.1098/rstb.2004.1567>
- Carvalho, G. R., & Hauser, L. (1995). Molecular genetics and the stock concept in fisheries. In: Carvalho, G. R., Pitcher, T. J (eds) *Molecular Genetics in Fisheries*. Chapman & Hall, London., 55-79 pp.
- Chow, S., Clarke, M.E., & Walsh, P.J. (1993). PCR-RFLP analysis on thirteen Western Atlantic snappers (Subfamily Lutjanidae): A simple method for species and stock identification. *Fishery Bulletin*, 91: 619-627.  
<http://dx.doi.org/10.7755/FB.111.2.1>
- Froese, R., & Pauly, D. (2015). Fish Base. Retrieved from <http://www.fishbase.org/html>
- Garcia, S. M., & Grainger, R.J.R. (2005). Gloom and doom? The future of marine capture fisheries. *Philosophical Transactions of the Royal Society of Biological Sciences*, 360: 21-46.  
<http://dx.doi.org/10.1098/rstb.2004.1580>
- Govinda Rao, V., Krishna, N.M., & Sujatha, K. (2014). Length weight relationship and length groups of two species of snappers (Pisces: Lutjanidae) represented in the catches of Visakhapatnam, Middle East coast of India. *Indian Journal of Experimental Zoology India*, 17: 127-132.
- Grant, W. S., & Utter, F.M. (1984). Biochemical population genetics of Pacific herring, *Clupea pallasii*. *Canadian Journal of Fisheries and Aquatic Sciences*, 41: 856-864.  
<http://dx.doi.org/10.1139/f84-102>
- Graves, J.E., Simovich, M.A., & Schaefer, K.M. (1988). Electrophoretic identification of early juvenile yellow fin tuna, *Thunnus albacares*. *Fishery Bulletin*, <http://dx.doi.org/00288330.1994.951660>
- Ihssen, P.E., Booke, H.E., Casselman, J.M., Mcglade, J.M., Pavne, N.R. and Utter, F.M. 1981. Stock identification: materials and methods. *Canadian Journal of Fisheries and Aquatic Sciences*, 38: 1838-1855.  
<http://dx.doi.org/10.1139/f81-230>
- Menezes, M. R. (1993). Interspecific genetic divergence in three pomfret species from the Goa region. *Aquaculture and Fisheries Management*, 24: 341-346
- Menezes, M. R. (1994). Little genetic variation in the oil sardine, *Sardinella longiceps* Val., from the western coast of India. *Australian Journal of Marine and Freshwater Research*, 45: 257-264.  
<http://dx.doi.org/10.1071/MF9940257>
- Miller, T.L., & Cribb, T.H. (2007a). Coevolution of Retrovarium n. gen. (Digenea: Cryptogonimidae) in Lutjanidae and Haemulidae (Perciformes) in the Indo-West Pacific. *International Journal for Parasitology*, 37: 1023-1045.  
<http://dx.doi.org/10.2478/s11686-010-0045-z>
- Muddula Krishna, N., Govindarao, V., & Venu, D. (2015). Taxonomic diversity, species composition, distribution, conservation and abundance of rocky intertidal fishes in the Visakhapatnam, East Coast of India. *Journal of Experimental Zoology India*, 19(1), 277-283.
- Nelson, J.S. (2006). *Fishes of the World*, 4th Edition., John Wiley & Sons, Inc., Mar 2006: 624 pp.
- Ormerod, S.J. (2003). Current issues with fish and fisheries: editor's overview and introduction. *Journal of Applied Ecology*, 40: 204-213.  
<http://dx.doi.org/1046/j.1365-2664.00824.x>
- Ovenden, J. R., & White, R.W.G. (1990). White Mitochondrial and Allozyme genetics of incipient speciation in a landlocked population of *Galaxias truttaceus* (Pisces: Galaxiidae). *Genetics*, 124: 701-716.  
<http://dx.doi.org/10.1038/hdy.1993.33>
- Ovenden, J.R., Lloyd, J.R., Keenan, S.J., & Slater, L.S. (2004). Pronounced genetic population structure in a potential vagile fish species (*Pristipomides multidens*, Teleostei; Perciformes; Lutjanidae) from East Indies triangle. *Molecular Ecology*, 13: 1991-1999.  
<http://dx.doi.org/10.1111/j.1365-294X.2004.02210.x>
- Ovenden, J.R., Lloyd, J., Newman, S.J., Keenan, C.P., & Slater, L.S. (2004). Spatial genetic subdivision between northern Australian and South East Asian populations of *Pristipomoides multidens*: a tropical marine reef fish species. *Fisheries Research* 59:57-69.  
<http://dx.doi.org/10.1186/1471-2148-9-40>
- Raymond, M.F., & Rousset, F. (1995a). An exact test for population differentiation. *Evolution*, 49: 1280-1283.  
<http://dx.doi.org/10.2307/241054>
- Ryman, N. (1991). Conservation genetics considerations in fishery management. *Journal of Fisheries Biology*, 39: 211-224.  
<http://dx.doi.org/1111/j.1095-8649.1991.tb05085.x>
- Ryman, N., & Utter, F. (1987). *Population Genetics and Fishery Management*. University of Washington Press, Washington, 420pp.  
<http://dx.doi.org/10.1017/S0025315400057180>
- Shaklee, J.B., & Tamaru, C.S. (1981). Biochemical and morphological evolution of Hawaiian bone fishes (*Albula*). *Systematic Zoology*, 30(2):125-146.  
<http://dx.doi.org/10.1093/SYSBIO/302.125>
- Shaklee, J.B., Allendorf, F.W., Morizot, D.C., & Whitt, G.S. (1990). Gene nomenclature for protein-coding loci in fish. *Transactions of the American Fisheries Society*, 119: 2-15.  
[http://dx.doi.org/10.1577/1548-8659\(1990\)119<0002:GNFPLI>2.3.CO;2](http://dx.doi.org/10.1577/1548-8659(1990)119<0002:GNFPLI>2.3.CO;2)
- Smith P.J., & Robertson, D.A. (1981). Genetic evidence for two species of sprat (*Sprattus*) in New Zealand waters.

- Marine Biology*, 62: 227-233.  
<http://dx.doi.org/10.1007/BF00397688>
- Smith, P.J., Francis, R.I.C.C., & McVeagh, M. (1991). Loss of genetic diversity due to fishing pressure. *Fisheries Research*, 10: 309-316.  
[http://dx.doi.org/10.1016/0165-7836\(91\)90082-Q](http://dx.doi.org/10.1016/0165-7836(91)90082-Q)
- Sulaiman, I.M., Fayer, R., Bern, C., Gilman, R.H., Trout, J.M., Schantz, P.M., Das, P., Lal, A.A., & Xiao, L. (2003). Triosephosphateisomerase gene characterization and potential zoonotic transmission of *Giardia duodenalis*. *Emerging Infectious Diseases journal*, 9:1444–1452.  
<http://dx.doi.org/10.3201/eid0911.030084>
- Utter, F.M. (1991). Biochemical genetics and fishery management: an historical perspective. *J Fish Biology*, 39: 1-20.  
<http://dx.doi.org/1111/j.1095-8649.1991.tb05063.x>
- Vasconcellos, A.V., Paulo Vianna, Paulo, C., Paiva, R. S., & Antonio, S. C. 2008. Genetic and morphometric differences between yellowtail snapper (*Ocyurus chrysurus*, lutjanidae) populations of the tropical West Atlantic. *Genetics and Molecular Biology*, 31(1): 308-316.  
<http://dx.doi.org/10.1590/S1415-7572008000200026>
- Ward, R.D., Elliott, N.G., & Grewe, P.M. (1995). Allozyme and mitochondrial DNA separation of Pacific Northern blue fin tuna, *Thunnus thynnus orientalis* (Temminck and Schlegel), from Southern blue fin tuna, *Thunnus maccoyii* (Castelnau). *Australian Journal of Marine and Fresh Water Research*, 46: 921-930.  
<http://dx.doi.org/10.1071/MF9950921>
- Whitmore, D.H. (1990). Electrophoretic and Isoelectric focusing techniques in fisheries management. CRC Press, Inc. (Florida), 23-80 pp.