



Genetic Diversity and Population Structure of the Asian Green Mussel (*Pernaviridis*) in the Waters of Sabah, Malaysia Based on Mitochondrial DNA D-Loop Sequences

Jen-Shi Lau¹, Julian Ransangan^{1,*}, Kenneth Francis Rodrigues²

¹ Fish Disease and Microbiology Laboratory, Borneo Marine Research Institute, Universiti Malaysia Sabah, 88400 Kota Kinabalu Sabah, Malaysia.

² Biotechnology Research Institute, Universiti Malaysia Sabah, 88400 Kota Kinabalu Sabah, Malaysia.

* Corresponding Author: Tel.: 320000-5599;
E-mail: kennethr@ums.edu.my

Received 24 January 2017
Accepted 18 May 2018

Abstract

The Asian green mussel (*Pernaviridis*) is a bivalve species with a high economic value. The waters surrounding the Malaysian state of Sabah hosts a wide range of geographically distributed populations of *P. viridis* whose population structure is unknown. This study was conducted to elucidate the genetic diversity and population structure of Asian green mussel from six sites located along the coastline of Sabah based on the mitochondrial DNA control region (D-loop). The complete D-loop sequences of 197 individuals were recovered by amplification using PCR, followed by DNA sequencing. Interpretation of the results revealed that the *P. viridis* populations exhibited high haplotype diversity ($H_d=0.912\pm 0.0002$) and low nucleotide diversity ($\pi=0.00890\pm 0.00066$). The AMOVA analysis showed Φ_{ST} was 0.02322 (P value < 0.05), which is indicative of low but significant structuring. Pairwise Φ_{ST} ranged from low to moderate indicating population differentiation. Five out of the fifteen population pairs showed significant population differentiation. Both AMOVA and pairwise Φ_{ST} suggest there were significant population subdivisions in Asian green mussel population in Sabah. Although the Asian green mussel population in Sabah has gone through a severe mass mortality, evidence for genetic bottleneck was not detected. However, a smaller population size after a severe demographic reduction can cause the population become vulnerable. Thus, regular monitoring of Asian green mussel population is required to keep track of fluctuations in population size and composition. This study provided new population genetic information which is crucial for establishing fisheries management strategies for this species.

Keywords: *Pernaviridis*, D-loop, control region, mitochondrial DNA, genetic diversity, high haplotype diversity, low nucleotide diversity, population structure, demographic history, Sabah, Malaysia.

Introduction

The Asian green mussel (*Pernaviridis*) is a type of marine bivalve mollusk under the family Mytilidae. The genus *Perna* consists of three extant species, *Pernaperma*, *Pernaviridis* and *Perna canaliculus* (Siddall, 1980). Some researchers have also included *Pernaindica* and *Pernapicta* under genus *Perna*. However, both of them have been synonymized with *Pernaperma* (Siddall, 1980; Vakily, 1989; Wood, Apte, Macvoy & Gardner, 2007).

The Asian green mussel has two hinged shells, which are connected with a posterior adductor muscle (Gosling, 2015). The native habitat of the Asian green mussel is in the Indo-Pacific region, which encompasses regions between Japan to New Guinea and from Persian Gulf to South Pacific Islands (FAO, 2013). Asian green mussel generally inhabits marine intertidal, subtidal and estuarine environments, which have high salinity and receive more nutrients from land run-off (Rajagopal, Venugopalan, van der Velde,

& Jenner 2006). Asian green mussel is able to tolerate a wide range of salinities and temperatures (Sivalingam, 1977). It is a dioecious organism which means the male and female reproductive organs are in separate individuals. *P. viridis* employs external fertilization whereby males and females Asian green mussels release gametes directly into the water during spawning. The Asian green mussel adult is sessile throughout its life whereas the larvae of Asian green mussel have long pelagic duration, which last around 21 to 35 days after fertilization (Laxmilatha *et al.*, 2011).

Asian green mussel is a popular type of seafood, especially in China, Philippines and Malaysia (FAO, 2013). Asian green mussel farming is considered one of the potential shellfish aquaculture in Malaysia. An experimental culture of Asian green mussel was initiated by the Fisheries Research Institute of Malaysia in Penang in 1977 (Mazuki, 1998). The aquaculture of this species has advanced since its first introduction; however, the production of Asian green

mussel in Malaysia lags behind as compared to countries such as Thailand and Philippines (FAO, 2013). There is a need to formulate fisheries management strategies for improving the aquaculture of the Asian green mussel aquaculture, but this is limited by the lack of knowledge with reference to the genetic diversity of the species. Therefore, there is a need to understand the population genetic of Asian green mussel in Sabah.

Genetic diversity and population structure of aquatic animals can be examined using a variety of molecular markers. The characteristics of mtDNA which include maternal inheritance and the absence of genetic recombination makes it the most popular molecular marker (Galtier, Nabholz, Glémin, & Hurst, 2009). Mitochondrial DNA is a super-coiled circular double stranded DNA located outside the cell nucleus (Freeland, Kirk, & Petersen, 2011). The control region of mtDNA, which also is known as D-loop region has the highest mutation rate (Wan, Wu, Fujihara, & Fang, 2004). This characteristic makes D-loop region a good marker for investigating genetic diversity in closely related species or within a species (Rosel, Haygood, & Perrin, 1995).

Several population genetic studies have been conducted on Asian green mussel populations in Thailand, India, Malaysia, China and Singapore (Yap, Tan, Ismail, & Omar, 2002; 2004; Ong, Yusoff, Yap, & Tan, 2009; Divya, Thomas, Gopalakrishnan, Sathianandan, & Paulton, 2012; Lin, Loong, & Gen, 2012; Yap, Cheng, Ong, & Tan, 2013; Gilg, Johnson, Gobin, Bright, & Ortolaza, 2013; Ibrahim, 2014; Ye, Li, & Wu, 2015). As far as Malaysian Borneo is concerned, a preliminary study was conducted in Santubong, Sarawak (Ibrahim, 2014) but no studies have been reported from Sabah. This investigation was directed towards elucidation of the genetic diversity of the population along the coastal waters of Sabah.

Materials and Methods

Sample Collection

A total of 197 Asian green mussels (*P. viridis*) were collected from six locations in the coastal waters of Sabah (Figure 1). Detailed information regarding the sampling location has been summarized in Table 1. The adductor muscle of each specimen was kept at -20°C prior to DNA extraction.

DNA Extraction

Doubly Uniparental Inheritance (DUI) had been discovered in several mollusc species. The F-type mtDNA exists in the somatic tissue whereas the M-type mtDNA is present mainly in the males' gonadal tissue and sperm (Chiesa et al., 2011; Plazzi & Passamonti, 2010). A study conducted by Wei, Kong, Wu, and Yu (2009) reported on the

absence of doubly uniparental inheritance in *P. viridis* and based on this report DNA was only extracted from the adductor muscle to avoid the possibility of M-type mtDNA extraction. Total genomic DNA was extracted from the Asian green mussel adductor muscle with DNeasy DNA extraction kit (Qiagen) according to manufacturer's instructions. Concentration and purity of DNA were verified using agarose gel electrophoresis and DNA Spectrophotometry (Pharmacia GeneQuant pro UV-Visible Spectrophotometer). The genomic DNA were diluted to 25ng/μL and stored at -20°C in 100μL AE buffer (Qiagen).

PCR Amplification and DNA Sequencing

The D-loop region of the mitochondrial DNA was amplified using forward primer 5'-GGGAGGCTATGGTGA GTCAA-3' reverse primer 5'-TGCCACATAAACTACCCTCATC-3'. The primers were designed using Primer3 (Koressaar & Remm, 2007; Untergrasser et al., 2012). Polymerase chain reactions (PCR) were carried out in volumes of 25 μL with 1x GoTaq Flexi Buffer (Promega), 0.2mM of each dNTP (Promega), 1.5mM of MgCl₂ (Promega), 0.1 μM of each primer, 1U of Taq DNA polymerase (Promega) and 50ng/μL of DNA. PCR was carried out using the Applied Biosystems Veriti™ thermal cycler under the following conditions: one cycle of 95°C for 3min, followed by 30 cycles at 95°C for 30s, 60°C for 30s and 72°C for 30s and with a final extension for 5 min at 72°C. PCR products were purified with PCRquick-spin™ PCR Product Purification Kit (iNtRON) according to manufacturer's instructions. Each of the PCR products was cloned into pGEM®-T Easy cloning vector (Promega). Bidirectional DNA sequencing was conducted by AIT Biotech Pt. Ltd (Singapore). The sequencing reactions for the DNA samples were carried out with BigDye Terminator v3.1 Cycle Sequencing kit and resolved using ABI 3730XL DNA analyzers (Applied Biosystems) sequencer.

Statistical Analysis

Forward and reverse D-loop sequences were assembled and further checked manually using the SeqMan software (DNASTAR package version 7.1.0, 2006). VecScreen (Altschul et al., 1997) in the National Center for Biotechnology Information (NCBI) <http://www.ncbi.nlm.nih.gov/> was used to eliminate vector contamination sites from the sequences. The flanking region encoding a fragment of the COI gene (136 base pair) and NADH dehydrogenase subunit 4 gene (35 base pair) were edited out from the consensus sequences. The DNA sequences were then aligned using the ClustalW in MEGA 6.0 (Tamura, Stecher, Peterson, Filipowski, &




Figure 1. Map of Sabah, Malaysia with the six sampling locations (black triangles).

Table 1. Sampling details, genetic diversity indices and neutral tests of green mussel (*P. viridis*) populations in Sabah

Geographic Location	Collection Date	<i>N</i>	<i>n</i>	<i>n_p</i>	<i>H_d</i> (mean ± SD)	π (mean ± SD)	Tajima's <i>D</i> (P-value)	Fu's <i>F_s</i> (P-value)
Kota Belud	Sept 2015	35	15	10	0.881±0.002	0.00766±0.00156	-	-2.565(0.151)
Kota Kinabalu	Oct 2013	30	22	17	0.966±0.020	0.01216±0.00126	0.916(0.194)	-5.786(0.062)
Kota Marudu	Sept 2013	32	21	16	0.919±0.040	0.01096±0.00156	0.861(0.203)	-3.821(0.096)
Kuala Penyu	Sept 2014	34	12	8	0.779±0.064	0.00602±0.00166	1.123(0.113)	-1.522(0.264)
Tawau	Jun 2014	35	21	16	0.919±0.034	0.00751±0.00131	1.411(0.057)	-5.821(0.053)
Tuaran	April 2013	31	19	14	0.920±0.034	0.00839±0.00169	1.318(0.074)	-4.948(0.078)
Total population	-	197	88	-	0.912±0.0002	0.00890±0.00066	1.214(0.105)	-4.077(0.092)
							1.141(0.124)	

Note: *N*, sample size; *n*, haplotype number; *H_d*, haplotype diversity and π , nucleotide diversity.

Kumar, 2013). The general genetic indices of mtDNA, including the haplotype number (*H*), haplotype diversity (*H_d*) and nucleotide diversity (π) were calculated using the DnaSP5.10.1 (Librado&Rozas, 2009).

The neighbor joining phylogenetic tree, maximum likelihood phylogenetic tree and maximum parsimony phylogenetic trees were constructed using the MEGA 6.0 to depict the genetic relationships among haplotypes. The statistical robustness of the nodes for all three phylogenetic trees was determined using bootstrap analysis with 1000 replicates. In addition, a haplotype network was created to depict the phylogenetic and geographical relationships of haplotypes using the median-joining method in PopART (Bandelt, Forster, & Röhl, 1999).

Pairwise Φ_{ST} , analysis of molecular variance (AMOVA) and Mantel test were used to examine

Asian green mussel population structure using the Arlequin 3.5.2.1. The analysis of molecular variance (AMOVA) was conducted to quantify the partitioning of genetic variation present within population and among populations, and its significance was determined with 1000 permutations. Pairwise Φ_{ST} , an analog of F-statistic was also calculated to determine the genetic differentiation between sampling locations and the significance of pairwise Φ_{ST} values was determined with 100 permutations. Besides, Mantel test (Mantel, 1967) was used to determine if there was any significant correlation between pairwise Φ_{ST} and geographic distance. Geographic distance was referred to the shortest distance between two locations along the coast. Significance of the Mantel test was determined with 10000 permutations.

Demographic history of Asian green mussel populations was examined using the neutral tests

including Tajima's D (Tajima, 1989) and Fu's F_s (Fu, 1997) incorporated in the Arlequin (Version 3.5.2.1). Both Tajima's D and Fu's F_s were used to determine whether the variation in DNA sequences deviate from the neutral model (Hamilton, 2009). Significance of the Tajima's D and Fu's F_s was determined with 1000 permutations.

Results

Sequence Variation and Genetic Diversity

The length of the D-loop fragment amplified of the mtDNA genome in the Asian green mussel in Sabah ranged from 657bp to 665bp. The actual size of the D-loop DNA fragment varied due to the presence of insertions and deletion (indels). All the DNA sequences were deposited in the EMBL Nucleotide Sequence Database under accession numbers KP731638-KP731799 and KU555309-KU555343. The average base composition was as follows: $T=30.4\%$, $C=6.6\%$, $A=33.8\%$ and $G=29.2\%$. The A/T base contents of the D-loop region were higher than the C/G base contents. This result was concordant with previous studies, which reported that the D-loop is an A-T rich region of the mitochondrial genome (Brown, Gadaleta, Pepe, Saccone, & Sbisà, 1986). In addition, 107 polymorphic sites were observed, including 73 singleton variable sites and 34 parsimony informative sites. Genetic diversity indices including haplotype number (n), haplotype diversity (H_d) and nucleotide diversity (π) are presented in Table 1. A total of 88 haplotypes were identified in the 197 samples. The number of haplotypes ranged from 15 in Kota Belud to 22 in Kota Kinabalu. The mean haplotype diversity (H_d) and nucleotide diversity (π) of the total population were 0.912 ± 0.0002 and 0.00890 ± 0.00066 respectively. The haplotype diversities (H_d) were very high, ranging from 0.779 ± 0.064 in Kuala Penyu to 0.966 ± 0.020 in Kota Kinabalu. On the other hand, the nucleotide diversity (π) is very low, ranging from 0.00602 ± 0.00166 in Kuala Penyu to 0.01216 ± 0.00126 in Kota Kinabalu.

Phylogenetic and Network Analysis

The topologies produced from neighbor joining phylogenetic tree (Figure 2), maximum likelihood phylogenetic tree (Figure 3) and maximum parsimony phylogenetic tree (Figure 4) were similar. All three phylogenetic trees were divided into two haplogroups. One haplogroup contained most of the haplotypes and the second haplogroup contained another 18 haplotypes. The median joining network was used to further depict the phylogenetic and geographical relationships among haplotypes (Figure 3). Resultant network exhibited star-like pattern surrounding haplotype 2, haplotype 3 and haplotype 4. Haplotype 4 was considered the ancestral haplotype, which was the most common haplotype (53.4%) encompassing

all six locations. Haplotype 3 (27.2%) encompassed all six locations whereas haplotype 2 (28.4%) encompassed all locations except Kota Kinabalu. The phylogenetic trees and median joining network provided some insight into the relationship among haplotypes of Asian green mussel in Sabah. Nevertheless, a clear spatial pattern was not able to be identified from phylogenetic trees or median joining network.

Population Structure Analysis

The AMOVA analysis based on haplotype frequencies revealed that 97.68% of the genetic variation occurred within populations while only 2.32% of the genetic variation occurred among populations (Table 2). The Φ_{ST} was 0.02322 ($P < 0.05$), which indicate low but significant population structuring of the *P. viridis* populations in Sabah.

Population pairwise Φ_{ST} showed that the Φ_{ST} between Kuala Penyu and Tuaran was the highest 0.07048 whereas the Φ_{ST} between Kota Marudu and Tawau was the lowest, at -0.01311 (Table 3). Mantel test with correlation coefficient, $r = -0.37$, ($P > 0.05$) showed there was no correlation between geographic distance and population pairwise Φ_{ST} between Asian green mussel populations.

Neutral Tests

The results of Tajima's D and Fu's F_s neutral tests are presented in Table 1. Both Tajima's D and Fu's F_s test showed consistent non significant negative D-value across populations, which indicate that, the demographic history of the D-loop region of *P. viridis* populations was in agreement with neutral expectation.

Discussion

Despite high haplotype diversity ($H_d = 0.912 \pm 0.0002$) of the Asian green mussel populations in Sabah, its nucleotide diversity ($\pi = 0.00890 \pm 0.00066$) was very low. High haplotype diversity but low nucleotide diversity of Asian green mussel population in Sabah could be attributed to the high mutational rate of the D-loop region (Wan, Wu, Fujihara, & Fang, 2004). The haplotype diversity of a relatively rapid evolving genome within a population often approach 1.0 as many individuals will tend to have unique haplotypes (Freeland, Kirk, & Petersen, 2011). Besides, low nucleotide diversity but high haplotype diversity may also be the indication of genetic bottleneck. Both haplotype and nucleotide diversities can be diminished in the event of genetic bottleneck (Frankham, Ballou, & Briscoe, 2004; Hamilton, 2009). High haplotype diversity and low nucleotide diversity can be observed in a population experiences rapid expansion from a low effective population size, assuming there is sufficient time for

haplotype to increase through mutation but insufficient time for accumulation of large sequence differences.(Grant & Bowen, 1998; Lowe, Harris, & Ashton, 2004).A rapid population growth can enhance the retention of new mutations in the population(Grant & Bowen, 1998). High haplotype diversity and low nucleotide diversity has been reported in many marine organisms such as *Cynoscionacoupa* (Rodrigues *et al.*, 2008), *Architeuthis dux* (Winkelmann *et al.*, 2013) and *Girellapunctata*(Saito *et al.*, 2008)that have

undergone a severe demographic reduction.

The population pairwise Φ_{ST} showed low to moderate population differentiation which ranged from -0.01311 to 0.07048. According to Wright (1978), fixation index, Φ_{ST} should range around 0.0 to 1.0. However, pairwise Φ_{ST} was negative between Tawau and Kota Marudu. Negative Φ_{ST} occurs when the genetic differentiation within population is higher than between populations, which indicate that gene flow between the populations is very high(Bortolotto,Bucklin, Mezzavilla, Zane,




Figure 2. Neighbour-joining tree based on D-loop haplotypes of the green mussel (*P. viridis*). (Hap: Haplotype).




Figure 3. Median-joining network based on the D-loop haplotypes of the *P. viridis*. Circles represent different haplotypes with relative size proportionate to its observed frequency. The colors of the circles indicate the geographic region. The number labels represent haplotypes names. The hatch marks indicate the mutation steps between haplotypes and the black circles represent missing haplotypes.

Table 2. Analysis of molecular variance (AMOVA) for green mussel populations in Sabah based on D-loop region

Source of variation	d.f	Sum of squares	Variance components	Percentage of variation	F_{ST}	P value
Among populations	5	3.980	0.01063	2.32	0.02322	0.0098
Within populations	191	85.416	0.44720	97.68		
Total	196	89.396	0.45783			

Table 3. Population pairwise F_{ST} between green mussel populations in Sabah based on D-loop region

	KB	KK	KM	KP	TA	TU
KB	-					
KK	0.03591*	-				
KM	0.01994	0.00904	-			
KP	0.01931	0.06527*	0.01559	-		
TA	0.01121	0.00661	-0.01311	0.01695	-	
TU	0.01113	0.02563*	0.03550*	0.07048*	0.02437	-

*Significant values at $P < 0.05$; Abbreviation: KB=Kota Belud; KK=Kota Kinabalu; KM=Kota Marudu; KP=Kuala Penyu; TA=Tawau; TU=Tuaran.

&Patarnello, 2011; Roesti, Salzburger, & Berner, 2012). Most of the genetic variation of marine invertebrates is observed within rather than between populations (Burton, 1997). This is likely to be the result of the spawning behaviour of adult invertebrates and high dispersal ability of marine invertebrate larvae. The larvae of this species have very long pelagic duration, which is around 21 to 35 days (Laxmilatha *et al.*, 2011). It is likely that the larvae of Asian green mussel tend to drift away from their natal habitat and that this in turn increases the gene flow between populations and lowers the degree of differentiation between populations. This study detected significant genetic differentiation between KK-KB, KK-KP, KK-TU, KM-TU and KP-TU. The significant population differentiation could be attributed to oceanographic features or the current patterns that limit the gene flow between two sites. Oceanographic features such as oceanographic front can prevent two adjacent sites to exchange migrants (Gilg & Hilbish, 2003). The surface current around Sabah is remarkably complex. The dominant surface circulation is the Mindoro strait outflow which drives the sea current from South China Sea to Sulu Sea through Mindoro strait. The Mindoro strait flow outwards to the South China Sea through the Balabac Strait and out to Sulu Sea through the Sibutu Passage (Han *et al.*, 2009). Nevertheless, we are not able to pinpoint the real reasons for the population subdivision in this study. Further studies will require identifying specific causative factors. The significant AMOVA and population differentiation indicated there was a significant subdivision among populations of Asian green mussels in Sabah. However, a clear spatial pattern could not be discerned. The significant subdivision among populations should be taken into consideration in the development and implementation of Asian green mussel aquaculture management strategies.

It is important to determine if the populations experienced bottleneck, as this would likely lead to inbreeding depression, loss of genetic variation and deleterious allele fixation, any of which can ultimately compromise species' adaptive potential and the probability of population persistence (Frankham, Ballou, & Briscoe, 2004). The Asian green mussel populations in Sabah have been reported to have gone through a severe mass mortality from 2010 (Tan

& Ransangan, 2015; Taib, Madin, & Ransangan, 2016). Studies carried out by researchers from several institutions of higher learning and fisheries department in Malaysia have not been able to determine the cause of the mass mortality. Although the green mussel experienced severe mass mortality for a long period of time, this study did not detect the evidence of genetic bottleneck in Asian green mussel populations. Under certain circumstances, a severe demographic bottleneck may not necessarily present itself as evidence in terms of a genetic signature. Several reasons could lead to this phenomenon. Firstly, effective population size that is large enough can diminish the bottleneck signature. Sastre *et al.* (2011) reported the mitochondrial DNA data can effectively detect bottleneck in the small isolated population but not in a large one. Secondly, small number of migrants can also diminish the effects of genetic bottleneck (Keller *et al.*, 2001; Busch, Waser, & DeWoody, 2007). Despite the absence of evidence of a genetic bottleneck, the smaller population size after severe demographic reduction could cause Asian green mussel populations to become more sensitive to environmental perturbations and stochastic demographic processes. Thus, regular monitoring of Asian green mussel population is recommended in order to keep track of fluctuations in population size and composition.

Acknowledgements

This study was funded by a research grant (COE0009) from the Ministry of Education, Malaysia. Our appreciation also goes to the Fisheries Department of Sabah for granting the permission for sample collection along the coastal waters of Sabah.

Conflict of Interest

The authors declare that they do not have any conflicting interests.

References

- Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W., & Lipman, D.J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research*,

- 25(17), 3389-3402. <https://dx.doi.org/10.1093/nar/25.17.3389>
- Bandelt, H., Forster, P., & Röhl, A. (1999). Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution*, 16(1), 37-48. <https://doi.org/10.1093/oxfordjournals.molbev.a026036>
- Brown, G.G., Gadaleta, G., Pepe, G., Saccone, C., & Sbisà, E. (1986). Structural conservation and variation in the D-loop-containing region of vertebrate mitochondrial DNA. *Journal of Molecular Biology*, 192(3), 503-511. [https://dx.doi.org/10.1016/0022-2836\(86\)90272-X](https://dx.doi.org/10.1016/0022-2836(86)90272-X)
- Bortolotto, E., Bucklin, A., Mezzavilla, M., Zane, L., & Patarnello, T. (2011). Gone with the currents: lack of genetic differentiation at the circum-continental scale in the Antarctic krill *Euphausiasuperba*. *BMC Genetics*, 12, 32-50. <https://dx.doi.org/10.1186/1471-2156-12-32>
- Burton, R.S. (1997). Genetic evidence for long term persistence of marine invertebrate populations in an ephemeral environment. *Evolution*, 51(3), 993-998. <https://dx.doi.org/10.2307/2411174>
- Busch, J.D., Waser, P.M., & DeWoody, J.A. (2007). Recent demographic bottlenecks are not accompanied by a genetic signature in banner-tailed kangaroo rats (*Dipodomys spectabilis*). *Molecular Ecology*, 16(12), 2450-2462. <https://dx.doi.org/10.1111/j.1365-294X.2007.03283.x>
- Chiesa, S., Marzano, F.N., Minervini, G., Lucrezia, D.D., Baccarani, G., Bordignon, G., & Argese, E. (2011). The invasive Manila clam *Ruditapes philippinarum* (Adams and Reeve, 1850) in Northern Adriatic Sea: population genetics assessed by an integrated molecular approach. *Fisheries Research*, 110(2), 259-267. <https://dx.doi.org/10.1016/j.fishres.2011.04.013>
- Divya, P.R., Thomas, P.C., Gopalakrishnan, A., Sathianandan, T.V., & Paulton, M.P. (2012). Stock structuring in Asian green mussel *Pernaviridis* population along the Indian coast based on shell morphometrics and RAPD markers. *Indian Journal of Animal Sciences*, 82(7), 775-778. <http://eprints.cmfri.org.in/id/eprint/9916>
- DNA STAR package version 7.1.0. (2006). Lasergene, DNA Star, Inc. Madison, WI, USA.
- FAO. (2013). Species fact sheets: *Pernaviridis* (Linnaeus, 1758). Retrieved from <http://www.fao.org/fishery/species/2691/en>
- Frankham, R., Ballou, J.D., & Briscoe, D.A. (2004). A primer of conservation genetics. United Kingdom, Cambridge University Press, 220 pp.
- Freeland, J.R., Kirk, H., & Petersen, S.D. (2011). Molecular ecology, second edition. Oxford, United Kingdom, John Wiley & Sons, Ltd., 400 pp.
- Fu, Y.X. (1997). Statistical tests for neutrality of mutations against population growth, hitchhiking and background selection. *Genetics*, 147(2), 915-925. <http://www.genetics.org/content/genetics/147/2/915.full.pdf>
- Galtier, N., Nabholz, B., Glémin, S., & Hurst, G.D.D. (2009). Mitochondrial DNA as a marker of molecular diversity: a reappraisal. *Molecular Ecology*, 18(22), 4541-4550. <https://doi.org/10.1111/j.1365-294x.2009.04380.x>
- Gilg, M.R. & Hilbish, T.J. (2003). The geography of marine larval dispersal: coupling with fine-scale physical oceanography. *Ecology*, 84(11), 2989-2998. <https://doi.org/10.1890/02-0498>
- Gilg, M.R., Johnson, E.G., Gobin, J., Bright, B.M., & Ortolaza, A.I. (2013). Population genetics of introduced and native populations of the Asian green mussel, *Pernaviridis*: determining patterns of introduction. *Biological Invasions*, 15(2), 459-472. <https://doi.org/10.1007/s10530-012-0301-2>
- Gosling, E. (2015). Marine bivalve molluscs, second edition. Oxford, United Kingdom, John Wiley & Sons, Ltd., 512 pp.
- Grant, W.S. & Bowen, B.W. 1998. Shallow population histories in deep evolutionary lineages of marine fishes: insights from sardines and anchovies and lessons for conservation. *The American Genetic Association*. 89: 415-426. <https://doi.org/10.1093/jhered/89.5.415>
- Hamilton, M.B. (2009). Population genetics. United Kingdom Oxford, United Kingdom, John Wiley & Sons, Ltd., 424 pp.
- Han, W.Q., Moore, A.M., Levin, J. Zhang, B., Arango, H.G., Curchitser, E., Lorenzo, E.D., Gordon, A.L. and Lin, J.L. (2009). Seasonal surface ocean circulation and dynamics in the Philippine Archipelago region during 2004-2008. *Dynamics of Atmospheres and Oceans*. 47. 114-137. <http://dx.doi.org/10.1016/j.dyatmoce.2008.10.007>
- Ibrahim, S.N. (2014). Preliminary study of morphological and genetic diversity on Asian green mussel (*Pernaviridis*) in Santubong, Sarawak (BSc thesis). Universiti Malaysia Sarawak, Sarawak, Malaysia. <http://ir.unimas.my/id/eprint/8276>
- Keller, L.F., Jeffery, K.J., Arcese, P., Beaumont, M.A., Hochachka, W.M., Smith, J.N.M., & Bruford, M.W. (2001). Immigration and the ephemerality of a natural population bottleneck: evidence from molecular markers. *Proceedings of the Royal Society B*, 268(1474), 1387-1394. <https://doi.org/10.1098/rspb.2001.1607>
- Koressaar, T., & Remm, M. (2007). Enhancements and modifications of primer design program Primer3. *Bioinformatics*, 23(10), 1289-1291. <https://doi.org/10.1093/bioinformatics/btm091>
- Laxmilatha, P., Rao, G.S., Patnaik, P., Nageshwara Rao, T., Prasad Rao, M., & Dash, B. (2011). Potential for the hatchery production of spat of the Asian green mussel *Pernaviridis* Linnaeus (1758). *Aquaculture*, 312(1-4), 88-94. <https://doi.org/10.1016/j.aquaculture.2010.12.031>
- Librado, P., & Rozas, J. (2009). DnaSPv5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25(11), 1451-1452. <https://doi.org/10.1093/bioinformatics/btp187>
- Lin, G., Loong, C.L., & Gen, H.Y. (2012). Genetic variations in populations from farms and natural habitats of Asian green mussel, *Pernaviridis*, in Singapore inferred from nine microsatellite markers. *Journal of the World Aquaculture Society*, 43(2), 270-277. <https://doi.org/10.1111/j.1749-7345.2012.00558.x>
- Lowe, A., Harris, S., & Ashton, P. (2004). Ecological genetics: design, analysis, and application. Oxford, United Kingdom, Blackwell Science Ltd., 432 pp.
- Mantel, N. (1967). The detection of disease clustering and a generalized regression approach. *Cancer Research*, 27(2), 209-220. http://cancerres.aacrjournals.org/content/27/2_Part_1/209.full-text.pdf
- Ong, C.C., Yusoff, K., Yap, C.K., & Tan S.G. (2009). Genetic characterization of *Pernaviridis* in peninsular

- Malaysia using microsatellite markers. *Journal of Genetics*, 88(2), 153-163. <https://doi.org/10.1007/s12041-009-0023-0>
- Plazzi, F., & Passamonti, M. (2010). Towards a molecular phylogeny of mollusks: bivalves' early evolution as revealed by mitochondrial genes. *Molecular Phylogenetics and Evolution*, 57(2), 641-657. <https://doi.org/10.1016/j.ympev.2010.08.032>
- Rajagopal, S., Venugopalan, V.P., van der Velde, G., & Jenner, H.A. (2006). Greening of the coasts: a review of the *Pernaviridis* success story. *Aquatic Ecology*, 40(3), 273-297. <https://doi.org/10.1007/s10452-006-9032-8>
- Rodrigues, R., Schneider, H., Santos, S., Vallinoto, M., Sain-Paul, U., & Sampaio, I. (2008). Low levels of genetic diversity depicted from mitochondrial DNA sequences in a heavily exploited marine fish (*Cynoscionacoupa*, Sciaenidae) from the northern coast of Brazil. *Genetics and Molecular Biology*, 31(2), 487-492. <https://doi.org/10.1590/s1415-47572008000300015>
- Roesti, M., Salzburger, W., & Berner, D. (2012). Uninformative polymorphisms bias genome scans for signatures of selection. *BMC Evolutionary Biology*, 12: 94-101. <https://doi.org/10.1186/1471-2148-12-94>
- Rosel, P.E., Haygood, M.G., & Perrin, W.F. (1995). Phylogenetic relationship among the true porpoises (Cetacea: Phocoenidae). *Molecular Phylogenetics and Evolution*, 4(4), 463-474. <https://doi.org/10.1006/mpev.1995.1043>
- Saito, T., Washio, S., Dairiki, K., Shimoio, S., Itoi, S., & Sugita, H. (2008). High gene flow in *Girellapunctata* (Perciformes, Kyphosidae) among the Japanese Islands inferred from partial sequence of the control region in mitochondrial DNA. *Journal of Fish Biology*, 73(8), 1937-1945. <https://doi.org/10.1111/j.1095-8649.2008.01997.x>
- Sastre, N., Vilà, C., Salinas, M., Bologov, V.V., Urios, V., Sánchez, A., & Ramirez, O. 2011. Signatures of demographic bottlenecks in European wolf populations. *Conservation Genetics*, 12(3), 701-712. <https://doi.org/10.1007/s10592-010-0177-6>
- Siddall, S.E. 1980. A clarification of the genus *Perna* (Mytilidae). *Bulletin of Marine Science*, 30(4), 858-870. <http://www.ingentaconnect.com/content/umrmsmas/bullmar/1980/00000030/00000004/art00008>
- Sivalingam, P.M. (1977). Aquaculture of the Asian green mussel, *Mytilusedulis* Linnaeus, in Malaysia. *Aquaculture*, 11(4), 297-312. [https://doi.org/10.1016/0044-8486\(77\)90079-5](https://doi.org/10.1016/0044-8486(77)90079-5)
- Taib, A.M., Madin, J., & Ransangan, J. (2016). Density, recruitment and growth performance of Asian green mussel (*Pernaviridis*) in Marudu Bay, Northeast Malaysian Borneo, three years after a massive mortality event. *Songklanakarinn Journal of Science and Technology*. <http://rdo.psu.ac.th/sjstweb/journal/38-6/38-6-5.pdf>
- Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, 123(3), 585-595. <http://www.genetics.org/content/genetics/123/3/585.full.pdf>
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., & Kumar, S. (2013). MEGA 6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Bioev and Evolution*, 30(12), 2725-2729. <https://doi.org/10.1093/molbev/mst197>
- Tan, K.S., & Ransangan, J. (2015). Factors influence the larvae distribution and spat settlement of *Pernaviridis* in Marudu Bay, Sabah, Malaysia. *Advances in Environmental Bioev*, 9(17), 18-23. [http://www.aensiweb.net/AENSIWEB/aeb/aeb/2015/Special%20IPN%20Langkawi%20\(August\)/18-23.pdf](http://www.aensiweb.net/AENSIWEB/aeb/aeb/2015/Special%20IPN%20Langkawi%20(August)/18-23.pdf)
- Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B.C., Remm, M., & Rozen, S.G. (2012). Primer3—new capabilities and interfaces. *Nucleic Acids Research*, 40(15), 115. <https://doi.org/10.1093/nar/gks596>
- Vakily, J.M. 1989. The biology and culture of mussels of the genus *Perna*. Philippines: International Center for Living Aquatic Resources Management. http://pdf.usaid.gov/pdf_docs/PNABF179.pdf
- Wan, Q.H., Wu, H., Fujihara, T., & Fang S.G. (2004). Which genetic marker for which conservation genetics issue? *Electrophoresis*, 25(14), 2165-2176. <https://doi.org/10.1002/elps.200305922>
- Wei, Z., Kong, X., Wu, X., & Yu, Z. (2009). Genetic analysis of the marine mussel *Pernaviridis* (Bivalvia: Mytilidae) based on mitochondrial COI sequence and its phylogenetic relationship with other congeneric species. *Journal of Tropical Oceanography*, 28, 72-78. <http://dx.doi.org/10.11978/j.issn.1009-5470.2009.06.072>
- Winkelmann, I., Campos, P.F., Strugnell, J. Chereil, Y., Smith, P.J., Kubodera, T., & Gilbert, M.T.P. (2013). Mitochondrial genome diversity and population structure of the giant squid *Architeuthis*: genetics sheds new light on one of the most enigmatic marine species. *Proceedings of the Royal Society B*, 280(1759), 203-213. <https://doi.org/10.1098/rspb.2013.0273>
- Wood, A.R., Apte, S., MacAvoy, E.S., & Gardner, J.P. (2007). A molecular phylogeny of the marine mussel genus *Perna* (Bivalvia: Mytilidae) based on nuclear (ITS1&2) and mitochondrial (COI) DNA sequences. *Molecular Phylogenetics Evolution*, 44(2): 685-698. <https://dx.doi.org/10.1016/j.ympev.2006.12.019>
- Wright, S. (1978). Variability within and among natural populations, volume 4. Chicago, U.S., University of Chicago Press, 590 pp.
- Yap, C.K., Tan, S.G., Ismail, A., & Omar, H. (2002). Genetic variation of the green-lipped mussel *Pernaviridis* (L.) (Mytilidae: Mytiloida: Mytilicae) from the west coast of Peninsular Malaysia. *Zoological Studies*, 41(4), 376-387. <http://zoolstud.sinica.edu.tw/Journals/41.4/376.pdf>
- Yap, C.K., Tan, S.G., Ismail, A., & Omar, H. (2004). Allozyme polymorphisms and heavy metal levels in the green-lipped mussel *Pernaviridis* (Linnaeus) collected from contaminated and uncontaminated sites in Malaysia. *Environment International*, 30(1), 39-46. [https://doi.org/10.1016/s0160-4120\(03\)00144-2](https://doi.org/10.1016/s0160-4120(03)00144-2)
- Yap, C.K., Cheng, W.H., Ong, C.C., & Tan, S.G. (2013). Heavy metal contamination and physical barrier are main causal agents for the genetic differentiation of *Pernaviridis* populations in Peninsular Malaysia. *Sains Malaysiana*, 42(11), 1557-1564. http://journalarticle.ukm.my/6618/1/03_C.K._Yap.pdf
- Ye, Y.Y., Li, J.J., & Wu, C.W. (2015). Genetic diversity and

population connectivity of the Asian green mussel *Perna viridis* in South China Sea, inferred from mitochondrial DNA markers. *Biochemical*

Systematics and Ecology, 61, 470-475.
<https://doi.org/10.1016/j.bse.2015.07.009>