



Effects of Three Different Dietary Binders on Juvenile Sea Cucumber, *Apostichopus japonicus*

Seonghun Won¹, Ali Hamidoghli¹, Jin-hyeok Lee¹, Jinho Bae¹ and Sungchul C. Bai^{1*}

¹ Pukyong National University, Department of Marine Bio-Materials & Aquaculture/Feeds & Foods Nutrition Research Center, Busan, 608-737 Korea

Tel.: 82 51 6206137/6874. Fax: 82 51 6286873

E-mail: scbai@pknu.ac.kr

Abstract

This study was conducted to evaluate the effects of three different dietary binders on growth performance and water analysis in the culture of juvenile sea cucumber, *Apostichopus japonicus*. The basal diet without binder was used as control (CON), three diets with each of three different binder sources such as guar gum (G₁₀), carrageenan (C₁₀) and xanthan gum (X₁₀) at 10%, and other three diets (G₅C₅, C₅X₅ and X₅G₅) were formulated to contain mixture of two different binders at 5% for each. Juvenile sea cucumber averaging 1.09 ± 0.01 g (mean ± SD) were fed one of the seven experimental diets for 12 weeks. Weight gain and specific growth rate of sea cucumber fed G₁₀ and X₅C₅ diets were significantly higher than those of sea cucumber fed the CON diet. Diet cumulative melt away rates for G₁₀, X₁₀, X₅C₅, G₅C₅ and G₅X₅ diets were significantly higher than CON diet. G₁₀ diet resulted in a significantly lower chemical oxygen demand comparing to CON diet. Therefore, the present study demonstrated that 10% of dietary guar gum and mixture of 5% xanthan gum and 5% carrageenan could improve growth and water quality in juvenile sea cucumber aquaculture.

Key words: *Apostichopus japonicus*, binder, growth performance, water quality

Introduction

Sea cucumbers with high commercial importance are distributed all around the world, and Japanese sea cucumber, *Apostichopus japonicus*, is one of the most harvested species that appears in Asian markets especially China, Japan and Korea (Sloan, 1985; Yang et al., 2005). This species is traditionally used as a food source and has been proven to have antimicrobial, anticoagulant, and anti-inflammatory activities (Nagase et al., 1995; Beauregard, Truong, Zhang, Lin, & Beck, 2001; Cho & Bureau, 2001). In addition, cancer prevention, wound healing, aphrodisiac and curative properties are other benefits brought by this invertebrate (Gu et al., 2010; Seo & Lee, 2011). Because of these reasons, high world demand for sea cucumber has resulted in severe losses in stocks (Conand, 2004; Uthicke, 2004). In 2016, the South Korean production of sea cucumber was 2,386 metric tons, which valued approximately 26 million dollars (National Statistical Office, 2016). Whereas, feed development of sea cucumber aquaculture for the current state in South Korea has known to be inadequate. Besides, sea cucumber culture depends on imported feeds that mostly come from untrusted sources that have not been tested in South Korean culture environment (Kim et al., 2016). According to what was mentioned, it seems necessary for researchers to develop a high quality feed with reasonable price.

Binders are feed additives incorporated into aquafeeds in order to increase water-stability, improve firmness and

prevent easy breakdown of feed particles (NRC, 2011; Lee et al., 2016). Generally, several carbohydrate sources including guar gum, carrageenan and xanthan have been used as binding sources in aquaculture (Storebakken, 1985; Hashim & Saat, 1992; O'mahoney, Mouzakitis, Doyle, & Burnell, 2011). Guar gum is the most common product derived from guar bean (*Cyamopsis tetragonoloba*) that is an annual legume belonging to the Fabaceae family. It has been previously shown that the addition of guar gum to fish feed can result in more stable feces and increasing water quality (Amirkolaie, Leenhouders, Verreth, & Schrama, 2005; Brinker, 2007). Carrageenan is extracted from red seaweed and has been used widely as a food additive for humans from many years ago (Klose & Glicksman, 1968). Bautista, Millamena & Kanazawa (1989) reported the successful use of a carrageenan microdiet for nutrition of *Penaeus monodon* larvae. This additive also has been previously used as a binder in the microdiets (Liu et al., 2008). Xanthan gum is an extracellular polysaccharide that is resulted from pathogenic bacteria (*Xanthomonas campestris*) of brassicas (Becker, Katzen, Puhler, & Ielpi, 1998). By addition of salts and reduction of temperature a conformation transition occurs in xanthan gum and it turns into a rigid coil (Morris, Rees, Young, Walkinshaw, & Darke, 1977). Xanthan gum has been previously tested as binder in pharmaceutical (Eyo-okon & Hilton, 2003) and aquafeeds (O'Mahoney, Mouzakitis, Doyle, & Burnell, 2011).

In order to reduce nutrient leakage, increase feed efficiency, reduce wastage and improve water stability, proper application of binders seems to be indispensable (Fagbenro & Jauncey, 1995; Hardy, 1989; Huang, 1989). However, current studies on the guar gum, carrageenan, xanthan gum as a binders have been scarce. Therefore, the aim of the present study was to evaluate the effects of different dietary binders such as guar gum, carrageenan, xanthan gum as potential binders on growth performance and water analysis in juvenile sea cucumber *Apostichopus japonicus* culture.

Materials and Methods

Experimental diets

Formulation and proximate composition of the experimental diets are shown in Table 1. A basal diet without binder was used as control (CON), three diets with each of three different binder sources such as guar gum (G₁₀), carrageenan (C₁₀) and xanthan gum (X₁₀) at 10%, and other three diets (G₅C₅, C₅X₅ and X₅G₅) were formulated to contain mixture of two different binders at 5% for each. The basal diets were formulated to contain 24% crude protein (Zhu, Mai, Zhang, Wang, & Xu, 2005) and 5% crude lipid. Fishmeal, blood meal, dehulled soybean meal, squid liver powder and corn gluten meal served as the major protein sources in the experimental diets. Fish oil was used as lipid source while umitorano-powder, comb pen shell by-product meal and seaweed powder were the carbohydrate sources. Briefly, after thoroughly mixing the dry ingredients, fish oil and filtered tap water were added to the experimental diets. After drying, the experimental diets were broken up and sieved into 150 µm, and stored at -20 °C in refrigerator until use.

Experimental Fish and Feeding Trial

The feeding trial was carried out at the National Institute of Fisheries Science (NIFS), Pohang, Republic of Korea. Juvenile sea cucumbers were obtained from a commercial hatchery, wando, Republic of Korea, and their health condition was checked until one weeks, upon arrival and one weeks thereafter. Prior to the start of the feeding trial, all sea cucumbers were acclimatized to the experimental status and facilities. At the start of the

experiment, 15 juvenile sea cucumbers with an initial average weight of 1.09 ± 0.01 g (mean \pm SD) were randomly distributed into each of the 21 plastic rectangular tanks (50 L) supplied with flow-through sea water at 2 L min^{-1} . Each tank was then randomly assigned to one of the three replicates of seven dietary treatments. Sea cucumbers were fed twice daily the experimental diets at the rate of 1~3% wet body weight per day for twelve weeks. Supplemental aeration was provided to maintain the dissolved oxygen (DO) near saturation, and water temperature and pH during the experiment were maintained at 14 ± 0.5 °C using an air conditioner and 8.0 ± 0.5 , respectively. A photoperiod of 12h light: 12h dark was used throughout the experimental period.

Sample Collection and Analysis

At the end of the feeding trial, all of sea cucumbers were starved for 24 h, and used weight method (Liao, Ren, He, Jiang, & Han, 2014). The number of sea cucumbers in each tank were checked for the calculation of weight gain (WG), specific growth rate (SGR) and survival. Five sea cucumbers per tank were randomly collected and individually weighed, then dissected to evaluate viscera for the determination of viscerosomatic index (VSI). Five sea cucumbers were randomly selected from each aquarium and frozen at -20 °C for analysis of whole body proximate composition. The proximate composition analyses of the experimental diets and whole-body of fish were performed by the standard methods of AOAC (AOAC, 1995). Samples of diets and fish were dried at 105 °C to determine their moisture contents. The ash content was determined by incineration at 550 °C, crude protein was determined using the Kjeldahl method ($N \times 6.25$) after acid digestion, and crude lipid was measured by ether extraction using the Soxhlet system 1046 (Tacator AB, Hoganas, Sweden) after freeze-drying the samples for 20 h.

Melt Away Rate

A piece of nylon net sheet (mesh size, 1 mm) was made to measure the underwater loss rate. Approximately 2 g of the experimental feed was placed in the nylon net sheet that was submerged in water. To evaluate the loss rate, dry feed was weighted before and after submersion at 0, 1, 2, 4, 8 and 12 hours using an oven.

Water Quality Analysis

About 2 g of feed was put into 1 L seawater to investigate loss rate of feed on time variation (0, 1, 2, 4, 8 and 12 hours) of chemical oxygen demand (COD) measure. COD was determined using a water quality analyzer (HACH, DR/850, USA).

Statistical Analysis

All data were analyzed by one-way ANOVA (Statistics 3.1; Analytical Software, St. Paul, MN, USA) to test for the dietary treatments. When a significant treatment effect was observed, a LSD test was used to compare means. Treatment effects were considered significant at $P < 0.05$ level.

Results and Discussions

Several studies have been conducted to develop artificial diets of sea cucumber using different plant materials (Seo, Shin, & Lee, 2011; Wu et al., 2015; Yu et al., 2015). In the present study, we evaluated the effects of different plant based ingredients as binders in the diet of Japanese sea cucumber. After 12 weeks of feeding trial, the growth performance and survival of sea cucumbers fed different experimental diets are presented in Table 3. WG of sea cucumber fed G₁₀ and X₅C₅ diets was significantly higher than those of sea cucumber fed CON diet ($P < 0.05$). However, there were no significant differences in WG of sea cucumber fed CON, C₁₀, X₁₀ and G₅C₅ diets ($P > 0.05$). On the other hand, the sea cucumber fed G₅X₅ diet had the lowest WG ($P < 0.05$). SGR indicated the same trend with WG and SGR of sea cucumber fed G₅X₅ diet was significantly lower than those of sea cucumber fed CON, G₁₀, X₁₀ and X₅C₅ diets ($P < 0.05$). Sea cucumber fed G₁₀ and X₅C₅ diets showed the highest SGR, although there was no significant difference among CON, C₁₀, X₁₀ and G₅C₅ diets ($P < 0.05$). Sea cucumber survival rate varied range from 68.9 to 82% and no significant differences were observed among the dietary treatments. In addition, VSI was not significantly affected by inclusion of different dietary binders. Previous researches demonstrated the effective results of binders such as guar gum, carrageenan, xanthan gum on growth performance of fish (D'Abramo, 2003; Kumar et al., 2014; O'mahoney et al., 2011; Pearce, Daggett, & Robinson, 2002). Results of our study showed that binders such as guar gum and mixture of carrageenan and xanthan gum were utilized more efficiently than seaweed powder by sea cucumber. Pearce *et al.* (2002) reported that guar gum based diet may be effective in terms of growth comparing with control diet in green sea urchin *Strongylocentrotus droebachiensis*. In addition, Leenhouders, Adjei-boateng, Verreth, and Schrama (2006) and Brinker and Reiter (2011) demonstrated that guar gum diet did not show adverse effects on growth performance as well as digesta viscosity of fish. Moreover, Slater, Lassudrie, and Jeffs (2011) indicated that best results on digestibility can be achieved by carrageenan-based diet as comparing with other carbohydrate sources in juvenile sea cucumber, *Australostichopus mollis*. In our study, the results for growth performance showed that some binder sources can improve growth performance of juvenile sea cucumber, *A. japonica*.

Whole-body proximate composition of sea cucumber is shown in Table 4. There were no significant differences in whole-body crude protein, crude lipid, moisture and ash content of sea cucumbers fed different experimental diets ($P > 0.05$). Seo et al. (2011) also reported that when different plant ingredients are used as carbohydrate sources, whole body proximate composition of sea cucumber is not significantly affected.

Binders play a very important role in the manufacture of aquaculture diets to protect heavy losses from feed disintegration, nutrient leaching, and water contamination (Paolucci, Fabbrocini, Volpe, Varricchio, & Coccia, 2012). Because of these features, many experiments on binders have been carried out for the artificial diets in aquaculture and it was shown that mostly binder-based diets caused improved water stability (Liu et al., 2008; Obaldo, Divakaran, & Tacon, 2002; Orire, Sadiku, & Tihamiyu, 2010). Cumulative melt away rate of the experimental diets of sea cucumber for 12 hours is shown in Fig 1. The cumulative melt away rate for G₁₀, X₅C₅, X₁₀, G₅C₅ and G₅X₅ diets showed significantly lower amounts than CON diet during 12 hours ($P < 0.05$). This is while, the cumulative melt away rate of C₁₀ diet was not significantly different with CON diet during 4 hours. However, after 8 hours C₁₀ diet showed lower melt away rate than the CON diet. Feed stability in water is considered as retention of physical pellet value with minimal decomposition and nutrient leaching during immersion before utilization. Experimental diets which easily break up into small particles can cause nutrient

leakage that could lead to induced water pollution, impaired growth, inefficient feed conversion, and poor survival rate (Obaldo et al., 2002). Caltagirone, Francour, and Fernandez (1992) found that 23.9% of guar gum in the feed could improve stability after 48 h immersion in seawater. In addition, Ali, Gropal, Ramana, and Nazer (2005) reported that guar gum and wheat flour-based pellets showed good water stability. Similar studies have been observed in many other fish species (Meyers & Zein-Eldin, 1972; Knauer, Britz, & Hecht, 1993; Brinker, 2007). In agreement with the results of this study, several studies have shown the reduction in water pollution by inclusion of different binders in the diet (Brinker, Koppe, & Rosch, 2005; Cho & Bureau, 2001; Rosas et al., 2008).

Variations of water chemical oxygen demand (COD) after for 12 hours is shown in Fig 2. According to the COD results, G₁₀ group showed significantly lower value than the CON group during 8 hours ($P < 0.05$). X₁₀ diet indicated the similar trend with G₁₀ diet until 8 hours and X₅C₅ and G₅C₅ diets caused no significant differences in COD with CON diet after 1~4 hours. However X₅C₅ diet showed significantly lower value than the CON group after 8 hours ($P < 0.05$). The COD is a value of water quality and defined as the quantity of consumed oxygen in relation to chemically oxidation of inorganic products (Zhao, Jiang, Zhang, Catterall, & John, 2004; Kim et al., 2017). However, no studies were found on estimation of the relation between dietary binders in aquafeeds and water COD value.

According to what was mentioned, binder based experimental diets gave better results in regard to growth and water quality compared with CON diet.

Therefore, the present study demonstrated that dietary guar gum for 10% and 5% xanthan gum + 5% carrageenan of diet could improve growth and water quality in juvenile sea cucumber culture. Future research could be dedicated to estimate optimum levels of guar gum in diet of Juvenile Sea cucumber *Apostichopus japonicus*.

Acknowledgments

This research has been financially supported by National Institute of Fisheries Science (20110117217-00) and Feeds and Foods Nutrition Research Center (FFNRC), Pukyong National University, Busan, Rep. of Korea.

References

- Ali, S.A., Gopal, C., Ramana, J.V., & Nazer, A.R. (2005). Effect of different sources of starch and guar gum on aqua stability of shrimp feed pellets. *Indian Journal of Fisheries*, 52, 301-305
- Amirkolaie, A.K., Leenhouders, J.I., Verreth, J.A.J., Schrama, J.W. (2005). Type of dietary fibre (soluble versus insoluble) influences digestion, faeces characteristics and faecal waste production in Nile tilapia (*Oreochromis niloticus* L.). *Aquaculture Research*, 36, 1157-1166. <http://dx.doi.org/10.1111/j.1365-2109.2005.01330.x>
- AOAC (Association of Official Analytical Chemists). (1995). Official Methods of Analysis, 16th ed. Association of Official Analytical Chemists, Arlington, VA.
- Bautista, M.N., Millamena, O.M., & Kanazawa, A. (1989). Use of kappa-carrageenan microbound diet (C-MBD) as feed for *Penaeus monodon* larvae. *Marine Biology*, 103, 169-173. <http://doi.org/10.1007/BF00543344>
- Beauregard, K.A., Truong, N.T., Zhang, H., Lin, W., & Beck, G. (2001). The detection and isolation of a novel antimicrobial peptide from the echinoderm *Cucumaria frondosa*. *Advances in Experimental Medicine and Biology*, 484, 55-62. http://doi.org/10.1007/978-1-4615-1291-2_5
- Becker, A., Katzen, F., Pühler, A., & Ielpi, L. (1998). Xanthan gum biosynthesis and application: a biochemical/genetic perspective. *Applied Microbiology and Biotechnology*, 50, 145-152. <http://doi.org/10.1007/s002530051269>
- Brinker, A., Koppe, W., & Rösch, R. (2005). Optimised effluent treatment by stabilised trout faeces. *Aquaculture*, 249, 125-144. <https://doi.org/10.1016/j.aquaculture.2004.12.029>
- Brinker, A. (2007). Guar gum in rainbow trout (*Oncorhynchus mykiss*) feed: the influence of quality and dose on stabilisation of faecal solids. *Aquaculture*, 267, 315-327. <https://doi.org/10.1016/j.aquaculture.2007.02.037>



- Brinker, A., & Reiter, R. (2011). Fish meal replacement by plant protein substitution and guar gum addition in trout feed, Part I: Effects on feed utilization and fish quality. *Aquaculture*, 310, 350-360. <https://doi.org/10.1016/j.aquaculture.2010.09.041>
- Caltagirone, A., Francour, P. & Fernandez, C. (1992). Formulation of an artificial diet for the rearing of the urchin *Paracentrotus lividus*: I. Comparison of different binding agents. In L. Scalera-Liaci & C. Canicatti (Eds.), *Proceedings of the Third European Conference on Echinoderms*, Lecce, Italy, A.A. Balkema press., 115–120 pp.
- Collin, P.D. (2004) U.S. Washington, DC: U.S. Patent and Trademark Office. United States Patent 6,767,890.
- Cho, C.Y., & Bureau, D.P. (2001). A review of diet formulation strategies and feeding systems to reduce excretory and feed wastes in aquaculture. *Aquaculture research*, 32, 349-360. <http://doi.org/10.1046/j.1355-557x.2001.00027.x>
- Conand, C. (2004). Present status of world sea cucumber resources and utilization: an international overview. In: A. Lovatelli, C. Conand, S. Purcell, S. Uthicke, J.F. Hamel and A. Mercier (Eds.), *Advances in sea cucumber aquaculture and management*, FAO, Rome: 13–23.
- D’Abramo, L.R. (2003). Microparticulate microbound diet for the culture of larval fish and crustaceans. United States Patent 6645536
- Eyo-okon, I.E., & Hilton, C.S. (2003). Use of Glucomannan Hydrocolloid as Filler Material in Prostheses. Konjac Technologies, LLC, South River, NJ. United States Patent 6537318
- Fagbenro, O., & Jauncey, K. (1995). Water stability, nutrient leaching and nutritional properties of moist fermented fish silage diets. *Aquacultural Engineering*, 14, 143-153. [https://doi.org/10.1016/0144-8609\(94\)P4432-B](https://doi.org/10.1016/0144-8609(94)P4432-B)
- Gu, M., Ma, H., Mai, K., Zhang, W., Ai, Q., Wang, X., & Bai, N. (2010). Immune response of sea cucumber *Apostichopus japonicus* coelomocytes to several immunostimulants in vitro. *Aquaculture*, 306, 49-56. <https://doi.org/10.1016/j.aquaculture.2010.05.024>
- Hardy, R.W. (1989). Diet preparation. In: J.E. Halver (Eds.), *Fish Nutrition*. San Diego, CA, Academic Press., 475–548 pp.
- Hashim, R., & Saat, M.A.M. (1992). The utilization of seaweed meals as binding agents in pelleted feeds for snakehead (*Channa striatus*) fry and their effects on growth. *Aquaculture*, 108, 299-308. [https://doi.org/10.1016/0044-8486\(92\)90114-Z](https://doi.org/10.1016/0044-8486(92)90114-Z)
- Huang, H.J. (1989). Aquaculture feed binders. *Proceedings of the People’s Republic of China Aquaculture and Feed Workshop* (pp. 316-319). American Soybean Association, Singapore, Library press., 433 pp.
- Kim, K.D., Bae, K.M., Han, H.S., Kim, K.W., Lee, B.J., Kim, S.S., Park, K.Y., & Kwon, O.N. (2016). Effects of the Dietary Inclusion of Sea Mud on the Growth and Body Composition of Juvenile Sea Cucumbers *Apostichopus japonicus*. *The Korean Society of Fisheries and Aquatic Science*, 49, 26-29. <https://doi.org/10.5657/KFAS.2016.0026>
- Klose, R.E., & Glicksman, M. (1968). Gums In Handbook of Food Additives. In T.E. Furia (Eds.), Cleveland, Ohio, CRC press., 313–375 pp.
- Knauer, J, Britz, P.J., & Hecht, T. (1993). The effect of seven binding agents on 24-hour water stability of an artificial weaning diet for the South African abalone, *Haliotis midae* (Haliotidae, Gastropoda). *Aquaculture*, 115, 327-334. [https://doi.org/10.1016/0044-8486\(93\)90146-P](https://doi.org/10.1016/0044-8486(93)90146-P)
- Kumar, V., Kumar, S., Pandey, P.K., Raman, R.P., Prasad, K.P., Roy, S., Kumar, A., & Kumar, K. (2014). Growth and hemato-immunological response to dietary i-carrageenan in *Labeo rohita* (Hamilton, 1822) juveniles. *The Israeli Journal of Aquaculture-Bamidgeh*, 66, 1-10. <http://hdl.handle.net/10524/49131>
- Leenhouders, J.I., Adjei-boateng, D., Verreth, J.A.J., & Schrama, J.W. (2006). Digesta viscosity, nutrient digestibility and organ weights in African catfish (*Clarias gariepinus*) fed diets supplemented with different levels of a soluble non-starch polysaccharide. *Aquaculture Nutrition*, 12, 111-116. <http://doi.org/10.1111/j.1365-2095.2006.00389.x>
- Lee, S., Moniruzzaman, M., Bae, J., Seong, M., Song, Y.J., Dosanjh, B., & Bai, S.C. (2016). Effects of extruded pellet and moist pellet on growth performance, body composition, and hematology of juvenile olive flounder, *Paralichthys olivaceus*. *Fisheries and Aquatic Sciences*, 19, 32. <http://doi.org/10.1186/s41240-016-0032-x>
- Liao, M., Ren, T., He, L., Jiang, Z., & Han, Y. (2014). Optimum dietary protein level for growth and coelomic fluid non-specific immune enzymes of sea cucumber *Apostichopus japonicus* juvenile. *Aquaculture nutrition*, 20, 443-450. <http://dx.doi.org/10.1111/anu.12098>
- Liu, F., Ai, Q., Mai, K., Tan, B., Ma, H., Xu, W., Zhang, W., & LiuFu, Z. (2008). Effects of Dietary Binders on Survival and Growth Performance of Postlarval Tongue Sole, *Cynoglossus semilaevis* (Gunther). *Journal of the World Aquaculture Society*, 39, 441–571. <http://doi.org/10.1111/j.1749-7345.2008.00177.x>
- Meyers, S.P., & Zein-Eldin, Z.P. (1972). Binders and Pellet Stability in Development of Crustacean Diets. *Journal of the World Aquaculture Society*. 3, 351-364. <http://doi.org/10.1111/j.1749-7345.1972.tb00079.x>
- Morris, E.R., Rees, D.A., Young, G., Walkinshaw, M.D., & Darke, A. (1977). Order-disorder transition for a bacterial polysaccharide in solution. A role for polysaccharide conformation in recognition between *Xanthomonas* pathogen and its plant host. *Journal of Molecular Biology*, 110, 1-16. [https://doi.org/10.1016/S0022-2836\(77\)80095-8](https://doi.org/10.1016/S0022-2836(77)80095-8)
- Ministry of Oceans and Fisheries. (2016). Survey on the status of fish culture. National Statistical Office, Daejeon, Korea
- Nagase, H., Enjyoji, K.I., Minamiguchi, K., Kitazato, K.T., Kitazato, K., Saito, H., & Kato, H. (1995). Depolymerized holothurian glycosaminoglycan with novel anticoagulant actions: antithrombin III-and heparin cofactor II-independent inhibition of factor X activation by factor IXa-factor VIIIa complex and heparin cofactor II-dependent inhibition of thrombin. *Blood*, 85, 1527-1534. [http://doi.org/0006-4971/9.5/8506-0005\\$3.00/0](http://doi.org/0006-4971/9.5/8506-0005$3.00/0)
- NRC (2011). Nutrient requirements of fish. National Academy Press, Washington, DC, USA.
- Obaldo, L.G., Divakaran, S., & Tacon, A.G. (2002). Method for determining the physical stability of shrimp feeds in water. *Aquaculture Research*, 33, 369-377. <http://doi.org/10.1046/j.1365-2109.2002.00681.x>
- O’mahoney, M., Mouzakitis, G., Doyle, J., & Burnell, G. (2011). A novel konjac glucomannan–xanthan gum binder for aquaculture feeds: the effect of binder configuration on formulated feed stability, feed palatability and growth



- performance of the Japanese abalone, *Haliotis discus hannai*. *Aquaculture Nutrition*, 17, 395-407. <http://doi.org/10.1111/j.1365-2095.2010.00816.x>
- Orire, A.M., Sadiku, S.O.E., & Tihamiyu, L.O. (2010). Evaluation of Yam Starch (*Discorea rotundata*) as Aquatic Feed Binder. *Pakistan Journal of Nutrition*, 9, 668-671. <http://doi.org/10.3923/pjn.2010.668.671>
- Paolucci, M., Fabbrocini, A., Volpe, M.G., Varricchio, E., & Coccia, E. (2012). Development of biopolymers as binders for feed for farmed aquatic organisms. In Z. Muchlisin (Eds.), *Aquaculture*, Rijeka, Croatia, InTech press., 1-25 pp.
- Pearce, C.M., Daggett, T.L., & Robinson, S.M. (2002). Effect of binder type and concentration on prepared feed stability and gonad yield and quality of the green sea urchin, *Strongylocentrotus droebachiensis*. *Aquaculture*, 205, 301-323. [https://doi.org/10.1016/S0044-8486\(01\)00685-8](https://doi.org/10.1016/S0044-8486(01)00685-8)
- Rosas, C., Tut, J., Baeza, J., Sánchez, A., Sosa, V., Pascual, C., Arena, L., Domingues, P., & Cuzon, G. (2008). Effect of type of binder on growth, digestibility, and energetic balance of *Octopus maya*. *Aquaculture*, 275, 291-297. <https://doi.org/10.1016/j.aquaculture.2008.01.015>
- Seo, J.Y., Shin, I.S., Lee, S.M. (2011). Effect of dietary inclusion of various plant ingredients as an alternative for *Sargassum thunbergii* on growth and body composition of juvenile sea cucumber *Apostichopus japonicus*. *Aquaculture Nutrition*, 17, 549-556. <http://doi.org/10.1111/j.1365-2095.2010.00849.x>
- Seo, J.Y., & Lee, S.M. (2011). Optimum dietary protein and lipid levels for growth of juvenile sea cucumber *Apostichopus japonicus*. *Aquaculture Nutrition*, 17, 56-61. <http://doi.org/10.1111/j.1365-2095.2009.00728.x>
- Slater, M.J., Lassudrie, M., & Jeffs, A.G. (2011). Method for determining apparent digestibility of carbohydrate and protein sources for artificial diets for juvenile sea cucumber, *Australostichopus mollis*. *Journal of the World Aquaculture Society*, 42, 714-725. <http://doi.org/10.1111/j.1749-7345.2011.00510.x>
- Sloan, N.A. (1985). Echinoderm Fisheries of the World: A Review. In B.F. Keegan & B.D.S. O'Connor (Eds.), *Echinodermata*, Rotterdam, Netherlands, AA Balkema press., 109-124 pp.
- Storebakken, T. (1985). Binders in fish feeds: I. Effect of alginate and guar gum on growth, digestibility, feed intake and passage through the gastrointestinal tract of rainbow trout. *Aquaculture*, 47, 11-26. [https://doi.org/10.1016/0044-8486\(85\)90004-3](https://doi.org/10.1016/0044-8486(85)90004-3)
- Uthicke, S. 2004. Overfishing of holothurians: lessons from the Great Barrier Reef. In A. Lovatelli, C. Conand, S. Purcell, S. Uthicke, J.H. Hamel & A. Mercier (Eds.), *Advances in sea cucumber aquaculture and management*, FAO, Rome :163-173.
- Kim, J., Park, H., Hwang, I., Han, J., Kim, D., Oh, C.W., Lee, J.S., & Kang, J. (2017). Alterations of growth performance, hematological parameters, and plasma constituents in the sablefish, *Anoplopoma fimbria* depending on ammonia concentrations. *Fisheries and Aquatic Sciences*, 20, 4. <http://doi.org/10.1186/s41240-017-0049-9>
- Wu, B., Xia, S., Rahman, M.M., Rajkumar, M., Fu, Z., Tan, J., & Yang, A. (2015). Substituting seaweed with corn leaf in diet of sea cucumber (*Apostichopus japonicus*): Effects on growth, feed conversion ratio and feed digestibility. *Aquaculture*, 444, 88-92. <https://doi.org/10.1016/j.aquaculture.2015.03.026>
- Yang, H., Yuan, X., Zhou, Y., Mao, Y., Zhang, T., & Liu, Y. (2005) Effects of body size and water temperature on food consumption and growth in the sea cucumber *Apostichopus japonicus* (*Selenka*) with special reference to aestivation. *Aquaculture Research*, 36, 1085-1092. <http://doi.org/10.1111/j.1365-2109.2005.01325.x>
- Yu, H.B., Gao, Q.F., Dong, S.L., Wen, B., Hou, Y.R., & Ning, L.G. (2015). Utilization of corn meal and extruded soybean meal by sea cucumber *Apostichopus japonicus* (*Selenka*): Insights from carbon stable isotope analysis. *Aquaculture*, 435, 106-110. <https://doi.org/10.1016/j.aquaculture.2014.09.036>
- Zhao, H., Jiang, D., Zhang, S., Catterall, K., & John, R. (2004). Development of a direct photoelectrochemical method for determination of chemical oxygen demand. *Analytical chemistry*, 76, 155-160. <http://doi.org/10.1021/ac0302298>
- Zhu, W., Mai, K.S., Zhang, B.G., Wang, F.Z., & Xu, G.Y. (2005). Study on dietary protein and lipid requirement for sea cucumber, *Stichopus japonicus*. *Marine Sciences*, 29, 54-58.

Table 1. Composition of the basal diet

Ingredients	
Fish meal ¹	5.00
Pork blood meal ¹	3.10
Dehulled soybean meal ¹	9.00
Squid liver powder ¹	6.00
Squid meal ¹	2.65
Corn gluten meal ¹	3.84
Umitorano-powder ²	3.00
Comb pen shell by-product meal ²	2.00
Seaweed powder ¹	20.5
Wheat flour	20.5
Fish oil ¹	1.41
Tideland ²	20.5
Saccharomyces cerevisiae ²	0.50
Vitamin premix ³	1.00
Mineral premix ⁴	1.00
Moisture (%)	9.05
Crude Protein (%)	24.0
Crude Lipid (%)	5.14
Ash (%)	34.0

¹ Supplied by Suhyup Co., Buan, Korea.

² Supplied by Myeong cheon Co., Wando, Korea.

³ Contains (as mg/kg in diets) : Ascorbic acid, 300; dl-Calcium pantothenate, 150 ; Choline bitartrate, 3000; Inositol, 150; Menadione, 6; Niacin, 150; Pyridoxine.HCl, 15; Riboflavin, 30; Thiamine mononitrate, 15; dl- α -Tocopherol acetate, 201; Retinyl acetate, 6; Biotin, 1.5; Folic acid, 5.4; B₁₂, 0.06.

⁴ Contains (as mg/kg in diets): NaCl, 437.4; MgSO₄.7H₂O, 1379.8; NaH₂P₄.2H₂O, 877.8; Ca(H₂PO₄)₂.2H₂O, 1366.7; KH₂PO₄, 2414; ZnSO₄.7H₂O, 226.4; Fe-Citrate, 299; Ca-lactate, 3004; MnSO₄, 0.016; FeSO₄, 0.0378; CuSO₄, 0.00033; Calcium iodate, 0.0006; MgO, 0.00135; NaSeO₃, 0.00025.

Table 2. Proximate composition of Seaweed & binders¹

	Seaweed	Guar gum	Carrageenan	Xanthan gum
Moisture (%)	19.4	10.63	9.94	12.07
Protein (%)	6.7	4.23	0.58	4.45
Lipid (%)	4.1	0.71	0.24	0.33
Ash (%)	25.0	4.73	0.64	5.06

¹Results reported on the dry matter basis, average of triplicate**Table 3.** Percent weight gain, specific growth rate and survival for sea cucumber fed the experimental diets (CON, G₁₀, C₁₀, X₁₀, X₅C₅, G₅C₅ and G₅X₅) for 12 weeks¹

	IW ² (g)	FW ³ (g)	WG ⁴ (%)	SGR ⁵ (%/day)	VSI ⁶	Survival (%)
CON	1.11	3.87 ^{bc}	249 ^{bc}	1.78 ^{bc}	12.6	78.9
G ₁₀	1.10	4.77 ^a	332 ^a	2.09 ^a	12.6	81.0
C ₁₀	1.11	3.44 ^{bcd}	210 ^{bcd}	1.62 ^{bcd}	12.7	72.0
X ₁₀	1.09	3.83 ^{bc}	246 ^{bc}	1.77 ^{bc}	12.4	82.0
X ₅ C ₅	1.11	4.51 ^a	311 ^a	2.02 ^a	12.8	74.5
G ₅ C ₅	1.11	3.41 ^{cd}	207 ^{cd}	1.60 ^{cd}	12.5	73.3
G ₅ X ₅	1.12	3.24 ^d	190 ^d	1.52 ^d	13.0	68.9
Pooled SEM ⁷	0.00	0.13	11.9	0.05	0.16	2.03

¹ Values are means from triplicate groups of fish where the values in each row with different superscripts are significantly different (P<0.05).² Initial weight (g/fish)³ Final weight (g/fish)⁴ Weight gain (%) = (final weight - initial weight) × 100/initial weight⁵ Specific growth rate (%) = 100 × (loge final weight - loge initial weight) / days⁶VSI: visceral somatic index = visceral weight / body weight × 100⁷ Pooled standard error of means: SD/√n**Table 4.** Proximate analysis of whole-body composition for sea cucumber fed the experimental diets (CON, G₁₀, C₁₀, X₁₀, X₅C₅, G₅C₅, G₅X₅) for 12 weeks (% dry matter basis)¹.

	Moisture (%)	Crude protein (%)	Crude lipid (%)	Ash (%)
--	--------------	-------------------	-----------------	---------



CON	90.3	42.6	4.3	41.2
G ₁₀	91.5	43.0	4.2	40.4
C ₁₀	90.2	43.6	4.3	41.4
X ₁₀	90.6	42.7	3.7	42.2
X ₅ C ₅	90.6	43.3	4.2	42.2
G ₅ C ₅	90.7	43.4	3.9	42.3
G ₅ X ₅	90.8	43.1	3.6	41.0
Pooled SEM ⁷	0.18	0.09	0.30	0.33

¹Values are means from triplicate groups of fish where the means in each row with a different superscripts are significantly different (P<0.05)

²Pooled standard error of means: SD/\sqrt{n}

Accepted Manuscript




Figure 1. Cumulate melt away rate for experimental diets (CON, G₁₀, C₁₀, X₁₀, X₅C₅, G₅C₅ and G₅X₅) for 12 hrs into the water




Figure 2. Water COD changes after the addition of diets (CON, G₁₀, C₁₀, X₁₀, X₅C₅, G₅C₅ and G₅X₅) for 12 hrs into the water