



Bacterial and Viral Fish Diseases in Turkey

Rafet Çağrı Öztürk¹, İlhan Altınok^{1,*}

¹ Karadeniz Technical University, Faculty of Marine Science, Department of Fisheries Technology Engineering, 61530 Sürmene, Trabzon, Turkey.

* Corresponding Author: Tel.: +90.462 3778083; Fax: +90.462 7522158;
E-mail: ialtinok@ktu.edu.tr

Received 1 January 2014
Accepted 28 February 2014

Abstract

This review summarizes the state of knowledge about the major bacterial and viral pathogens of fish found in Turkey. It also considers diseases prevention and treatment. In this study, peer reviewed scientific articles, theses and dissertations, symposium proceedings, government records as well as recent books, which published between 1976 and 2013 were used as a source to compile dispersed literature. Bacterial and viral disease problems were investigated during this period in Turkey. Total of 48 pathogen bacteria and 5 virus species have been reported in Turkey. It does mean that all the bacteria and virus present in fish have been covered since every year new disease agents have been isolated. The highest outbreaks occurred in larval and juvenile stages of the fish. This article focused on geographical distribution, host range, and occurrence year of pathogenic bacteria and virus species. Vibriosis, Furunculosis, Motile Aeromonas Septicemia, Yersiniosis, Photobacteriosis and Flavobacteriosis are among the most frequently reported fish diseases. Meanwhile, *Vagococcus salmoninarum*, *Renibacterium salmoninarum*, *Piscirickettsia salmonis* and *Pseudomonas luteola* are rarely encountered pathogens and might be emerging disease problems. Finally, the current status in fish diseases prevention and their treatment strategies are also addressed.

Keywords: Disease transfer, vaccines, disease treatment, disease prevention.

Türkiye’de Görülen Bakteriyel ve Viral Balık Hastalıkları

Özet

Bu derlemede Türkiye’de bulunan başlıca bakteriyel ve viral balık patojenleri hakkında ki bilgi durumu özetlenmiştir. Bu bağlamda 1976 ve 2013 yılları arasında yayımlanan bilimsel makaleler, yüksek lisans ve doktora tezleri, sempozyum bildirileri, resmi kayıtlar ve kitaplarda dağınık olarak bulunan kayıtlar bir araya getirilmiştir. Türkiye’de bu dönemdeki bakteriyel ve viral hastalıkları incelenmiş ve toplam 48 patojenik bakteri ve 5 virüs türü rapor edilmiştir. Her yıl yeni hastalık etkenleri izole edilmesinden dolayı bu çalışmanın mevcut tüm bakteri ve virüsleri kapsadığı anlamı ortaya çıkmamalıdır. En yüksek salgınlar balıkların larval ve yavru dönemlerinde meydana gelmektedir. Bu makalede, patojenik bakteri ve virüs türlerinin coğrafik dağılımı, konakçı ve ortaya çıktıkları yıllara odaklanmıştır. Rapor edilen hastalıklar arasında en sık bildirilen balık hastalıkları Vibriosis, Furunculosis, Hareketli Aeromonas Septisemi, Yersiniosis, Photobacteriosis ve Flavobacteriosis olarak yer almaktadır. Bu arada *Vagococcus salmoninarum*, *Renibacterium salmoninarum*, *Piscirickettsia salmonis* ve *Pseudomonas luteola* nadiren karşılaşılan patojenler olmasına rağmen balıklarda sorunlar oluşturabileceği düşünülmektedir. Son olarak, mevcut balık hastalıklarını önleme ve tedavi stratejileri de ele alınmıştır.

Anahtar Kelimeler: Hastalık transferi, aşılarda, hastalıkların tedavisi, hastalıklardan korunma.

Introduction

Fish diseases are playing one of the roles as a limiting factor in fish production and causing heavy mortalities especially in hatcheries thus affecting profit negatively. Both researchers and farmers in Aquaculture area are looking for a ways to get maximum amount of yield from per unit volume of

water to lower the cost in aquaculture operations. One simple way to get maximum amount of yield from per unit volume of water is basically overcrowding the fish and this condition causes stress in fish which are accustomed to live freely in nature and causes unexpected losses (Toksen, 1999). Therefore, it is essential that proper measurements are taken against stress causing agents such as bad water

quality, bad feed nutrition, and heavy stocking rate etc. The best way to deal with diseases is to prevent both infectious and noninfectious outbreaks. Beyond that, correct diagnosis and economically acceptable treatment methods should be carried out.

Some of the bacterial species that inhabit the aquatic environment are essential to the balance of nature with no direct consequence in causing disease in fish. However, approximately 125 different bacterial species belonging to 34 different bacterial families have been associated with various fish diseases in the world. Furthermore, nearly 100 fish viruses belonging to 16 virus families have been isolated worldwide while only 5 fish viruses identified in Turkey. The list of viral and bacterial fish pathogens keeps extending. The water used in aquaculture operations provides a natural habitat for growth and proliferation of bacteria which can be influenced by nutrient availability, pH, temperature, and other factors that affect their growth pattern, virulence, and pathogenicity. In order to grow, bacteria need an organic substrate that provide nutrients; some survive as free living organisms or exist as fish pathogens, while others are fastidious and survive indefinitely only within a host (Plumb and Hanson, 2011). Most bacteria responsible for causing disease in fish are gram negative rods but some pathogens that are gram-positive rods or cocci and a few that are acid-fast rods also cause disease in aquatic animals.

Since rainbow trout (*Oncorhynchus mykiss*), Gilt-head sea bream (*Sparus aurata*), and Sea bass (*Dicentrarchus labrax*) are the most predominant fish species cultured in Turkey (TUIK, 2012), most of the diseases agents have been isolated from these three species. Fish diseases identification or etiological agent isolation is mainly concentrated in certain location where Faculty members or Fisheries Institute employee work on fish diseases. Furthermore, most of the fish diseases agents isolated in Turkey after 2000s because of the increasing number of fish diseases professional.

Compared with bacterial and parasitic diseases of fish, studies about viral diseases of fish in Turkey is relatively new. First proven viral fish disease in Turkey was infectious pancreatic necrosis virus (IPNV). Although comparatively few fish viruses cause severe disease in aquaculture, results can be devastating when viral diseases outbreak. Most known fish viruses have been reported in freshwater cultured species, some occur in marine fish only, and others are found in both environments (Plumb and Hanson, 2011).

Main reason for spreading diseases from one location to another or from country to country is uncontrolled fish transfer. To avoid transferring pathogen from one location to another, specific pathogen free stocks should be developed and fish transfer should be controlled. Unfortunately, fish transferred between fish farms are not controlled in

the Turkey. Although transportation of fish is regulated by the state, implementation of the regulations are mostly lacking. Based on regulations imposed, whenever a farmer wants to transport fish, an application must be filed to the provincial directorate of agriculture to appoint a veterinarian to check the fish health for virus, parasite, bacteria and fungus. However, whoever wants to transport fish pays a visit to a veterinarian who works for the provincial directorate of agriculture to obtain transport certificate. In this case veterinarian may give the certificate without seeing fish. Therefore, infectious diseases have been spreading region to region. Yersiniosis caused by *Yersinia ruckeri* and lactococcosis diseases caused by *Lactococcus garvieae* were first reported in Aegean region and they have been spread all over the Turkey where fish cultured. Newly arrived fish or eggs should be given a prophylactic treatment with appropriate drugs to remove any external pathogens. When possible, new fish should be segregated from the residential population until they are shown to be disease free. Certain fish production facilities should produce certified disease-free fish or eggs (Plumb and Hanson, 2011). Because of the production systems, lack of legal restraints to shipping and limited number of pathogens to target, no facilities currently market disease free eggs or fry in the Turkey. Another problem is the fish eggs or ornamental fish import. When live fish are imported from any country, there is no quarantine procedure taken at the port of entry. Therefore, exotic pathogens are directly being imported to country and spread all the regions. These problem have to be solved by responsible government authorities.

Since the most common causative agents of infectious diseases in aquaculture, are bacteria and viruses, in this review only both of them are covered. The other fish diseases caused by parasites and fungi are discarded. This review examines the major bacterial and viral pathogens of fish found in Turkey. It also considers diseases prevention and treatment. For this purpose peer reviewed scientific articles, theses and dissertations, symposium proceedings, government records as well as recent books, which published between 1976 and 2013 were used as a source to compile dispersed literature.

Bacterial Diseases

Vibriosis

Vibriosis is a disease caused by bacteria belonging to the genus *Vibrio*. This disease possesses wide distribution and host range worldwide. Losses associated with it has been reported in many fish species, including sea bass, sea bream, salmonid spp., cod (*Gadus morhua*), European eel (*Anguilla anguilla*), turbot (*Psetta maxima*), and tilapia (*Oreochromis niloticus*) (Toranzo and Barja, 1990).

Although the causative agents of vibriosis are *Listonella anguillarum*, *V. ordalii*, *V. vulnificus*, *V. harveyi*, and *V. alginolyticus* (Austin and Austin, 2012; Toranzo *et al.*, 2005), the main causative agent of the disease is *L. anguillarum*. Ten different serotypes of this pathogen described (O1-O10) so far. Serotype 1 and 2 are responsible for the most of the outbreaks. Among the *Vibrio* species *L. anguillarum* was first described in 1909 by Bergamn as the etiological agent of the 'red pest of eels' in the Baltic Sea (Woo and Bruno, 2011). According to published literature in Turkey, *L. anguillarum* was first isolated from diseased sea bream in Mugla (Candan, 1991a) and couple of years later the disease causing agent was also isolated from sea bass, red porgy (*Pagrus pagrus*) and rainbow trout (Cagirgan, 1993a; Korun and Gokoglu, 2007a; Tanrikul, 2007). The pathogen possesses wide distribution throughout Turkey (Table 1) because of uncontrolled fish transfer.

Vibrio ordalii is another important etiological agent of the vibriosis. It was initially described in coho salmon (*O. kisutch*) in USA (Schiewe and Crosa, 1981). First isolation of *V. ordalii* in Turkey was reported from sea bream in Mugla (Akayli, 2001). It was seven years after when same bacteria was also reported from sea bass in Aegean Sea

(Korun and Timur, 2008).

Other identified pathogen species of *Vibrio* causing fish diseases also described in Turkey including *V. alginolyticus*, *V. vulnificus*, *V. harveyi* and *V. parahaemolyticus*. *V. alginolyticus* first reported from both sea bass and sea bream in Mugla (Cagirgan, 1993a). Other susceptible species are rainbow trout and cultured horse mackerel (*Trachurus mediterraneus*) (Savaş *et al.*, 2006; Boran *et al.*, 2013). *V. vulnificus* was first isolated from sea bream in Aegean Sea (Turk, 2002) while *V. harveyi* was first observed in sea bass in Antalya (Korun and Akayli, 2004a). The other *Vibrio* species is *V. parahaemolyticus*, first isolated from rainbow trout in Aydin (Aydin, 2000a).

Some of the vibrio species can be isolated in freshwater fish. It does not mean that *Vibrio* species can infect fish in fresh water or facultative vibrio species can live in fresh water. In Black Sea region, especially North East cost of the Turkey, rainbow trout have been cultured in freshwater raceways and when fish weight reached to 50 g or more especially fish farmers prefer 150 g or more and water temperature decrease to 18°C in October or November, they are transferred to sea cages. They are kept there until the water temperature reached to 20°C

Table 1. Distribution and host range of *Vibrio* spp.

Common name of the disease	Etiological agents	Host	Geographical distribution	References
Red-pest, Vibriosis, Salt-water furunculosis, Boil disease	<i>Listonella anguillarum</i> (<i>Vibrio anguillarum</i>)	<i>Sparus aurata</i>	Mugla	Candan (1991)
		<i>Dicentrarchus labrax</i>	Aegean Sea	Cagirgan (1993a)
		<i>Salmo salar</i>	Black Sea	Candan (2000)
		<i>Dicentrarchus labrax</i>	Izmir, Mugla, Aydin	Tanrikul <i>et al.</i> (2004)
		<i>Dicentrarchus labrax</i>	Antalya	Korun (2004a)
		<i>Dicentrarchus labrax</i>	Ordu	Savaş <i>et al.</i> (2006)
		<i>Pagrus pagrus</i>	Mugla	Korun and Gokoglu (2007a)
		<i>Aulonocara maylandi</i>	Istanbul (Aquarium)	Akayli (2007)
		<i>Oncorhynchus mykiss</i>	Mugla, Aydin, Denizli	Aksit and Kum (2008)
			<i>Sparus aurata</i>	Izmir, Mugla
	<i>Oncorhynchus mykiss</i>	Black Sea	Timur <i>et al.</i> (2011)	
Vibriosis, Eye disease, Septicemia	<i>Vibrio alginolyticus</i>	<i>Dicentrarchus labrax</i>	Aegean Sea	Cagirgan (1993a)
		<i>Sparus aurata</i>	Aegean Sea	Cagirgan (1993a)
		<i>Oncorhynchus mykiss</i>	Ordu	Savaş <i>et al.</i> (2006)
		<i>Dicentrarchus labrax</i>	Ordu	Savaş <i>et al.</i> (2006)
		<i>Trachurus trachurus</i>	Mersin	Ozer <i>et al.</i> (2008a)
	<i>Trachurus mediterraneus</i>	Trabzon	Boran <i>et al.</i> (2013)	
Vibriosis	<i>Vibrio ordalii</i>	<i>Sparus aurata</i>	Mugla	Akayli (2001)
		<i>Dicentrarchus labrax</i>	Aegean Region	Turk (2002)
		<i>Dicentrarchus labrax</i>	Izmir	Korun (2004a)
		<i>Dicentrarchus labrax</i>	Canakkale	Tanrikul <i>et al.</i> (2004)
		<i>Dicentrarchus labrax</i>	Antalya	Korun (2004a)
Vibriosis	<i>Vibrio vulnificus</i>	<i>Sparus aurata</i>	Aegean Region	Turk (2002)
		<i>Dicentrarchus labrax</i>	Aegean Region	Turk (2002)
		<i>Trachurus mediterraneus</i>	Trabzon	Boran <i>et al.</i> (2013)
Vibriosis, Eye disease, Granuloma	<i>Vibrio harveyi</i>	<i>Dicentrarchus labrax</i>	Mugla	Korun and Akayli (2004)
		<i>Sparus aurata</i>	Mugla, Izmir	Canak (2011)
Vibriosis	<i>Vibrio parahaemolyticus</i>	<i>Oncorhynchus mykiss</i>	Aydin	Aydin (2000a)

or until the harvest to sell. Sometimes, some farmer cannot sell their fish when water temperature increase in April or May; Therefore, they transfer their fish from sea cages to freshwater tank or raceways. Meanwhile, *Vibrio* species can also transferred from seawater to freshwater with the fish. Although *Vibrio* species cannot stay in freshwater for a long time, they can be isolated from fish since fish osmolality closed to 9‰ salinity. It does not mean that *vibrio* species can infect fresh water fish in fresh water.

Furunculosis

Aeromonas salmonicida subsp. *salmonicida* is known as an etiologic agent of furunculosis, one of the oldest known diseases and its origin is unclear and questionable. Etiological agent is also known as “typical *A. salmonicida*” and it should not be confused with “atypical *A. salmonicida*”. The early isolation of the etiologic agent was made in 1930s (Mackie *et al.*, 1930; Austin and Austin, 2012). Furunculosis is a prevalent disease of cultured salmonids, which causes serious economic crisis in the industry. Although salmonids seem to be most susceptible species, this particular disease has been reported more than 50 different fish species worldwide (McFadden, 1970; Bernoth, 1977; Barker and Kehoe, 1995; Kirkan *et al.*, 2003).

The pathogen bacterium was first isolated from rainbow trout fry in Marmara Region, Turkey (Timur *et al.*, 1999). Following reports have been documented from different parts of Turkey (Table 2) but so far, only it has been observed in rainbow trout. The outbreaks generally occurred in hatcheries and ended up with high mortalities (Timur *et al.*, 1999; Kirkan *et al.*, 2003).

Goldfish Ulcer Disease

The causative agent of the disease is *Aeromonas salmonicida* subsp. *achromogenes*, which is also known as atypical *A. salmonicida*. Ulcer disease was initially diagnosed in the early 1960s in trout in U.K. (Smith, 1963). Atypical *A. salmonicida* has become a major disease in cyprinids (Plumb and Hanson, 2011). The disease has been confirmed in many countries

including European countries, USA and Australia (Plumb and Hanson, 2011).

Few outbreaks of atypical *A. salmonicida* occurred in Turkey (Table 2). Korun and Timur (2001) were first reported the pathogen from rainbow trout in Balıkesir. Subsequently, it was reported from sea bass and turbot in Black Sea region (Karatas *et al.*, 2005; Savas and Ture, 2008).

Motile Aeromonas Septicemia (MAS)

MAS is associated with infections caused by *Aeromonas hydrophila*, *A. sobria*, *A. veronii*, and *A. caviae*. *Aeromonas hydrophila* is the predominant causative agent of MAS. These pathogens exist worldwide in fresh and brackish waters and occasionally in salt waters and they have a diverse host range. Motile *Aeromonas* spp. are considered as opportunistic pathogens and could easily found in organically rich waters. Thus, stress and poor water quality play a key role in occurrence. *A. hydrophila* has been recognized as a pathogen of fish since early 1960s in Europe (Levis and Bender, 1960) and in USA (Snieszko and Bullock, 1965). *A. sobria* has been recognized as a fish pathogen since 1987. Toranzo *et al.* (1989) was first reported it from wild gizzard shad (*Dorosoma cepedianum*) in USA.

In Turkey, *A. hydrophila* was first isolated from rainbow trout in Eskisehir (Baran *et al.*, 1980). The pathogen was subsequently isolated from other fish species such as sea bass, sea bream, eel, carp (*Cyprinus carpio*), sturgeon (*Acipenser gueldenstaedtii*), horse mackerel (*T. mediterraneus*) and from some ornamental fish species from different parts of Turkey (Table 3).

The other etiological agent of the disease, *A. sobria*, was reported from Atlantic salmon (*Salmo salar*) in Black Sea (Karatas, 1996). It also has a wide host range such as rainbow trout, sea bass and sea bream. Initially, *A. caviae* was isolated from different fish species in Keban Dam (Muz *et al.*, 1995). Afterwards, etiological agent was also isolated from rainbow trout, Atlantic salmon and some ornamental fish species (Candan *et al.*, 1995; Timur *et al.*, 2003; Korun and Toprak, 2010).

The only incidence of mortality caused by *A.*

Table 2. Distribution and host range of *A. salmonicida*

Common name of the disease	Etiological agent	Host	Geographical distribution	References
Ulcerative Disease	<i>A. salmonicida</i> subsp.	<i>Oncorhynchus mykiss</i> <i>Dicentrarchus labrax</i>	Balıkesir Black Sea	Korun and Timur (2001) Karatas <i>et al.</i> (2005)
	<i>achromogenes</i>	<i>Psetta maxima</i>	Eastern Black Sea	Savas and Ture (2008)
Furunculosis	<i>A. salmonicida</i> subsp. <i>salmonicida</i>	<i>Oncorhynchus mykiss</i>	Marmara Region	Timur <i>et al.</i> (1999)
			Aegean Region	Kirkan <i>et al.</i> (2003)
			Ankara	Ozkok (2005)
			Black sea	Timur <i>et al.</i> (2008)
			Mugla, Aydın, Denizli Trabzon, Rize	Aksit and Kum (2008) Kayis <i>et al.</i> (2009)

Table 3. Distribution and host of motile *Aeromonas* species

Common name of the disease	Etiological agents	Host	Geographical distribution	References
Motile Aeromonas septicemia (MAS),	<i>Aeromonas hydrophila</i>	<i>Oncorhynchus mykiss</i>	Eskisehir	Baran <i>et al.</i> (1980)
		<i>Anguilla anguilla</i>	Eskisehir	Timur (1983)
		Ornimental Fish spp.	Istanbul (Aquarium)	Guvener (2001)
		<i>Sparus aurata</i>	Aegean Region	Turk (2002)
		<i>Dicentrarchus labrax</i>	Aegean Region	Turk (2002)
		<i>Ctenopharyngodon idella</i>	Ankara	Uzbilek and Yildiz (2002)
		<i>Oncorhynchus mykiss</i>	Erzurum	Aydin and Ciltas (2004)
		<i>Xiphophorus hellerii</i>	Istanbul (Aquarium)	Akayli and Zeybek (2005)
		<i>Oncorhynchus mykiss</i>	Ordu	Savaş <i>et al.</i> (2006)
		<i>Dicentrarchus labrax</i>	Ordu	Savaş <i>et al.</i> (2006)
Red fin disease,	<i>Aeromonas hydrophila</i>	<i>Cyprinus carpio</i>	Sanliurfa	Tel <i>et al.</i> (2007)
		<i>Carassius auratus</i>	Antalya (Aquarium)	Korun and Toprak (2007b)
		<i>Cyprinus carpio</i>	Burdur	Ozturk <i>et al.</i> (2007)
Red pest,	<i>Aeromonas hydrophila</i>	<i>Oncorhynchus mykiss</i>	Mersin	Ozer <i>et al.</i> (2009)
		<i>Oncorhynchus mykiss</i>	Eastern Black Sea	Durmaz and Turk (2009)
		<i>Oncorhynchus mykiss</i>	Elazig	Aksoy (2009)
Infectious abdominal dropsy	<i>Aeromonas hydrophila</i>	<i>Acipenser gueldenstaedtii</i>	Istanbul	Timur <i>et al.</i> (2010)
		<i>Oncorhynchus mykiss</i>	Antalya	Korun and Toprak (2010)
Infectious abdominal dropsy	<i>Aeromonas sobria</i>	<i>Trachurus mediterraneus</i>	Trabzon	Boran <i>et al.</i> (2013)
		Unknown	Elazig	Muz <i>et al.</i> (1995)
		<i>Salmo salar</i>	Black Sea	Karatas (1996)
		<i>Oncorhynchus mykiss</i>	Ankara	Ozkok (2005)
		<i>Oncorhynchus mykiss</i>	Erzurum	Saglam <i>et al.</i> (2006)
		<i>Carassius auratus</i>	Antalya (Aquarium)	Korun and Toprak (2007b)
		<i>Oncorhynchus mykiss</i>	Black Sea	Timur <i>et al.</i> (2008)
		<i>Oncorhynchus mykiss</i>	Trabzon, Rize	Kayis vd. (2009)
		<i>Oncorhynchus mykiss</i>	Mersin	Ozer <i>et al.</i> (2009)
		<i>Oncorhynchus mykiss</i>	Antalya	Korun and Toprak (2010)
		<i>Sparus aurata</i>	Aegean Region	Avsever <i>et al.</i> (2012)
		<i>Dicentrarchus labrax</i>	Aegean Region	Avsever <i>et al.</i> (2012)
		-	Elazig	Muz <i>et al.</i> (1995)
		<i>Salmo salar</i>	Black Sea	Candan <i>et al.</i> (1995)
		<i>Poecilia reticulata</i>	Istanbul (Aquarium)	Timur <i>et al.</i> (2003)
<i>Carassius auratus</i>	Antalya (Aquarium)	Korun and Toprak (2007b)		
<i>Oncorhynchus mykiss</i>	Trabzon, Rize	Kayis <i>et al.</i> (2009)		
<i>Oncorhynchus mykiss</i>	Antalya	Korun and Toprak (2010)		
<i>Aeromonas schubertii</i>	<i>Oncorhynchus mykiss</i>	Mediterranean Region	Akayli <i>et al.</i> (2011)	

schubertii was occurred in Mediterranean region and pathogen was isolated from rainbow trout (Akayli *et al.*, 2011). This bacteria also belongs to motile *Aeromonas* species and might cause MAS.

Yersiniosis

It is one of the most important diseases in salmonids causing serious economic losses both in freshwater and marine salmonid aquaculture (Austin and Austin, 2012; Bullock and Snieszko, 1975; Toranzo, 2004). It also known as enteric red mouth disease (ERM), or yersiniosis. Causative agent of the disease is *Yersinia ruckeri*. It was first isolated from rainbow trout in Hagerman Valley, USA in 1950s (Ross *et al.*, 1966). Yersiniosis is commonly observed from all over the world where salmonids are cultured (Plumb and Hanson, 2011).

In Turkey, first outbreak of this disease occurred

in one of the rainbow trout farm in southern Anatolia (Timur and Timur, 1991b). Disease agent came from Africa with imported rainbow trout summer eggs. After two decades, it became one of the widest distributed fish pathogen in Turkey (Table 4). Like the rest of the world, it was mostly seen in salmonids but it was also isolated from sea bass (Savaş *et al.*, 2006). Recently, number of *Y. ruckeri* isolation were decreased may be due to vaccination of fish with *Y. ruckeri* bacterin.

Photobacteriosis

The disease is also known as pseudotuberculosis (formerly pasteurellosis) which is caused by *Photobacterium damsela* subsp. *piscicida*. The causative agent of the disease was first observed in white perch (*Morone americanus*) and striped bass (*M. saxatilis*) in Chesapeake Bay, USA (Snieszko and

Bullock, 1965). Until now, the bacteria has been isolated from diseased fish in England (Ajmal and Hobbs, 1967), Norway (Hastein and Bullock, 1976), Taiwan (Toranzo *et al.*, 1989), Spain (Toranzo *et al.*, 1991), Greece (Baudin *et al.*, 1991) and Italy (Ceschia *et al.*, 1991).

Photobacterium damsela subsp. piscicida has been observed in numerous fish species in Turkey (Table 5). It was first isolated from diseased sea bream in Aegean Sea (Cagirgan, 1993a). Subsequent isolations were reported from sea bass (Candan, 1996), *Mugil sp.* (Tanrikul and Cagirgan, 2001) and rainbow trout (Savas and Ture, 2008).

Flavobacteriosis

There are three main *Flavobacterium spp.* that are primary pathogen of freshwater fish within the genus. These are; *F. psychrophilum*, the causative agent of cold-water disease, *F. columnare*, the causative agent of columnaris disease and *F. branchiophilum*, the causative agent of bacterial gill

disease (BGD). Other *Flavobacterium* species including *F. scophthalmum*, *F. balustinum*, *F. hydatis*, *F. johnsoniae* and *F. oncorhynchi* were also recognized as a fish pathogens (Austin and Austin, 2012). Among them, *F. psychrophilum*, *F. columnare*, *F. branchiophilum*, *F. johnsoniae* and *F. hydatis* were identified in Turkey (Table 6).

Bacterial Cold-water Disease (BCD)

Flavobacterium psychrophilum (*Cytophaga psychrophilum* or *Flexibacter psychrophilum*) is the etiological agent of the rainbow trout fry syndrome (RTFS) and also known as the causative agent of bacterial cold-water disease in larger cultured fish. The disease had this name because it particularly occurs in low temperatures when water temperature decrease to below 12°C. The first outbreak occurred in coho salmon and isolated from diseased fish in USA in 1948 (Lehmann *et al.*, 1991). RTFS has been confirmed in many countries. During last twenty years, it had economically devastating impact on

Table 4. Distribution and host range of *Yersinia ruckeri*

Common name of the disease	Etiological agent	Host	Geographical distribution	References
		<i>Oncorhynchus mykiss</i>	Mediterranean Region	Timur and Timur (1991b)
		<i>Oncorhynchus mykiss</i>	Marmara Region	Karatas and Candan (1997)
		<i>Oncorhynchus mykiss</i>	Mugla	Diler <i>et al.</i> (1998)
		<i>Oncorhynchus mykiss</i>	Aydin	Kirkan <i>et al.</i> (2000)
		<i>Oncorhynchus mykiss</i>	Black Sea	Karatas <i>et al.</i> (2004)
		<i>Oncorhynchus mykiss</i>	Malatya	Seker <i>et al.</i> (2005)
		<i>Dicentrarchus labrax</i>	Ordu	Savaş <i>et al.</i> (2006)
		<i>Oncorhynchus mykiss</i>	Erzurum	Saglam <i>et al.</i> (2006)
Enteric red mouth	<i>Yersinia ruckeri</i>	<i>Oncorhynchus mykiss</i>	Adana, Antalya, Burdur, Isparta Mersin and Kahramanmaraş	Mefut <i>et al.</i> (2007)
		<i>Oncorhynchus mykiss</i>	Elazig	Kilic <i>et al.</i> (2007)
Yersiniosis		<i>Dicentrarchus labrax</i>	Eastern Black Sea	Savas and Ture (2008)
		<i>Oncorhynchus mykiss</i>	Mugla, Aydin, Denizli	Aksit and Kum (2008)
		<i>Oncorhynchus mykiss</i>	Mersin	Ozer <i>et al.</i> (2008b)
		<i>Oncorhynchus mykiss</i>	Isparta, Bilecik, Denizli, Kirkklareli Bolu, Tekirdag, Canakkale, Elazig Aydin, Bursa, Yalova, Trabzon, Rize Kutahya, Kayseri, Urfa, Ordu Manisa, Denizli	Altun <i>et al.</i> (2010)
		<i>Oncorhynchus mykiss</i>	Malatya and Elazig	Seker <i>et al.</i> (2012)

Table 5. Distribution and host range of *Ph. damsella subsp. P. piscicida*

Common name of the disease	Etiological agent	Host	Geographical distribution	References
Photobacteriosis	<i>Photobacterium damsella subsp. piscicida</i>	<i>Sparus aurata</i>	Aegean Sea	Cagirgan (1993b)
		<i>Dicentrarchus labrax</i>	Canakkale	Candan <i>et al.</i> (1996)
Pseudotuberculosis	<i>Pasteurella piscicida</i>	<i>Mugil sp.</i>	Aegean Sea	Tanrikul and Cagirgan (2001)
		<i>Dicentrarchus labrax</i>	Aegean Region	Turk (2002)
Pasteurellosis	<i>Pasteurella piscicida</i>	<i>Dicentrarchus labrax</i>	Antalya	Korun (2004b)
		<i>Oncorhynchus mykiss</i>	Eastern Black Sea	Savaş <i>et al.</i> (2006)
		<i>Dicentrarchus labrax</i>	Eastern Black Sea	Savaş <i>et al.</i> (2006)

Table 6. Distribution and host range of *Flavobacterium* species

Common names of the disease	Etiological agents	Host	Geographical distribution	References
Cold-water disease	<i>Flavobacterium psychrophilum</i>	<i>Oncorhynchus mykiss</i>	Aydın, Canakkale, Bilecik, Manisa, Samsun, Ordu, Kayseri,	Balta and Cagırgan (1998)
		<i>Oncorhynchus mykiss</i>	Sakarya	Korun and Timur (2001)
		<i>Oncorhynchus mykiss</i>	Mediterranean Region	Diler <i>et al.</i> (2003)
		<i>Oncorhynchus mykiss</i>	Marmara Region	Timur <i>et al.</i> (2004)
Rainbow trout fry syndrome (RTFS)	<i>Flavobacterium psychrophilum</i>	<i>Oncorhynchus mykiss</i>	Elazığ, Malatya, Erzincan	Ispir <i>et al.</i> (2004)
		<i>Dicentrarchus labrax</i>	Mugla	Ayaz (2006)
		<i>Oncorhynchus mykiss</i>	Aegean Sea	Kum <i>et al.</i> (2008)
		<i>Oncorhynchus mykiss</i>	Trabzon, Rize	Kayis <i>et al.</i> (2009)
Columnaris disease	<i>Flavobacterium columnare</i> (<i>Flexibacter columnaris</i> , <i>Cytophaga columnaris</i>)	<i>Oncorhynchus mykiss</i>	Rize, Trabzon, Aydın, Kayseri, Ordu, Manisa, Denizli	Balta and Cagırgan (1998)
Saddleback disease		<i>Oncorhynchus mykiss</i>	Isparta	Kubilay <i>et al.</i> (2008)
Cotton wool disease		<i>Oncorhynchus mykiss</i>	Mugla	Kubilay <i>et al.</i> (2009)
Mouth fungus		<i>Oncorhynchus mykiss</i>	Mersin	Yildirim and Ozer (2010)
Gill and skin disease	<i>Flavobacterium johnsoniae</i>	<i>Oncorhynchus mykiss</i>	Mersin	Yildirim and Ozer (2010)
		<i>Acipenser gueldenstaedtii</i>	Sakarya	Karatas <i>et al.</i> (2010)
Bacterial gill disease (BGD)	<i>Flavobacterium branchiophilum</i>	<i>Oncorhynchus mykiss</i>	Mersin	Yildirim ve Ozer (2010)
-	<i>Flavobacterium hydatis</i>	<i>Acipenser gueldenstaedtii</i>	Sakarya	Timur <i>et al.</i> (2010)

cultivated fish, especially on rainbow trout in Europe and considered as a serious disease in farmed rainbow trout fry and fingerlings (Austin and Stobie, 1992; Bernardet *et al.*, 1988; Santos *et al.*, 1992; Toranzo and Barja, 1993).

The pathogen was confirmed in most of the rainbow trout hatcheries in Turkey and caused heavy mortalities. First isolation of *F. psychrophilum* was from rainbow trout which sampled from different farms in 1990 (Balta and Cagırgan, 1998) (Table 6). Especially, when fish are small (0.1-4 g) and water temperature decrease to below 10°C, BCD or RTFS causes high mortality.

Columnaris Disease

Causative agent of the columnaris disease was described as *Flavobacterium columnare* (formerly *Flexibacter columnaris*, *Cytophaga columnaris*, *Chondrococcus columnaris*). It also referred to as cotton wool disease, saddleback disease, or mouth

fungus. The disease was first reported from different fish species in Mississippi River and named *Bacillus columnaris* (Davis, 1922). After two decades, Ordal and Rucker (1944) isolated and described the causative agent. Unlike cold-water disease, columnaris disease generally occurs when the water temperature gets above 15°C.

Although *F. columnare* was isolated in Aegean, Mediterranean and Black Sea regions of Turkey, initial isolation of columnaris disease was from farmed rainbow trout in Mugla (Balta and Cagırgan, 1998) (Table 6).

Bacterial Gill Disease

Flavobacterium branchiophilum is the causative agent of the Bacterial Gill Disease (BGD). It was first described in 1926 (Davis, 1926). Afterwards, it was isolated from salmonids in USA (Kimura *et al.*, 1978). Subsequently it was also isolated in Korea (Ko and Heo, 1997), Canada (Ostland *et al.*, 1994) and

Japan (Wakabayashi *et al.*, 1980).

In Turkey, it was claimed that the pathogen was isolated from rainbow trout hatcheries and caused mortalities up to %50 with other *Flavobacterium spp.* (Yildirim and Ozer, 2010) (Table 6). However, some of the biochemical test results reported in this particular study does not match with *F. branchiophilum* (Austin and Austin, 2012; Plumb and Hanson, 2011; Woo and Bruno, 2011). Therefore, there might be a misidentification of *F. branchiophilum*.

Other *Flavobacterium spp.*

Two other *Flavobacterium* species were also isolated from diseased fish in Turkey. One of them is *F. johnsoniae* which causes disease in gills and skin. The etiological agent was first isolated from Russian sturgeon (*Acipenser gueldenstaedtii*) in Sakarya (Karatas *et al.*, 2010). Diseased sturgeons were characterized by hemorrhages and erosions in the ventral part of the body. At the same year, pathogen was also reported from rainbow trout cultured in fish farm, located in Mersin (Yildirim and Ozer, 2010). The other *Flavobacterium* species is *F. hydatis*. It was also first isolated from Russian sturgeon in Sakarya (Table 6). Mixed infection with *F. hydatis* and *A. hydrophila* caused low mortalities in sturgeon (Timur *et al.*, 2010). Findings suggest that *F. hydatis* might be playing a role as secondary pathogen.

Flexibacteriosis

Flexibacteriosis is a severe disease caused by *Tenacibaculum maritimum* (formerly *Flexibacter maritimus*). Several other names, such as tenacibaculosis, marine columnaris, eroded mouth syndrome, black patch necrosis, and gliding bacterial disease were used to designate the disease caused by this bacterium. It has been recognized as a pathogen of fish since 1979 (Hikida *et al.*, 1979). The pathogen has been isolated in Australia (Handler *et al.*, 1997), USA (Chen *et al.*, 1985), Canada (Ostland *et al.*, 1999) and Europe (Pazos *et al.*, 1996). Many fish species would appear to be susceptible to infections by *T. maritimum*. To date, it has been isolated from sea bass, Dover sole (*Solea solea*), turbot (*Scophthalmus maximus*), salmonids, striped trumpeter (*Latris lineata*), greenback flounder (*Rhombosolea tapiriña*), sardine (*Sardinops sagax*), anchovy (*Engraulis mordax*) (Santos *et al.*, 1999).

The first report of *T. maritimum* infection in Turkey was in sea bass and sea bream from Aegean Region (Turk, 2002). Timur *et al.* (2011) isolated the pathogen from Black Sea Region from rainbow trout.

Coccal Infections

Streptococcosis stands for a common name of a disease caused by different genera and species. The

disease which is known as "warm water" streptococcosis (causes morbidity and mortality above 15°C) is caused by *Lactococcus garvieae*, *Streptococcus iniae*, *Streptococcus parauberis* and *Streptococcus agalactiae*. The other one, vagococcosis, (causes morbidity and mortality below 15°C) is caused by *Vagococcus salmoninarum*. Hoshina *et al.* (1958) was reported *Streptococcus spp.* as a fish pathogen in 1950s in Japan.

Lactococcus garvieae (formerly *Enterococcus seriolicida*) was initially isolated from a trout farm in 1988 in Spain (Palacios *et al.*, 1993). Subsequently, it was reported from many other countries such as, South Africa (Casrson *et al.*, 1993), Italy (Ghittino and Prearo, 1992), Iran (Soltani *et al.*, 2008) and USA (Evas *et al.*, 2009). *L. garvieae* was first isolated from rainbow trout in Aegean region of Turkey (Diler *et al.*, 2002). Subsequently, the outbreaks have been reported from rainbow trout, turbot and sea bass (Table 7) in different parts of the country.

Vagococcus salmoninarum was first isolated from diseased rainbow trout in USA in 1968 and named as lactobacillus (Austin and Austin, 2012). In Turkey, Didinen *et al.* (2011) was performed the first isolation from rainbow trout farmed in Mediterranean region. No following outbreak or isolation was reported since.

Staphylococcus epidermidis was first reported from farmed yellowtail and red sea bream in Japan (Kusuda and Sugiyama, 1981). In Turkey, initial isolation was performed by Turk (2002) from sea bass cultured in cage, located at the coastal region of the Aegean Sea. Subsequently, Timur *et al.* (2008) proved the existence of the pathogen in Turkey by isolating it from sea bass in cage farm, located in the coastal region of the Black Sea. *Staphylococcus cohnii subsp. cohnii* was first isolated from farmed rainbow trout and common dentex (*Dentex dentex*) in Aegean region of Turkey (Akayli *et al.*, 2011).

Pseudomoniasis

Pseudomonas infections or Pseudomoniasis refer to a disease caused by *Pseudomonas* species. *Pseudomonas spp.* are found in normal microbial flora of both freshwater and saltwater fish. It was believed that these bacteria could be opportunistic pathogens. Most of the time, *Pseudomonas spp.* isolated with other bacteria. For instance, when rainbow trout is infected with *Y. ruckeri*, both *Ps. pseudoalcaligenes* and *Y. ruckeri* were isolated from the fish (Austin and Stobie, 1992). Findings suggest that *Pseudomonas spp.* can be secondary infections. Confirmed pathogenic species of this genus are *Ps. chlororaphis*, *Ps. anguilliseptica*, *Ps. fluorescens*, *Ps. putida*, *Ps. plecoglossicida*, *Ps. aeruginosa* and *Ps. luteola* (Muroga and Nakajima, 1981; Prosyanyaya, 1981; Toranzo and Barja, 1993; Altınok *et al.*, 2006; Kayis *et al.*, 2009). Among these pathogens, *Ps. anguilliseptica* is considered the most significant

Table 7. Distribution and host of some gram positive pathogens

Common names of the disease	Etiological agent	Host	Geographical distribution	References
Lactococcosis	<i>Lactococcus garvieae</i>	<i>Oncorhynchus mykiss</i>	-	Cagırgan and Tanrikul (1995)
		<i>Oncorhynchus mykiss</i>	Aegean Region	Diler <i>et al.</i> (2002)
		<i>Oncorhynchus mykiss</i>	Mugla, Denizli, Antalya, Konya	Altun <i>et al.</i> (2005)
		<i>Oncorhynchus mykiss</i>	Konya, Antalya, Isparta,	Kubilay <i>et al.</i> (2005)
		<i>Psetta maxima</i>	Eastern Black Sea	Savas and Ture (2008)
		<i>Oncorhynchus mykiss</i>	Eastern Black Sea	Savas and Ture (2008)
		<i>Dicentrarchus labrax</i>	Eastern Black Sea	Savas and Ture (2008)
		<i>Oncorhynchus mykiss</i>	Aydin, Mugla, Denizli	Aksit and Kum (2008)
		<i>Oncorhynchus mykiss</i>	Mersin	Ozer <i>et al.</i> (2008b)
"Warm-water" Streptococcosis		<i>Oncorhynchus mykiss</i>	Marmara Region	Timur <i>et al.</i> (2011)
		<i>Oncorhynchus mykiss</i>	Kutahya, Bilecik, Isparta,	
		<i>Oncorhynchus mykiss</i>	Bursa, Samsun, Mugla, Antalya	Altun <i>et al.</i> (2013b)
Vagococcosis, "Cold-water" Streptococcosis	<i>Vagococcus salmoninarum</i>	<i>Oncorhynchus mykiss</i>	Mediterranean Region	Didinen <i>et al.</i> (2011)
Staphylococcosis	<i>Staphylococcus epidermidis</i>	<i>Dicentrarchus labrax</i>	Aegean Region	Turk (2002)
		<i>Sparus aurata</i>	Aegean Region	Kubilay and Ulukoy (2004)
Staphylococcosis	<i>Staphylococcus cohnii subsp. cohnii</i>	<i>Dicentrarchus labrax</i>	Black Sea	Timur <i>et al.</i> (2008)
		<i>Dentex dentex</i>	Aegean Region	Akayli <i>et al.</i> (2011)
		<i>Oncorhynchus mykiss</i>	Aegean Region	Akayli <i>et al.</i> (2011)

pathogen for fish and causes winter disease when the water temperature gets below 15°C. *Ps. anguilliseptica* was first reported from Japanese eel in Japan in 1972 (Wakabayashi and Egusa, 1972). It mainly affects Japan and European eels, causes red spot disease (also known as sekiten-byo) (Plumb and Hanson, 2011). In Turkey, initial isolation of this pathogen was performed by Turk (2002) from sea bass from Aegean Sea. Recognized hosts of the disease are sea bass and sea bream.

Ps. fluorescens is the causative agent of the pseudomonas septicemia. As it was mentioned earlier, it is an opportunistic pathogen. It shows similar clinical signs of *Aeromonas* septicemia. In Turkey, the pathogen was first reported from sea bream in Aegean Sea (Turk, 2002). Afterwards, it was also isolated from sea bass, rainbow trout and from some ornamental fish species (Table 8).

The first outbreak of a disease caused by *Ps. luteola* was recorded by Altınok *et al.* (2007) in cultured rainbow trout. To date, it is the only report implicating *Ps. luteola* as a fish pathogen in the world. Diseased fish were externally characterized by exophthalmia, dark pigmentation, and ulcerative lesions near dorsal fin.

Ps. aeruginosa is known as a causative agent of gill disease. It was reported from sea bass in Aegean Sea and rainbow trout in Mersin, Trabzon and Rize (Ayaz, 2006; Kayis *et al.*, 2009). *Ps. chlororaphis* was first reported from sea bream in Aegean Sea but biochemical properties of the isolate was not mentioned (Turk, 2002). *P. plecoglossicida* was first reported from rainbow trout fries in Turkey (Akayli *et*

al., 2010). *P. putida* causes ulcerative infection in fish. It was first isolated from common carp in Turkey (Aydin *et al.*, 1998). It was also isolated from rainbow trout and sea bass (Table 8).

Mycobacteriosis

Mycobacteriosis (Fish Tuberculosis) is severe disease known to affect wide range of freshwater and saltwater fish. Primary etiological agent of the disease is *Mycobacterium marinum* but other *Mycobacterium spp.* were also described as a fish pathogen (Bragg *et al.*, 1990. Chinabut, 1999; Talaat *et al.*, 1999). It was reported initially by Aronson (1926) from tropical aquarium fish in USA.

Mycobacterium can be found in the water and sediment and it has worldwide distribution. Until now, two outbreaks of the mycobacteriosis from different hosts were reported in Turkey. Candan (1991a) was first who mentioned existence of *Mycobacterium sp.* in sea bream in Mugla. Pathogen was diagnosed as a *Mycobacterium sp.* based on clinical signs of the disease and no biochemical or other tests were carried out to identify bacteria.

Korun *et al.* (2005) was claimed the first isolation of mycobacteriosis. The outbreak of the disease was occurred in sea bass farm located in Aegean coast of Canakkale. The etiological agent of the disease was reported as a *Mycobacterium sp.* Acid-fast rods in granulomas were seen in histological section but authors failed to isolate the pathogen.

Piscirickettsiosis

Table 8. Distribution and host range of *Pseudomonas* spp

Common names of the disease	Etiological agents	Host	Geographical distribution	References
Haemorrhagic septicaemia, Fin Rot	<i>Pseudomonas</i> sp.	<i>Stizostedion lucioperca</i>	Eğirdir	Timur and Timur (1985)
		<i>Sparus aurata</i>	Aegean Region	Turk (2002)
	<i>Pseudomonas fluorescens</i>	<i>Dicentrarchus labrax</i>	Aegean Region	Turk (2002)
		<i>Oncorhynchus mykiss</i>	Sakarya	Akayli and Timur (2004)
		<i>Poecilia reticulata</i>	Istanbul (Aquarium)	Akayli and Korun (2004)
		<i>Oncorhynchus mykiss</i>	Ankara	Ozkok (2005)
		<i>Dicentrarchus labrax</i>	Adana	Matyar (2007)
		<i>Oncorhynchus mykiss</i>	Trabzon, Rize	Kayis <i>et al.</i> (2009)
Ulcerative disease, Haemorrhagic ascites	<i>Pseudomonas putida</i>	<i>Cyprinus carpio</i>	-	Aydin <i>et al.</i> (1998)
		<i>Carassius auratus</i>	-	Aydin <i>et al.</i> (1998)
		<i>Oncorhynchus mykiss</i>	Ankara	Ozkok (2005)
		<i>Oncorhynchus mykiss</i>	Trabzon	Altınok <i>et al.</i> (2006)
		<i>Dicentrarchus labrax</i>	Adana	Matyar (2007)
Red spot disease, Sekiten-byo	<i>P. anguilliseptica</i>	<i>Dicentrarchus labrax</i>	Aegean Sea	Turk (2002)
		<i>Sparus aurata</i>	Izmir, Mugla	Canak (2011)
Gill disease	<i>P. aeroginasa</i>	<i>Dicentrarchus labrax</i>	Mugla	Ayaz (2006)
		<i>Oncorhynchus mykiss</i>	Mersin	Yildirim and Ozer (2010)
		<i>Oncorhynchus mykiss</i>	Trabzon, Rize	Kayis <i>et al.</i> (2009)
-	<i>P. chlororaphis</i>	<i>Dicentrarchus labrax</i>	Aegean Region	Turk (2002)
Haemorrhagic ascites	<i>P. plecoglossicida</i>	<i>Oncorhynchus mykiss</i>	-	Akayli <i>et al.</i> (2010)
-	<i>P. luteola</i>	<i>Oncorhynchus mykiss</i>	Trabzon	Altınok <i>et al.</i> (2007)

Piscirickettsiosis is a bacterial infection that caused by *Piscirickettsia salmonis*. The disease was also known as salmonid rickettsial septicemia. It primarily affects salmonids reared in sea cages. The disease has been known since 1939 (Plumb and Hanson, 2011). After half decade of piscirickettsiosis outbreaks, it has been observed in cultured coho salmon in Chile and caused heavy mortalities (Bravo and Campos, 1989). Subsequently, the bacterium was isolated from Atlantic salmon in Norway, Canada and Ireland (OIE, 2013; Olsen *et al.*, 1997; Rodger and Drinan, 1993) and tilapia in Taiwan (Chern and Chao, 1994).

During 2003, first outbreak of the disease occurred in sea bass farm on the coast of the Black Sea in Turkey. Isolated bacterium was described as rickettsia-like organism (RLO) (Timur *et al.*, 2005). Different pathogen species, *L. anguillarum* and *P. damsela* subsp. *piscicida* were also isolated from examined fish. To date, it is the only confirmed report of the pathogen in Turkey.

Bacterial Kidney Disease (BKD)

Bacterial kidney disease is a chronic disease that causes high mortalities particularly in salmonids. Etiological agent of the disease is *Renibacterium salmoninarum* (syn. *Corynebacterium salmonis*) which targets the fish kidney and causes white spots.

Thus, it was named as kidney disease. The disease is generally observed in salmonids and first reported in Atlantic salmon in 1990 in Scotland (Austin and Austin, 2012), but etiological agent was first isolated in 1956 (Ordal and Earp, 1956). Subsequently, BKD occurred in Finland (Rimaila, 2002), Japan (Kimura and Awakura, 1977) and Canada (Evelyn *et al.*, 1973).

In Turkey, first attempt was made to isolate *R. salmoninarum* from diseased rainbow trout in Bayindir Dam, Ankara (Halici *et al.*, 1977) but biochemical tests used to isolate the bacterium did not match with *R. salmoninarum*. Therefore, it should be misidentification of *R. salmoninarum*. Later, mixed colonies of *Y. ruckeri* and *R. salmoninarum* were isolated from cultured Black Sea salmon (*Salmo trutta labrax*) in the Eastern Black Sea Region and results were confirmed by sequencing 501 base pairs of DNA fragment from 16s rDNA gene of the *R. salmoninarum* (Savas *et al.*, 2006).

Other Bacterial Fish Pathogens

Some bacteria are well-studied by scientists and known as fish pathogens. Most of the bacteria classified as well-recognized pathogens are economically important and causes heavy mortalities or has a wide host range and distribution. For other bacterial fish pathogens there are limited information available. Whenever there is limited information

available, it could be possible that either the disease is new emerging disease or it is rarely encountered. It is also possible that some of them are recognized as secondary or opportunistic pathogens and unusual stress factors may play a key role in their occurrence. Bacteria, which were reported as a fish pathogens in Turkey without adequate information were also discussed (Table 9).

Edwardsiella Infections (Enteric Septicemia)

Two species of genus *Edwardsiella* are recognized as fish pathogens. These are *E. ictaluri* and *E. tarda*. *E. tarda* is not only a fish pathogen but also human pathogen. *E. tarda* causes red disease of eels and fish gangrene of catfish. The disease was first isolated from cultured eel in Asia (Hoshina, 1962). Edwardsiellosis caused by *E. tarda* which causes serious mortality in marine and freshwater fish, including catfish, carp, eel, flounder, seabream tilapia,

and yellowtail (Plumb and Hanson, 2011). It has not been isolated from any fish species in the Turkey.

On the other hand, *E. ictaluri* is the causative agent of enteric septicemia of catfish or hole in the head disease was first reported in catfish in USA (Hawke *et al.*, 1981). Subsequently, the disease, was observed in Australia (How *et al.*, 1983) and Taiwan (Chung and Kou, 1983).

E. ictaluri was first isolated from rainbow trout in Ankara in Turkey (Keskin *et al.*, 2004), however, infected fish didn't show any clinical sign. To date, it remains as the only *E. ictaluri* isolation in Turkey (Table 9).

Ph. damsela subsp. *damsela* Infection

The disease caused by *Photobacterium damsela* subsp. *damsela* (Formerly known as *Vibrio damsela*) is skin ulcer. It was initially isolated from blacksmith (*Chromis punctipinnis*) (Love *et al.*, 1981). The

Table 9. Distribution and host range of other bacterial pathogens

Common name of the disease	Etiological agent	Host	Geographical distribution	References
-	<i>Arcobacter cryaerophilus</i> (<i>Campylobacter cryaerophila</i>)	<i>Oncorhynchus mykiss</i>	Marmara Region	Bulut and Aydin (2004)
-	<i>Aerococcus viridans</i>	<i>Oncorhynchus mykiss</i>	Mersin	Ozet <i>et al.</i> (2008)
-	<i>Burkholderia cepacia</i>	<i>Oncorhynchus mykiss</i> <i>T. mediterraneus</i>	Trabzon, Rize Trabzon	Kayis <i>et al.</i> (2009) Boran <i>et al.</i> (2013)
Necrotic kidney, Hyperemia of the mouth	<i>Citrobacter braakii</i>	<i>Oncorhynchus mykiss</i>	Bursa	Altun <i>et al.</i> (2013c)
-	<i>Citrobacter freundii</i>	<i>Oncorhynchus mykiss</i> <i>Oncorhynchus mykiss</i>	Erzurum Trabzon, Rize	Saglam <i>et al.</i> (2006) Kayis <i>et al.</i> (2009)
Enteric septicemia of catfish, hole in the head disease	<i>Edwardsiella ictaluri</i>	<i>Oncorhynchus mykiss</i>	Ankara	Keskin <i>et al.</i> (2004)
	<i>Enterobacter cloacae</i>	<i>Oncorhynchus mykiss</i>	Trabzon, Rize	Kayis <i>et al.</i> (2009)
Septicemia	<i>Escherichia vulneris</i>	<i>Oncorhynchus mykiss</i> , <i>Carassius Carassius</i> , <i>Poecilia sp.</i>	Erzurum	Aydin <i>et al.</i> (1997)
Hemorrhagic septicemia	<i>Hafnia alvei</i>	<i>Oncorhynchus mykiss</i> <i>Oncorhynchus mykiss</i>	Ankara Trabzon, Rize	Ozkok (2005) Kayis <i>et al.</i> (2009)
Vibriosis	<i>Photobacterium damsela</i> subsp. <i>damsela</i> (<i>Vibrio damsela</i>)	<i>Sparus aurata</i> <i>Dicentrarchus labrax</i> <i>Dicentrarchus labrax</i> <i>Oncorhynchus mykiss</i> <i>Oncorhynchus mykiss</i> <i>T. mediterraneus</i>	Mugla Aegean Region Antalya Black Sea Trabzon, Rize Trabzon	Akayli (2001) Turk (2002) Korun (2004b) Timur <i>et al.</i> (2008) Kayis <i>et al.</i> (2009) Boran <i>et al.</i> (2013)
-	<i>Plesiomonas shigelloides</i>	<i>Poecilia reticulata</i> <i>Oncorhynchus mykiss</i>	Istanbul (Aquarium) Trabzon, Rize	Akayli and Zeybek (2005) Kayis <i>et al.</i> (2009)
Septicemia	<i>Serratia liquefaciens</i>	<i>Oncorhynchus mykiss</i> <i>Oncorhynchus mykiss</i>	Erzurum Trabzon, Rize	Aydin <i>et al.</i> (2001) Kayis <i>et al.</i> (2009)
Ulcer, Hemorrhage	<i>Shewanella putrefaciens</i> grp	<i>Dicentrarchus labrax</i> <i>Oncorhynchus mykiss</i> <i>Carassius auratus auratus</i>	Aegean Region Trabzon, Rize Bursa (Aquarium)	Korun <i>et al.</i> (2009) Kayis <i>et al.</i> (2009) Altun <i>et al.</i> (2013a)

bacterium is not only a fish pathogen but also human pathogen. It effects variety of marine fish spp. Damsel fish seems to be most susceptible to this pathogen.

It is one of the most important diseases in cultured European sea bass (Korun, 2004b). Initial isolation was fulfilled from sea bream in Mugla, Turkey (Akayli, 2001). In same region, it was also isolated from sea bass (Turk, 2002). Subsequently, the pathogen was reported from rainbow trout and horse mackerel in Black Sea (Timur *et al.*, 2008; Kayis *et al.*, 2009; Boran *et al.*, 2013) (Table 9). It should be noted that only mixed bacteria colonies were isolated in all of these disease cases. Thus, pathogenicity of the bacterium is unclear in Turkey and the causative agent probably plays secondary pathogen role in most of the cases.

***Plesiomonas shigelloides* Infection**

Plesiomonas shigelloides was first isolated from rainbow trout in Portugal in 1984 (Cruz *et al.*, 1986). Subsequently, pathogen also isolated from catfish, sturgeon, rainbow trout, eel and gourami in Germany (Klein *et al.*, 1993). Hemorrhages around the vent and protruded anus generally occur after infection (Austin and Austin, 2012). The pathogen is also the etiological agent of diarrhea in humans (Klein *et al.*, 1993).

Akayli and Zeybek (2005) was first reported the isolation of *P. shigelloides* from guppies (*Poecilia reticulata*) which were collected from different aquarium stores. After four years, the pathogen was reported from rainbow trout in Trabzon (Kayis *et al.*, 2009) (Table 9).

***Serratia liquefaciens* Infection**

Serratia liquefaciens is recognized as a potential fish pathogen belonging to *Enterobacteriaceae* family and it is the causative agent of bacterial septicemia (Woo and Bruno, 2011). The bacterium was isolated in different countries, for instance, from turbot in France (Vigneulle and Baudin, 1995), arctic char (*Salvelinus alpinus*) in U.S.A (Starliper, 2001) and from Atlantic salmon in Scotland (McIntosh and Austin, 1990).

It was first isolated from rainbow trout in 2001 in Erzurum, Turkey (Aydin *et al.*, 2001). Following isolation was reported in Trabzon from rainbow trout (Kayis *et al.*, 2009). So far, the pathogen has never associated with serious outbreaks in Turkey (Table 9).

***Citrobacter* Infection**

Citrobacter freundii was first reported as an emerging fish pathogen from sunfish (*Mola mola*) in Japan (Sato *et al.*, 1982). After initial isolation, it was also isolated from Atlantic salmon in USA (Baya *et al.*, 1990), rainbow trout in Spain which caused high mortalities with IPNV (Sanz, 1991) and from carp in

India (Karunasagar *et al.*, 1992). *C. freundii* is an opportunistic pathogen. Thus, stress, and pollution play a key role in infection occurrence (Sanz, 1991; Whalen *et al.*, 2007).

In Turkey, it was isolated from rainbow trout in Erzurum (Saglam *et al.*, 2006) but there is no clear information about bacterial pathogenicity. After few years, it was also isolated from rainbow trout in Trabzon (Kayis *et al.*, 2009). In both events, no heavy mortalities were observed. Besides that, another *Citrobacter* species, *Citrobacter breakii* was first reported as a fish pathogen in Bursa (Altun *et al.*, 2013c) (Table 9).

***Arcobacter cryaerophilus* Infection**

The bacterium is part of the normal micro flora of freshwater fish (Aydin *et al.*, 2000b). Since the bacterium is ubiquitous in the environment and fish, it is not surprising that it might be opportunistic fish pathogen. It was first described as an emerging fish pathogen and isolated from rainbow trout in Balıkesir and Canakkale (Aydin *et al.*, 2000). After that, pathogenicity of the bacterium was confirmed by experimental infections which were performed on rainbow trout, and albino crosses (Bulut and Aydin, 2004) (Table 9). Infected fish showed pale gills, watery spleens, hemorrhages and bloody fluid in intestine.

***Shewanella putrefaciens* Infection**

Disease caused by *S. putrefaciens* was first occurred in farmed rabbit fish (*Siganus rivulatus*) in Red Sea in 1985 and caused heavy mortalities (Saeed *et al.*, 1987). After initial report, it was isolated from both common carp (*Cyprinus carpio*) and rainbow trout farms located in Poland (Kozinska and Pekala, 2004).

Korun *et al.* (2009) was first reported the outbreak of a disease caused by *S. putrefaciens* from European sea bass in Aegean Region of Turkey. In the same year, the bacterium was also reported from rainbow trout in Trabzon (Kayis *et al.*, 2009) (Table 9).

Aerococcus viridans

The bacterium was described as a pathogen and caused low mortalities in lobsters (*Homarus gammarus* L.) (Gjerde, 1984). Ozer *et al.* (2008) isolated *A. viridans* from cultured rainbow trout in Mersin (Table 9). Fish infected with *A. viridans* showed darkened skin, exophthalmia and hemorrhages in liver.

Hafnia alvei

Hafnia alvei is one of the causative agents of the hemorrhagic septicemia. It was first reported as a fish

pathogen from rainbow trout in Bulgaria (Gelev and Gelev, 1988). Subsequent outbreak occurred in cherry salmon (*O. masou*) in Japan and confirmed its pathogenicity. If API Biochemical Test System is used for identification, the bacterium could be misidentified as *Y. ruckeri* or vice versa. The pathogen was isolated from rainbow trout in Ankara, Rize and Trabzon (Ozkok, 2005; Kayis *et al.*, 2009) (Table 9).

Enterobacter cloacae

Enterobacter cloacae was isolated from mullet (*Mugil cephalus*) in India. Bath challenge was performed and pathogenicity was determined (Sekar *et al.*, 2008). In Turkey, Kayis *et al.* (2009) recovered the disease organism from rainbow trout in Trabzon and Rize (Table 9).

Viral Fish Diseases

IPNV

The causative agent is Infectious pancreatic necrosis virus (IPNV) which belongs to *Birnaviridae* group. It is icosahedral in shape, unenveloped and has a double stranded RNA genome. IPN is an infectious systemic disease that has been recognized to have worldwide distribution in a wide range of fish. It causes high mortalities especially in fry and fingerling salmonids. The virus can be vertically and horizontally transmitted. Thus, the only way to get rid of the virus is the destruction of the stock (Olson and Thomas, 1994). Wood *et al.* (1955) was first described the IPNV.

In Turkey, first IPNV particles were reported from rainbow trout organs after histopathological examination (Timur *et al.*, 1993). After a decade, existence of IPNV was proved by RT-PCR (Candan, 2002). Subsequently, IPNV has been reported from different parts of Turkey and findings suggest wide distribution of the virus in Turkey (Table 10).

VHSV

Viral hemorrhagic septicemia (VHS) is a severe viral disease caused by VHS virus. The virus belongs to the family *Rhabdoviridae*, genus *Novirhabdovirus*. The disease was first occurred in rainbow trout in Germany (Schaperclaus, 1938). It is responsible for severe economic losses in rainbow trout farms in Europe (Woo and Bruno, 2011). VHSV was thought to have a predilection for salmonids, especially rainbow trout. Over the years, host range of the pathogen has expanded. For instance, Atlantic cod, pike, turbot, sea bass, Coho salmon, brown trout (Castric and Kinkelin, 1980; Jensen *et al.*, 1979; Meier and Vestergard, 1980; Rasmussen, 1965; Winton *et al.*, 1989). Unlike IPNV, there is no clear evidence about vertical transmission of the virus, but

it is likely that, it can be transmitted horizontally and in many cases survivors become asymptomatic carriers. Salmonid fries are the most susceptible fish species to the virus. Mortality rates can reach up to 80-90%.

VHSV was first isolated in Trabzon, Turkey from cultured turbot fry and brood stock in 2004 (Kalayci *et al.*, 2006; Nishizawa *et al.*, 2006). Subsequent isolations (Table 10) of VHSV in the Black Sea Region proved the existence of VHSV in Turkey (Altuntas and Ogut, 2010).

Viral Erythrocytic Necrosis (VEN)

The etiological agent of the disease is an iridovirus, named as erythrocytic necrosis virus (ENV). It was first reported in 1969 and identified as piscine erythrocytic necrosis (Laird and Bullock, 1969). Since its initial recognition, the virus has been recovered as a pathogen from a wide variety of fish species from different countries, for instance, sea bass in Spain (Pinto *et al.*, 1989), coho salmon in Japan (Takahashi *et al.*, 1982), pacific herring in Alaska (Meyers *et al.*, 1986), coho salmon in Chile (Reyes and Campalans, 1987) and eel in Taiwan (Chen *et al.*, 1985). Unlike VHSV and IPNV, VEN has not been recognized as a severe disease, it causes nominal mortalities in regard to other viral infections.

In 2008, viral erythrocytic infection occurred in cultured Mediterranean sea bass in Black Sea (Timur *et al.*, 2008). It is the only recorded report that has been made to date in Turkey (Table 10).

Lymphocystis

Lymphocystis is caused by a lymphocystis virus. It has been reported from wide variety of both freshwater and marine fish species from all over the world. It is one of the oldest and the best known fish virus. It has been known as a causative agent of a disease since 1874 (Plumb and Hanson, 2011). It was reported from ornamental fish in USA (Niggrelli and Ruggieri, 1965), Red Sea, Bering Sea, Mediterranean Sea (Anders, 1989) and Korea (Hossain *et al.*, 2007). The causative agent of the virus was isolated from Sea Bream in Mugla, Turkey (Candan, 1991b). There is not any report of severe disease outbreak associated with Lymphocystis in Turkey (Table 10).

Carp Pox

It is also known as a fish pox or epithelioma papillosum, caused by a herpes virus. Like lymphocystis, it is one of oldest known fish disease. The virus was first recorded in 15th century (Hedrick and Sano, 1989). It was mainly reported from carp producer countries such as European countries, United States and Far East countries. Sano *et al.* (1985) was reported that, the most susceptible hosts of the virus are common carp and koi carp.

Table 10. Distribution and host range of viral diseases

Virus	Abbreviation	Genome	Classification	Identified host	Geographical distribution	References
Infectious pancreatic necrosis virus	IPNV	dsRNA	<i>Birnaviridae</i> , <i>Aquabirnavirus</i>	<i>Oncorhynchus mykiss</i>	Different parts of Turkey	Candan (2002)
				<i>Oncorhynchus mykiss</i>	Ordu, Samsun, Tokat, Trabzon	Albayrak and Ozan (2010)
				<i>Oncorhynchus mykiss</i>	Eastern Black Sea	Ogut and Altuntas (2012)
				<i>Oncorhynchus mykiss</i>	Adana, Adiyaman, Ankara, Antalya, Artvin, Aydin, Bilecik, Bolu, Burdur, Denizli, Duzce, Elazig, Eskisehir, Gumushane, Hatay, Izmir, Kahramanmaraş, Kayseri, Mugla, Ordu, Sakarya, Samsun, Tokat, Trabzon, Usak, Zonguldak,	Kalayci <i>et al.</i> (2012)
				<i>Dicentrarchus labrax</i>	Antalya, Mugla, Ordu	Kalayci <i>et al.</i> (2012)
				<i>Oncorhynchus mykiss</i>	Elazig, Malatya, Sivas, Erzincan, Sanliurfa, Kahramanmaraş, Erzurum	Gurcay <i>et al.</i> (2013)
Viral haemorrhagic septicemia virus	VHSV	ssRNA	<i>Rhabdoviridae</i> , <i>Novirhabdovirus</i>	<i>Psetta maxima</i>	Trabzon	Nishizawa <i>et al.</i> (2006)
				<i>Scophthalmus maximus</i>	Trabzon	Kalayci <i>et al.</i> (2006)
				<i>Merlangius merlangus</i>	Eastern Black Sea	Altuntas and Ogut (2010)
				<i>Oncorhynchus mykiss</i>	Bolu	Kalayci <i>et al.</i> (2012)
				<i>Sparus aurata</i>	Izmir	Kalayci <i>et al.</i> (2012)
				<i>Dicentrarchus labrax</i>	Antalya, Mugla, Izmir	Kalayci <i>et al.</i> (2012)
Lymphocystis disease virus	-	dsDNA	<i>Iridoviridae</i> , <i>Lymphocystis virus</i>	<i>Psetta maxima</i>	Trabzon, Mugla	Kalayci <i>et al.</i> (2012)
				<i>Sparus aurata</i>	Mugla	Candan (1991b)
Erythrocytic necrosis virus	ENV	dsDNA	<i>Iridoviridae</i>	<i>Dicentrarchus labrax</i>	Black Sea	Timur <i>et al.</i> (2008)
Cyprinid herpes virus 1(Carp pox)	CHV-1	ssRNA	<i>Rhabdoviridae</i>	<i>Cyprinus carpio</i>	-	Timur (1991a)

Since it is restricted to cyprinids and carp production rates are low in Turkey, it is not considered as a serious disease problem in Turkey. It was first reported by Timur (1991a) from mirror carp in Antalya (Table 10).

Prevention and Treatment

The ideal way to control infectious fish diseases is to prevent exposure to pathogenic agents whenever possible, thus avoiding most devastating health problems through biosecurity (Plumb and Hanson, 2011). However, when dealing with the aquatic environment, it is virtually impossible to define all disease-causing agents and keep them isolated from the fish host. Water provides an excellent medium for transfer of many communicable agents from fish to fish or from locality to locality. Moreover, many disease-causing organisms are endemic to the aquatic environment and are opportunistic, facultative pathogens that remain viable under various conditions (Plumb and Hanson, 2011).

Aquaculture is a vital source of food and still the fastest growing animal food producing sector. As a result of intensive fish farming and stressful conditions, infectious diseases, especially bacterial and viral origins, became one of the most important

limiting factor in aquaculture facilities. Some diseases do not always reveal themselves in a clinical form. These types of diseases, e.g. BKD, furunculosis and ERM pose real risk of transferring the pathogen with fish movements (Hirvela *et al.*, 2006). Therefore, it is obligatory to apply transportation restrictions. These may prevent the spread of the diseases or slow down the transmission. It may require efficient monitoring, dissuasive and financial sanctions. In some instances, it is really hard to get rid of the disease. Complete disposition of infected fish and disinfection of facilities, especially hatcheries, may be more economical than the losses associated with mortalities. The best practice for the elimination of IPNV, VHSV, BKD and furunculosis is prevention by the use of disease-free stocks if possible.

Since fish are poikilothermic, they react more quickly to environmental change than homothermic terrestrial animals. Fish respond quickly any kind of environmental changes such as temperature change, excessive or insufficient dissolved gases in the water, metabolites, or chemical additives, and so forth, to which they are unable to adapt. These factors increase fish susceptibility to infectious agents and compromise their immune response (Plumb and Hanson, 2011).

In aquaculture, the usage of antimicrobial

compound was started in 1940s against furunculosis (Austin and Austin, 2012; Gutsell, 1946). Intensive fish farming is the main reason for the use of high amount of antimicrobial drugs. Incorrect usage of antimicrobials in veterinary medicine and in aquaculture as a growth promoter, prophylaxis and therapeutic purposes leads to bacterial resistance to the antibiotics and also, accumulation antibiotic residue in fish. Bacteria may develop resistance to antimicrobials if used too often, over an extended period of time, or if applied improperly. Oxytetracycline, sulfadimethoxine, tetracycline, tetracycline are among the most frequently used antibiotics (Plumb and Hanson, 2011). Consequently, most of the bacteria has already acquired resistance against these antibiotics (unpublished data). On the other hand, florfenicol is a newly introduced antibiotic to aquaculture (Kayis, 2009). Therefore most of the bacteria strains are susceptible against it. In Both in EU member states and the USA, limited antimicrobial agents are licensed for use in finfish and their use in aquaculture products that are for human consumption is very limited (Matyar, 2007). In contrast, in most of the developing countries, antimicrobial drug usage regulations are virtually absent, inadequate or unrestrained.

Several different kinds of antibiotics are used around the world in aquaculture for the control of bacterial diseases by adding them directly to the water or incorporating them into the feed. High frequencies of bacteria that are resistant to the antimicrobial agent have been found in aquaculture, including multiple resistant bacteria, found in fish farms and the surrounding aquatic environment. Accumulation of surplus antimicrobials and antimicrobial residues may occur in fish farms. Antimicrobial build up could establish selective pressure favoring selection and growth of antimicrobial-resistant bacteria. There is a potential risk that antimicrobial resistance genes could be disseminated into a wide range of aquatic environmental bacteria. There is also a possible flow of antimicrobial resistance genes between animal and human pathogens (Petersen *et al.*, 2002).

Initial effectiveness of using antibiotics against various fish diseases somewhat decreased the interest in vaccine development for particular diseases. However, with the emergence of antibiotic-resistant strains of bacteria as a result of antibiotics use has drawn significant attention back to vaccines. Under intensive rearing conditions, vaccines can provide protection against specific disease when fish are the most susceptible and provides long-term immunoprotection (Plumb and Hanson, 2011). Unlike antibiotics, vaccines do not leave any residue and they are safe for applied fish. Depending on the fish and environmental conditions, vaccination can be performed orally, by immersion or by injection. Like antibiotics, vaccines cannot totally eliminate all of the disease organism. Vaccinated fish can become carriers and can play role as a disease reservoir. That

might have an impact on wild stocks, which interact with vaccinated fish especially in sea cages.

Antibiotics are commonly used for treatment of these diseases but bacteria develop antibiotic resistant. To overcome these problems, fish should be vaccinated. Many attempts have been made to vaccinate fish against vibriosis and yersiniosis by using simple killed bacterin preparations. Immunization with killed bacteria has been attempted and protection obtained by immersion by feeding as a feed additive or by injection with the killed bacteria has been minimal. Better protection has been obtained by administering the killed bacteria by injection in combination with an emulsified adjuvant (Cane, 2013). However, immune responses to live vaccines are generally of greater magnitude and of longer duration than those produced by killed or subunit vaccines. Attenuated vaccines have some advantages over killed bacterin vaccines. They are living and invasive, thereby facilitating vaccine uptake and they establish low-grade infections resulting in the stimulation of cellular immunity and typically establish longer-lasting immunity (unpublished data).

Nowadays, live attenuated bacterial vaccines have been developed to immunize fish against several bacterial diseases. Direct and random approaches can be used to induce mutations into bacterial pathogens to achieve attenuation. Direct approaches include mutation or deletion of genes involved in metabolic pathways and/or pathogenesis, while random approaches include genetic methods such as transposon mutagenesis or the use of chemicals such as antibiotics (Cane, 2013). Sometime mutations are reversible especially when bacteria mutated by chemicals or by passages. This situation poses a risk for both host and environment. To encounter such problems, bacteria should be mutated by inactivation of genes involved in the metabolic pathways which are necessary for bacterial growth and survival in vivo. Examples of biochemical pathways targeted to produce attenuated strains include the following: aromatic amino acid biosynthesis, purine biosynthesis, capsule biosynthesis, galactose epimerase and adenylate cyclase (Tatum and Briggs, 2005). These mutant bacteria were unable to increase their number to make disease; therefore they cannot survive long enough to create diseases (Roberts *et al.*, 1990). Bacteria have a linear biochemical network for the synthesis of aromatic amino acid. The biosynthesis of aromatic amino acids from core primary metabolism initiates via the shikimate pathway, leading to the synthesis of chorismate. Shikimate pathway is catalyzed by 5-enolpyruvylshikimate 3-phosphate synthase, which leads to the synthesis of enolpyruvylshikimate 3-phosphate (EPSP) encoded by *aroA* gene. The final step in the shikimate pathway is catalyzed by chorismate synthase which converts EPSP to chorismate encoded by *aroC* gene (Oyston *et al.*, 1996; Kitzing *et al.*, 2004; Johansson and Liden,

2006).

Novel live attenuated vaccines have been developed against pathogenic *L. anguillarum* and *Y. ruckeri* by deleting their *aroA* and *aroC* genes. Therefore bacteria are incapable of expressing *aroA* and *aroC* metabolic genes. This mutation is a deletion that the mutation leads to the failure to chorismate synthase that catalyzes the last step in the common shikimate pathway leading to aromatic compounds. The vaccination of rainbow trout with the *aroA* and *aroC* mutant as a live vaccine conferred significant protection (relative percentage survival $\geq 92\%$) against the pathogenic wild-type strain of *L. anguillarum* or *Y. ruckeri*. *aroA* and *aroC* mutant *L. anguillarum* and *aroA* and *aroC* mutant *Y. ruckeri* can be used as a live vaccine to protect fish from vibriosis and yersiniosis, respectively when fish are 3 grams or older (TUBITAK, Project no. 110O886).

Acknowledgements

This project was funded by The Scientific and Technological Research Council of Turkey (TUBITAK, Project no. 110O886). Manuscript was reviewed by Dr. Huseyin Kucuktas at Auburn University, USA.

References

- Ajmal, M. and Hobbs, B.C. 1967. Species of corynebacterium and pasteurilla isolated from diseased salmon, trout and rudd. *Nature*, 215: 142-143
- Akayli, T. 2001. Diagnosis of vibriosis by elisa and methods in cultured Sea bream (*Sparus auratus*, L 1758). Phd. thesis. Istanbul University.
- Akayli, T. and Korun, J. 2004. Bir lepiştes üretim ünitesindeki balıklarda (*Poecilia reticulata*) *Pseudomonas fluorescens* ile birlikte görülen flavobacteriosis olgusu. *Istanbul Üniversitesi Veteriner Fakültesi Dergisi*, 30 (2): 133-142
- Akayli, T. and Timur, G. 2004. Yavru alabalıklarda (*Oncorhynchus mykiss*) pseudomonas septisemi üzerinde bir çalışma. *Istanbul Üniversitesi Veteriner Fakültesi Dergisi*, 30 (1): 121-131
- Akayli, T. and Zeybek, Z. 2005. Bazı akvaryum balıklarında *Plesiomonas shigelloides* enfeksiyonu üzerinde bir araştırma. *Ege Üniversitesi Su Ürünleri Dergisi*, 22 (1-2): 31-34
- Akayli, T. 2007. Çiklet (*Aulonocara maylandi*) balıklarında görülen bir vibriosis olgusu. *Sumder Su Ürünleri Dergisi*, 27:38-40
- Akayli, T., Colak, S. and Yardimci, R.E. 2008. Bacterial infection in *Sparus aurata* cultured in Turkey. In: Middle East and North Africa (MENA) Region Animal Wealth Research Conference, Egypt.
- Akayli, T., Canak, O. and Basaran, B. 2010. A new pseudomonas species observed in cultured young rainbow trout (*Oncorhynchus mykiss* walbaum, 1792): *Pseudomonas plecoglossicida*. *Biyoloji Bilimleri Araştırma Dergisi*, 4(1):107-111
- Akayli, T., Çanak, O. and Başaran, B. 2011. Gökkuşuğu alabalıklarında (*Oncorhynchus mykiss* Walbaum, 1792) görülen *Aeromonas schubertii* enfeksiyonu üzerine bir çalışma. *Biyoloji Bilimleri Araştırma Dergisi*, 4(1): 99-106
- Akayli, T., Urku, C. and Başaran, B. 2011. Kültür balıklarında *Staphylococcus cohnii subsp. cohnii* enfeksiyonu. *Istanbul University Journal of Fisheries and Aquatic Sciences*, 26(1): 1-12
- Aksit, D. and Kum, C. 2008. Gökkuşuğu alabalıkları (*Oncorhynchus mykiss*, Walbaum 1792)'nda sık görülen patojenik mikroorganizmaların tespiti ve antibiyotik duyarlılık düzeylerinin belirlenmesi. *YYU Veterinerlik Fakültesi Dergisi*, 19(1): 1-7
- Aksoy, S. 2009. Bir gökkuşuğu alabalığı (*Oncorhynchus mykiss*) üretim tesisinde (Elazığ) görülen motil *Aeromonas septisemi*. In: Ulusal Su Ürünleri Sempozyumu, Rize.
- Albayrak, H. and Ozan, E. 2010. Investigating the presence of infectious haematopoietic necrosis and infectious pancreatic necrosis virus infections in rainbow trout (*Oncorhynchus mykiss* walbaum, 1792). *Ankara Üniversitesi Veteriner Fakültesi Derg.*, 57: 125-129
- Altınok, I., Kayis, S. and Capkin, E. 2006. *Pseudomonas putida* infection in rainbow trout. *Aquaculture*, 261 (3): 850-855. doi: 10.1016/j.aquaculture.2006.09.009
- Altınok, I., Balta, F., Capkin, E. and Kayis, S. 2007. Disease of rainbow trout caused by *Pseudomonas luteola*. *Aquaculture*, 273 (4): 393-397. doi: 10.1016/j.aquaculture.2007.10.025
- Altun, S., Diler, A., Diler, O., Basak, K. and Isikli, B.T. 2005. Histopathology of streptococcosis in rainbow trout (*Oncorhynchus mykiss* walbaum). *Bulletin of the European Association of Fish Pathologists*, 25(3): 131-135
- Altun, S., Kubilay, A. and Diler, O. 2010. *Yersinia ruckeri* suslarının fenotipik ve serolojik özelliklerinin incelenmesi. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi*, 16(1): 223-229
- Altun, S., Buyukekiz, A.G., Duman, M., Ozyigit, M.O., Karatas, S. and Turgay, E. 2013a. First isolation of *Shewanella putrefaciens* from Gold Fish (*Carrasius auratus auratus*). *Israeli journal of Aquaculture*, (in press, accepted in 3.4.2013)
- Altun, S., Buyukekiz, A.G., Duman, M., Ozyigit, M.O., Karatas, S. and Turgay, E. 2013a. First isolation of *Shewanella putrefaciens* from Gold Fish (*Carrasius auratus auratus*). *Israeli journal of Aquaculture*, (in press, accepted in 3.4.2013)
- Altun, S., Buyukekiz, A., Onuk, E.E., Duman, M. and Ciftci, A. 2013b. Phenotypic, genotypic characterisation and antimicrobial susceptibility determination of *Lactococcus garvieae* strains. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi*, 19 (3): 375-381. doi: 10.9775/kvfd.2012.7754

- Karatas, S. 2013c. First isolation of *citrobacter braakii* from rainbow trout (*Oncorhynchus mykiss*). The Israeli Journal of Aquaculture - Bamidgah, 1: 9
- Altuntas, C. and Oğut, H. 2010. Monthly occurrence and prevalence of viral haemorrhagic septicaemia virus (VHSV) in Whiting *merlangius merlangus*. Diseases of Aquatic Organisms, 88: 107-113. doi: 10.3354/dao02161.
- Anders, K. 1989. Lymphocystis disease of fishes. In: W. Ahne and E. Kurstak (Eds.), Viruses of lower vertebrates, Springer-Verlag, Berlin: 141-160.
- Aronson, J.D. 1926. Spontaneous tuberculosis in salt water fish. Journal of Infectious Diseases, 39: 315-320
- Austin, B. and Stobie, M. 1992. Recovery of *Serratia plymuthica* and presumptive *Pseudomonas pseudoaeruginosa* from skin lesions in rainbow trout, *Oncorhynchus mykiss* (Walbaum), otherwise infected with enteric redmouth. Journal of Fish Diseases of The Aquatic Organisms, 15: 541-543. doi: 10.1111/j.1365-2761.1992.tb00687.x
- Austin, B. and Austin, D.A. 2012. Bacterial fish pathogens; diseases of farmed and wild fish. Springer, New York, London
- Avsever, M.L., Onuk, E.E., Turk, N., Tunaligil, S., Eskiizmirli, S., Incoglu, S. and Yabanli, M. 2012. Pasteurellosis cases in sea bass and sea bream cultured in the Aegean region and other bacteria isolated from these cases. Bornova Veteriner Kontrol ve Araştırma Enstitüsü Dergisi, 34(48): 9-16.
- Ayaz, A. 2006. Dominant aerobic bacterial intestinal flora of sea bass (*Dicentrarchus labrax* L. 1758) larvae in culture conditions. MSc. thesis. İstanbul: İstanbul University.
- Aydin, S., Celebi, S. and Akyurt, I. 1997. Clinical, haematological and pathological investigations of *Escherichia vulneris* in Rainbow trout (*Oncorhynchus mykiss*). Fish Pathology, 32: 29-34.
- Aydin, S., Ciltas, A. and Erman, Z. 1998. *Pseudomonas putida* infections in Scattered mirror carp (*Cyprinus carpio* L.) and Gold fish (*Carassius auratus* L.). In: First International Symposium on Fisheries and Ecology Proceedings, Trabzon.
- Aydin, E. 2000a. Investigation of high mortalities in eyed eggs and fry of rainbow trout (*Oncorhynchus mykiss* Walbaum) and Brown trout (*Salmo trutta*). Turkish Journal of Marine Sciences, 6(1): 245-254.
- Aydin, S., Gultepe, N. and Yildiz, H. 2000b. Natural and experimental infectious of *Campylobacter cryaerophila* in rainbow trout: gross pathology, bacteriology, clinical pathology and chemotherapy. Fish pathology, 35: 117,123
- Aydin, S., Erman, Z. and Bilgin, O.C. 2001. Investigations of *Serratia liquefaciens* infection in rainbow trout (*Oncorhynchus mykiss* Walbaum). Turkish Journal of Veterinary and Animal Sciences, 25 (5): 643-650
- Aydin, S. and Ciltas, A. 2004. Systemic infection of *Aeromonas hydrophila* in rainbow trout: Gross pathology, bacteriology, clinical pathology, histopathology and chemotherapy. Journal of Animal and Veterinary Advances, 12(3): 810-819
- Balta, F. and Cagiran, H. 1998. Kültürü yapılan alabalıklarda görülen *Flexibacter columnaris* enfeksiyonu. In: Doğu Anadolu Bölgesi III. Su Ürünleri Sempozyumu. Erzurum.
- Baran, I., Timur, M., Aydın, N., İstanbulluoğlu, E. and Aydin, M.K. 1980. Çifteler-Sakaryabaşı balık üretim ve araştırma istasyonunda, alabalıklarda (*Salmo gairdneri*) görülen bakteriyel hemorajik septisemi hastalığı üzerine incelemeler. Ankara Üniversitesi Veteriner Fakültesi Dergisi, 27(1): 467-473
- Barker, G.A. and Kehoe, E. 1995. Assessment of disc diffusion methods for susceptibility testing of *Aeromonas salmonicida*. Aquaculture, 134: 1-8
- Baudin, F., Pepin, J.F. and Raymond, J.C. 1991. First observation of an epizootic of pasteurellosis in farmed and wild fish of the French Mediterranean coasts. Paper read at European Association of Fish Pathologists Fifth International Conference on Diseases of Fish and Shellfish. August at Budapest, Hungary.
- Baya, A.M., Toranzo, A.E., Nunez, S., Barja, J. and Hetrick, F.M. 1990. Association of a moraxella sp. and a reo-like virus with mortalities of Atriped bass, *Morone saxatilis*. In: F.O. Perkins and C. Cheng (Eds.) Pathology in marine science. Academic Press, San Diego: 91-99.
- Bernardet, J.F., Baudin Laurencin, F. and Tixerant, G. 1988. First isolation of *Cytophaga psychrophila* in France. Bulletin of the European Association of Fish Pathologists, 8: 104-105
- Bernoth, E.M. 1977. Diagnosis of furunculosis: The tools. In: E.M. Bernoth, A.E. Ellis, P.J. Midtlyng, G. Oliver and P. Smith (Ed.), Furunculosis: Multidisciplinary fish disease research, Academic Press, London: 98-1158
- Boran, H., Terzi, E., Altınok, I., Capkin, E. and Bascinar, N. 2013. Bacterial diseases of cultured mediterranean Horse mackerel (*Trachurus mediterraneus*) in sea cages. Aquaculture, 396: 8-13. doi: 10.1016/j.aquaculture.2013.02.025
- Bragg, R.R., Huchzermeyer, H.F. and Hanisch, M.A. 1990. *Mycobacterium fortuitum* isolated from three species of fish in South Africa. Onderstepoort J. Vet. Res., 57: 101-102
- Bravo, S. and Campos, M. 1989. Coho salmon syndrome in Chile. American Fisheries Society/Fish Health Section Newsletter. 17: 3
- Bullock, G.L. and Snieszko, S.F. 1975. Hagerman redmouth, a disease of salmonids caused by a member of the enterobacteriaceae. United States Department of the Interior, Fish and Wildlife Service, Fish Disease Leaflet. 42: 1-5
- Bulut, M. and Aydin, S. 2004. *Arcobacter cryaerophilus* infection in cross albino Rainbow trout and Rainbow trout: Bacteriology, gross pathology and clinical pathology. Journal of Animal and Veterinary Advances, 6(3): 378-383
- Cagiran, H. 1993a. Kültürü yapılan Cipura (*Sparus aurata* L.) ve Levrek (*Dicentrarchus labrax* L.) balıklarında görülen bakteriyel hastalıkların teşhis ve tedavisi üzerine bir araştırma. Phd. thesis. İzmir: Ege University.
- Cagiran, H. 1993b. The first isolation *Pasteurella piscicida* from cultured Sea bream. Hay. Araştırma Dergisi, 3: 82-83
- Cagiran, H. and Tanrikul, T.T. 1995. Türkiye'deki alabalık (*Oncorhynchus mykiss*, w.) çiftliklerinde yeni bir sorun: Enterococcus benzeri bir bakteri enfeksiyonu. Veteriner Kontrol ve Araştırma Enstitüsü Dergisi, 33 (1): 9-19
- Canak, O. 2011. Determination of the protein profiles of some pathogenic bacteria isolated from cultured Gilt-head sea bream (*Sparus aurata*, L. 1758). Msc. thesis.

- Istanbul: Istanbul University.
- Candan, A. 1991a. Çipura (*Sparus aurata* L. 1758) yetiştiriciliğinde mevsimsel olarak görülen hastalık etkenlerinin tespit ve tedavi yönteminin geliştirilmesi. PhD Thesis. İstanbul: İstanbul Üniversitesi.
- Candan, A. 1991b. Lymphocystis disease of *Sparus aurata* in marine culture at Aegean and Mediterranean coast of Turkey. Mediterranean Regional Aquaculture Project. Basic Level Training Course on Disease, İstanbul.
- Candan, A., Kucuker, M.A. and Karatas, S. 1995. Motile aeromonad septicemia in *Salmo salar* cultured in the Black Sea in Turkey. Bulletin of the European Association of Fish Pathologists, 15(6): 195-196.
- Candan, A., Kucker, A. and Karatas, F. 1996. Pasteurellosis in cultured Sea bass (*Dicentrarchus labrax*) in Turkey. Bulletin of the European Association of Fish Pathologists, 16(5): 150-153.
- Candan, A. 2000. Türkiye'de üretilen Atlantik salmону (*Salmo salar* L.)'nda tespit edilen ilk vibriosis olgusu. Türkiye Mikrobiyoloji Cemiyeti Dergisi, 30: 107-108
- Candan, A. 2002. First report on the diagnosis of infectious pancreatic necrosis (ipn) based on reverse transcription polymerase chain reaction (RT-PCR) in Turkey. Bulletin of the European Association of Fish Pathologists, 22(1): 45-48.
- Cane, K.D. 2013. Attenuated fish vaccine following culture in iron limited media. US patent publication number WO2013028400 A1.
- Casron, J., Gudkovs, N. and Austin, B. 1993. Characteristics of an enterococcus-like bacterium from Australia and South Africa, pathogenic for Rainbow trout, *Oncorhynchus mykiss*. Journal of Fish Diseases, 16: 381-388.
- Castric, J. and Kinkelin, P. 1980. Occurrence of viral haemorrhagic septicemia in Rainbow trout *Salmo gairdneri* Richardson reared in sea water. Journal of Fish Diseases, 3: 21-27.
- Ceschia, G., Quaglio, F., Giorgetti, G., Bertoja, G. and Bovo, G. 1991. Serious outbreak of pasteurellosis (*Pasteurella piscicida*) in euryhaline species along the Italian coasts. In: European Association of Fish Pathologists Fifth International Conference on Diseases of Fish and Shellfish. Budapest, Hungary.
- Chen, S.N., Kou, G.H., Hedrick, R.P. and Fryer, J.L. 1985. The occurrence of viral infections of fish in Taiwan. In: A.E. Ellis (Ed.). Fish and Shellfish Pathology. First International Conference of the European Association of Fish Pathologists, Plymouth, England: Academic Press, London, England.
- Chern, R.S. and Chao, C.B. 1994. Outbreaks of a disease caused by rickettsia-like organism in cultured Tilapias in Taiwan. Fish Pathology, 29: 61-71.
- Chinabut, S. 1999. Mycobacteriosis and nocardiosis. In: P. T. K. Woo and D. W. Bruno (Ed.), Fish diseases and disorders, CAB International Publication, United Kingdom: 319-340.
- Chung, H.J. and Kou, G.H. 1983. *Edwardsiella ictaluri* isolated from cultured eel in Taiwan. Fish Disease Research, 5: 69-70.
- Cruz, J.M., Saraiva, A., Eiras, J.C., Branco, R. and Sousa, J.C. 1986. An outbreak of *Plesiomonas shigelloides* in farmed rainbow trout, *Salmo gairdneri* Richardson, in Portugal. Bulletin of the European Association of Fish Pathologists, 6: 20-22.
- Davis, H.S. 1922. A new bacterial disease in freshwater fishes. US Bureau of Fisheries Bulletin of the Japanese Society of Scientific Fisheries, 38: 261-280
- Davis, H.S. 1926. A new gill disease of trout. Transactions of the American Fisheries Society, 56: 156-159
- Didinen, B. I., Kubilay, A., Diler, O., Ekici, S., Onuk, E.E. and Findik, A. 2011. First isolation of *Vagococcus salmoninarum* from cultured rainbow trout (*Oncorhynchus mykiss*, walbaum) broodstocks in Turkey. Bulletin of the European Association of Fish Pathologists, 31 (6): 235-243
- Diler, O., Demirkan, T., Altun, S. and Çalıkları, F. 1998. Fethiye bölgesindeki bazı alabalık işletmelerinde yersiniosis'in mevsimsel dağılımı üzerine bir araştırma. In: Doğu Anadolu Bölgesi III. Su Ürünleri Sempozyumu: akademia.net.
- Diler, O., Altun, S., Adilgözü, A., Kubik, A. and Isikli, B.T. 2002. First occurrence of streptococcosis affecting farmed rainbow trout (*Oncorhynchus mykiss*) in Turkey. Bulletin of the European Association of Fish Pathologists, 22 (1): 21
- Diler, A., Altun, S. and Isikli, B.I. 2003. Kültürü yapılan gökkuşluğu alabalığı (*Oncorhynchus mykiss*)'ndan izole edilen *Flavobacterium psychrophilum*'un fenotipik karakterleri. SDU Fen Bilimleri Enstitüsü Dergisi, 7 (1): 1-8
- Durmaz, Y. and Türk, N. 2009. Alabalık işletmelerinden motil aeromonasların izolasyonu ve antibiyotiklere duyarlılıklarının saptanması. Kafkas Üniversitesi Veteriner Fakültesi Dergisi, 15 (3): 357-361
- Evas, J.J., Klesius, P.H. and Shiemaker, C.A. 2009. First isolation and characterization of *Lactococcus garviae* from Brazilian Nile tilapia, *Oreochromis niloticus* and pintado, *Pseudoplathystoma corruscans*. Journal of Fish Diseases, 32: 943-951
- Evelyn, T.P.T., Hoskins, G.E. and Bell, G.R. 1973. First record of bacterial kidney disease in an apparently wild salmonid in British Columbia. Journal of the Fisheries Research Board of Canada, 30: 1578-1580
- Gelev, I. and Gelev, E. 1988. A new species of fish-pathogenic bacterium antigenically related to classical brucellae. Zentralblatt für Bakteriologie und Hygiene, Reihe A, 269: 1-6
- Ghittino, C. and Prearo, M. 1992. Report of streptococcosis in Rainbow trout (*Oncorhynchus mykiss*) in Italy. Boll. Soc. Patol. Ittica. , 8: 4-11
- Gjerde, J. 1984. Occurrence and characterization of *Aerococcus viridans* from lobsters, *Homarus gammarus* L., dying in captivity. Journal of Fish Diseases, 7 (5): 355-362
- Gureay, M., Turan, A.T. and Parmaksiz, A. 2013. A study on the presence of infectious pancreatic necrosis virus infections in farmed rainbow trout (*Oncorhynchus mykiss* walbaum, 1792) in Turkey. Kafkas Üniversitesi Veteriner Fakültesi Dergisi, 19(1): 141-146. doi: 10.9775/kvfd.2012.7396
- Gutsell, J. 1946. Sulfa drugs and the treatment of furunculosis in trout. Science, 104: 85-86
- Guvener, R.P. 2001. A study on the diagnosis of the aeromonad infections in some aquarium fishes. Msc. thesis. İstanbul University.
- Halici, G., İstanbulluoğlu, E. and Arda, M. 1977. Bayındır barajı alabalık yetiştirme istasyonunda görülen bakteriyel böbrek hastalığı ve sağitımı. İstanbul Üniversitesi, Veteriner Fakültesi Dergisi, 3 (2): 22-27
- Handler, J., Soltani, M. and Percival, S. 1997. The pathology of *Flexibacter maritimus* in aquaculture species in Tasmania, Australia. Journal of Fish Diseases, 20: 159-168. doi: 10.1046/j.1365-

- 2761.1997.00288.x
- Hastein, T. and Bullock, A.M. 1976. An acute septicaemic disease of Brown trout (*Salmo trutta*) and Atlantic salmon (*Salmo salar*) caused by a pasteurilla-like organism. *Journal of Fish Biology*, 8: 23-26.
- Hawke, J.P., McWhorter, A.C., Steigerwalt, A.G. and Brenner, D.J. 1981. *Edwardsiella ictaluri* sp. Nov., the causative agent of enteric septicaemia of cat fish. *International Journal of Systematic Bacteriology*, 31: 396-400.
- Hedrick, R.P. and Sano, T. 1989. Herpesviruses of fishes. In: W. Ahne and E. Kurstak (Eds.), *Viruses of lower vertebrates*. Springer-Verlag, Berlin: 161-170.
- Hikida, M., Wakabayashi, H., Egusa, S. and Masumura, K. 1979. *Flexibacter* sp, a gliding bacterium pathogenic to some marine fishes in Japan. *Bulletin of Japanese Society of Science in Fisheries*, 45: 421-428.
- Hirvela, V., Pohjanvirta, T., Koski, P. and Sukura, A. 2006. Atypical growth of *Renibacterium salmoninarum* in subclinical infections. *Journal of Fish Diseases*, 29: 21-29. doi: 10.1111/j.1365-2761.2005.00677.x
- Hoshina, T., Sano, T., Morimoto, K. and Jofre, J. 1958. *A streptococcus* pathogenic to fish. *Journal of the Tokyo University of Fisheries*, 44: 57-68.
- Hoshina, T. 1962. On a new bacterium, *Paracolobactrum anguillimortiferum* sp. *Bulletin of the Japanese Society of Scientific Fisheries*, 28: 162-164.
- Hossain, M., Kim, S.R. and Oh, M.J. 2007. Lymphocystis disease virus persists in the epidermal tissues of olive flounder, *Paralichthys olivaceus* (Temminck and Schlegel), at low temperatures. *University Journal of Zoology*, 26: 59-62. doi: 10.1111/j.1365-2761.2009.01048.x
- How, R.A., Bradley, A.J., Iveson, J.B., Kemper, C.M., Kitchener, D.J. and Humphreys, W.F. 1983. The natural history of salmonellae in mammals of the tropical Kimberley region, Western Australia. *Ecological Diseases*, 2: 9-32.
- Ispir, U., Seker, E., Sağlam, N. and Dörücü, M. 2004. Doğu Anadolu bölgesinde bazı gökkuşağı alabalığı (*Oncorhynchus mykiss*) işletmelerinde görülen *Flavobacterium psychrophilum* enfeksiyonunun araştırılması. *F.U. Fen ve Mühendislik Bilimleri Dergisi*, 16(4): 718-724.
- Jensen, N.J., Bloch, B. and Larsen, J.L. 1979. The ulcer-syndrome in cod (*Gadus morhua*). A preliminary virological report. *Nordisk Veterinærmedicin*, 31: 436-442.
- Johansson, L. and Liden, G. 2006. Transcriptome analysis of a shikimic acid producing strain of *Escherichia coli* W3110 grown under carbon and phosphate limited condition. *Journal of Biotechnology*, 126: 528-545.
- Kalaycı, G., Incoglu, S. and Ozkan, B. 2006. First isolation of viral haemorrhagic septicaemia (VHS) virus from turbot (*Scophthalmus maximus*) cultured in the Trabzon coastal area of the Black Sea in Turkey. *Bulletin of the European Association of Fish Pathologists*, 26(4): 157.
- Kalaycı, G., Incioğlu, S., Özyer, B. and Küçükali, Y. 2012. Türkiye'de enfeksiyöz pankreatik nekrozis ve viral hemorajik septisemi hastalıklarının durumu. *Bornova Veteriner Kontrol ve Araştırma Enstitüsü Dergisi*, 34: 31-38.
- Karataş, S. 1996. *Salmo salar* L larda bağırsak florısından *Aeromonas* cinsi bakterilerin izolasyonu MSc Thesis İstanbul: İstanbul University.
- Karataş, S. and Candan, A. 1997. Marmara bölgesindeki bir gökkuşağı alabalığı (*Oncorhynchus mykiss*) işletmesinde yersiniozis vakası. In: IX. Ulusal Su Ürünleri Sempozyumu. Eğirdir, Isparta.
- Karatas, S., Candan, A. and Demircan, D. 2004. Enteric red mouth disease in cultured Rainbow trout (*Oncorhynchus mykiss*) on the Black Sea coast of Turkey. *Israeli Journal of Aquaculture-Bamidgeh*, 56(3): 226-231.
- Karatas, S., Candan, A. and Demircan, D. 2005. Atypical aeromonas infection in cultured sea bass (*Dicentrarchus labrax*) in the Black Sea. *The Israeli Journal of Aquaculture*, 57(4): 255-263.
- Karatas, S., Ercan, D., Steinum, T.M., Turgay, E., Memiş, D. and Candan, A. 2010. First isolation of a *Flavobacterium johnsoniae* like bacteria from cultured Russian sturgeon in Turkey. *Journal of Animal and Veterinary Advances*, 9(14): 1943-1946. doi: 10.3923/javaa.2010.1943.1946.
- Karunasagar, I., Karunasagar, I. and Pai, R. 1992. Systemic *Citrobacter freundii* infection in Common carp, *Cyprinus carpio* L., fingerlings. *Journal of Fish Diseases*, 15: 95-98. doi: 10.1111/j.1365-2761.1992.tb00642.x
- Kayis, S., Capkin, E., Fikri, B. and Altınok, I. 2009. Bacteria in rainbow trout (*Oncorhynchus mykiss*) in the Southern Black Sea Region of Turkey - a survey. *Israeli Journal of Aquaculture-Bamidgeh*. 61 (4): 339-344.
- Keskin, O., Secer, S.U., Izgur, M., Turkyilmaz, S. and Mkakosya, R.S. 2004. *Edwardsiella ictaluri* infection in Rainbow trout (*Oncorhynchus mykiss*). *Turkish Journal of Veterinary and Animal Sciences*, 28 (4): 649-653.
- Kılıç, A., Seker, E., Özcan, M. and İspir, U. 2007. Elâzığ'daki gökkuşağı alabalığı (*Oncorhynchus mykiss*) işletmelerinin bakteriyel yönden incelenmesi. *F. U. Fen ve Mühendislik Bilimleri Dergisi*, 19(2): 129-132.
- Kimura, N., Wakabayashi, H. and Kudo, S. 1978. Studies on bacterial gill disease in salmonids. I. Selection of bacterium transmitting gill disease. *Fish Pathology*, 12: 233-242.
- Kimura, T. and Awakura, T. 1977. Bacterial kidney disease of salmonid; first observation in Japan. *Bulletin of Japanese Society of Science in Fisheries*, 50: 771-774.
- Kirkan, S., Göksoy, E.O. and Kaya, O. 2000. Aydın bölgesinde gökkuşağı alabalıklarından *Yersinia ruckeri* izolasyonu. *Pendik Veteriner Mikrobiyoloji Dergisi*, 31 (1): 23-30.
- Kirkan, S., Goksoy, E.O. and Kaya, O. 2003. Isolation and antimicrobial susceptibility of *Aeromonas salmonicida* in rainbow trout (*Oncorhynchus mykiss*) in Turkey hatchery farms. *Journal of Veterinary Medicine Series B-Infectious Diseases and Veterinary Public Health*, 50(7):339-342.10.1046/j.1439-0450.2003.00671.x
- Kitzing, K., Auweter, S., Amrhein, N. and Macheroux, P. 2004. Mechanism of chorismate synthase, role of the two invariant histidine residues in the Active Site. *Journal of Biological Chemistry*, 279: 9451-9461. doi: 10.1074/jbc.M312471200
- Klein, B.U., Kleingeld, D.W. and Bohm, K.H. 1993. First isolations of *Plesiomonas shigelloides* from samples of cultured fish in Germany. *Bulletin of the European Association of Fish Pathologists*, 13: 70-72.
- Ko, Y.M. and Heo, G.J. 1997. Characteristics of *Flavobacterium branchiophilum* isolated from

- rainbow trout in Korea. Fish Pathology, 32: 97-102.
- Korun, J. and Timur, G. 2001. Gökkuşluğu alabalıklarında (*O. mykiss*) fry mortalite sendromu (FMS) üzerine bir çalışma. Istanbul University Journal of Fisheries and Aquatic Sciences, 12: 15-30.
- Korun, J. 2004a. A study on diagnosis of vibriosis and pasteurellosis some diagnostic kits and laboratory methods in cultured Sea bass (*Dicentrarchus labrax*). Phd. Thesis. Istanbul University.
- Korun, J. and Akaylı, T. 2004b. Kültür levrek balıklarında paraziter bir isopod: *Ceratothoa oestroides* ve sekonder bakteriyel enfeksiyonlar olgusu. İstanbul Üniversitesi, Veteriner Fakültesi Dergisi, 30(2): 123-132.
- Korun, J., Olgac, V., Akgun, K., Colomi, A. and Diamant, A. 2005. Mycobacteriosis in European sea bass, *Dicentrarchus labrax* L., cultured in turkey. Israeli Journal of Aquaculture-Bamidgeh, 57(4): 215-222.
- Korun, J. and Gokoglu, M. 2007a. *Listonella anguillarum* isolated from hatchery-cultured Red porgy *pagrus pagrus* in Turkey. Journal of Animal and Veterinary Advances, 6(6): 823-827.
- Korun, J. and Toprak, B.H. 2007b. Japon (*Carassius auratus*) balıklarından izole ve identifiye edilen *Aeromonas hydrophila*, *A. caviae* ve *A. sobria* türlerinin antibiyotik hassasiyetleri, hemolitik aktiviteleri ve siderofor üretimleri üzerine bir çalışma. Türk Sucul Yasam Dergisi, 3(5): 776-782.
- Korun, J. and Timur, G. 2008. Marine vibrios associated with diseased sea bass (*Dicentrarchus labrax*) in Turkey. Journal of Fisheries Sciences, 2(1): 66-76. doi: 3153/jfscm.2008007
- Korun, J., Akgun, K. and Yazici, M. 2009. Isolation of *Shewanella putrefaciens* from cultured European sea bass, (*Dicentrarchus labrax*) in turkey. Revue Med. Vet., 11(1): 532-536
- Korun, J. and Toprak, H.B. 2010. Kültürü yapılan gökkuşluğu alabalıkları (*Oncorhynchus mykiss*)'nin bağırsağından izole edilen hareketli *Aeromonas* suşlarının antibiyotik hassasiyetleri üzerine NACI'un etkisi. Kafkas Üniversitesi Veteriner Fakültesi Dergisi, 16(2): 193-198
- Kozinska, A. and Pekala, A. 2004. First isolation of *Shewanella putrefaciens* from freshwater fish - a potential new pathogen of fish Bulletin of the European Association of Fish Pathologists, 24 (4):189-193
- Kubilay, A. and Ulukoy, G. 2004. First isolation of *Staphylococcus epidermidis* from cultured Gilthead sea bream (*Sparus aurata*) in Turkey. Bulletin of the European Association of Fish Pathologists, 24(3): 137
- Kubilay, A., Altun, S., Ulukoy, G. and Diler, O. 2005. *Lactococcus garvieae* suşlarının antimikrobiyal duyarlılıklarının belirlenmesi. Eğirdir Su Ürünleri Fakültesi Dergisi, 1(1): 39-48
- Kubilay, A., Altun, S., Diler, O. and Ekici, S. 2008. Isolation of *Flavobacterium columnare* from Cultured Rainbow Trout (*Oncorhynchus mykiss*) Fry in Turkey. Turkish Journal of Fisheries and Aquatic Sciences, 8: 165-169
- Kubilay, A., Altun, S., Didinen, B.I., Ekici, S. and Diler, O. 2009. Gökkuşluğu alabalığı (*Oncorhynchus mykiss*) işletmelerinde *Flavobacterium psychrophilum* izolasyonu. Kafkas Üniversitesi Veteriner Fakültesi Dergisi, 15 (5): 709-715
- Kum, C., Kirkan, S., Sekkin, S., Akar, F. and Boyacioglu, M. 2008. Comparison of in vitro antimicrobial susceptibility in *Flavobacterium psychrophilum* isolated from rainbow trout fry. Journal of Aquatic Animal Health, 20 (4): 245-251. doi: 10.1577/H07-040.1.
- Laird, M. and Bullock, W.L. 1969. Marine fish hematozoa from New Brunswick and New England. Journal of the Fisheries Research Board of Canada, 26: 1075-1102
- Lehmann, J., Mock, D., Stuenkel, F. and Bernardet, J.F. 1991. First isolation of *Cytophaga psychrophila* from a systemic disease in eel and cyprinids. Diseases of Aquatic Organisms, 10: 217-220
- Levis, W. and Bender, M. 1960. Heavy mortality due a bacterium of the genus aeromonas. The Prog. Fish Cult., 22 (1): 11-14
- Love, M., Teebeken, D., Hose, J., Farmer, J., Hickman, F. and Fanning, G. 1981. *Vibrio damsela*, a marine bacterium, causes skin ulcers on the damselfish *Chromis punctipinnis*. Science, 214: 1139-1140
- Matyar, F. 2007. Distribution and antimicrobial multiresistance in gram-negative bacteria isolated from Turkish sea bass (*Dicentrarchus labrax* L., 1781) farm. Annals of Microbiology, 57 (1): 35-38
- McFadden, T.W. 1970. Furunculosis in non-salmonids. Journal of the Fisheries Research Board of Canada, 27: 2365-2370
- McIntosh, D. and Austin, B. 1990. Recovery of an extremely proteolytic form of *Serratia liquefaciens* as a pathogen of Atlantic salmon, *Salmo salar*, in Scotland. Journal of Fish Biology, 36: 765-772
- Mefut, A., Emre, Y., Diler, O., Altun, S. and Ince, I. 2007. Akdeniz bölgesindeki bazı gökkuşluğu alabalığı (*Oncorhynchus mykiss*) işletmelerinden bakteriyel balık patojenlerinin tespiti ve kontrolü. Turk Sucul Yasam Dergisi, 5 (8): 9-18
- Meier, W. and Vestergard, P.E. 1980. Isolation of VHS virus from pike fry (*Esox lucius*) with hemorrhagic symptoms. In: W. Ahne (Ed.), Fish Diseases, Springer-Verlag, Berlin: 8-17
- Muroga, K. and Nakajima, K. 1981. Red spot disease of cultured eel-method for artificial infection. Fish Pathology, 15: 315-318
- Muz, A., Sarıyüpoğlu, M., Ertaş, H.B. and Simsek, A. 1995. Keban baraj gölünden yakalanan bazı balıkların serobik ve mikroaerofilik bakteriler yönünden incelenmesi. Fırat Üniversitesi Sağlık Bilimleri Dergisi, 9 (2): 212-219
- Niggrelli, R.F. and Ruggieri, G.D. 1965. Viral nervous necrosis of larval and juvenile marine fish. In: Proceedings of the international symposium on biotechnology applications in aquaculture. Asian Fisheries Society Special Publication, 10: 147-152
- Nishizawa, T., Savas, H., Isidan, H., Ustundag, C., Iwamoto, H. and Yoshimizu, M. 2006. Genotyping and pathogenicity of viral hemorrhagic septicemia virus from free-living turbot (*Psetta maxima*) in a Turkish coastal area of the Black Sea. Applied and Environmental Microbiology, 72 (4): 2373-2378. doi: 10.1128/aem.72.4.2373-2378.2006
- Ogut, H. and Altuntas, C. 2012. Occurrence and prevalence of infectious pancreatic necrosis virus in rainbow trout (*Oncorhynchus mykiss*) cultured in cages in the Black Sea. Aquaculture Research, 43 (10): 1550-1556. doi: 10.1111/j.1365-2109.2011.02959.x
- OIE. 2013. Piscirickettsiosis, chapter 2.1.13. Diagnostic tests for aquatic animals.
- Olsen, A.B., Melby, H. P., Speilberg, L., Evensen, O. and

- Hastein, T. 1997. *Piscirickettsia salmonis* infection in Atlantic salmon *Salmo salar* in Norway - epidemiological, pathological and microbiological findings. *Diseases of Aquatic Organisms*, 31(1):35-48 doi: 10.3354/dao031035
- Olson, C. and Thomas, J. 1994. An outbreak of infectious hematopoietic necrosis in the baker river system affecting two year classes of sockeye Fish Health Section, American Fisheries Society Newsletter, 22: 1-3.
- Ordal, E.J. and Rucker, R.R. 1944. Pathogenic myxobacteria. *Society of Experimental Biology and Medicine Proceedings*, 56: 15-18.
- Ordal, E.J. and Earp, B.J. 1956. Cultivation and transmission of the etiological agent of bacterial kidney disease. *Proceedings of the Society of Experimental Biology and Medicine*, 85-88.
- Ostland, V.E., Lumsden, J.S., MacPhee, D.D. and Ferguson, H.W. 1994. Characteristics of *Flavobacterium branchiophilum*, the cause of bacterial gill disease in Ontario. *Journal of Aquatic Animal Health*, 6: 13-26. doi: 10.1577/1548-8667(1994)006
- Ostland, V.E., LaTrace, C., Morrison, D. and Ferguson, H.W. 1999. *Flexibacter maritimus* associated with a bacterial stomatitis in Atlantic salmon smolts reared in net-pens in british columbia. *Journal of Aquatic Animal Health*, 11: 35-44. doi: 10.1577/1548-8667(1999)011
- Oyston, P.C., Russel, P. and Williamson, E.D. 1996. An aroA Mutant of *Yersinia pestis* is Attenuated in Guinea-Pigs, but Virulent in Mice. *Microbiology*, 142: 1847-1853. doi: 10.1099/13500872-142-7-1847
- Ozer, S., Aslan, G., Tezcan, S. and Bulduklu, P. 2008a. Genetic heterogeneity and antibiotic susceptibility of *Vibrio alginolyticus* strains isolated from Horse-mackerel (*Trachurus trachurus* L., 1758). *Turkish Journal of Veterinary and Animal Sciences*, 32(2): 107-112.
- Ozer, S., Bulduklu, P.S. and Dönmez, E. 2008b. Mersin ilinde yetiştiriciliği yapılan gökkuşağı alabalıklarında (*Oncorhynchus mykiss*, Walbaum) streptokokozis varlığı. *Journal of Fisheries Sciences*, 2(3): 272-283.
- Ozer, S., Bulduklu, P., Tezcan, S., Dönmez, E., Aydın, E., Aslan, G. and Emekdas, G. 2009. Genetic diversity and antimicrobial susceptibility of motile aeromonads isolated from rainbow trout (*Oncorhynchus mykiss*, walbaum) farms. *Journal of Applied Ichthyology*, 25 (2): doi: 195-200.10.1111/j.1439-0426.2008.01222.x
- Özkök, S. 2005. Gökkuşağı alabalıklarında (*Oncorhynchus mykiss*) görülen önemli bakteriyel etkenlerin tespiti ve antibiyotiklere duyarlılıklarının saptanması. *Etlık Veteriner Mikrobiyoloji Dergisi*, 16 (1-2):1-11.
- Ozturk, D., Adanır, R. and Turutoğlu, H. 2007. Bir sazan (*Cyprinus carpio*) işletmesinde *Aeromonas hydrophila* izolasyonu ve antibiyotik duyarlılığı. *Ulusal Su Ürünleri Sempozyumu*, at Muğla.
- Palacios, M.A., Zamora, M.J., Vasquez, J., Zamora, E. and Duran, A. 1993. Streptococcosis in rainbow trout (*Oncorhynchus mykiss*) in Spain. *Boll. Soc. Patol. Ittica*, 13: 11-16
- Pazos, F., Santos, Y., Macias, A.R., Nunez, S. and Toranzo, A.E. 1996. Evaluation of media for the successful culture of *Flexibacter maritimus*. *Journal of Fish Diseases*, 19: 193-97. doi: 10.1111/j.1365-2761.1996.tb00701.x
- Petersen, A., Andersen, J.S., Kaewmak, T., Somsiri, T. and Dalsgaard, A. 2002. Impact of integrated fish farming on antimicrobial resistance in a pond environment. *Applied and Environmental Microbiology*, 68: 6036-6042. doi: 10.1128/AEM.68.12.6036-6042.2002.
- Plumb, J.A. and Hanson, L.A. 2011. Health maintenance and principal microbial diseases of cultured fish. Third ed. Wiley-Blackwell, Iowa, USA. 244 pp doi: 10.1002/9780470958353.ch11
- Prosyanyaya, V.V. 1981. Some pathogenic bacteria of freshwater fishes. In: J. Olah, K. Molnar and Z. Jeney (Ed.), *Fish pathogens and environment in European polyculture*. Szarvas, Hungary: 82-94.
- Pinto, P.M., Alvarez, P., Bosch, A. and Jofre, J. 1989. Occurrence of a viral erythrocytic infection in the Mediterranean Sea bass, *Dicentrarchus labrax*. *Journal of Fish Diseases*, 12: 185-191. doi: 10.1111/j.1365-2761.1989.tb00291.x
- Rasmussen, C.J. 1965. A biological study of the egg disease (inul). *Annals of the New York Academy of Sciences*, 126: 427-460
- Reyes, X.P. and Campalans, M.B. 1987. Necrosis eritrocítica viral (VEN) presunta en salmon coho de la decima region, Chile. *Investigaciones Marinas, Valparaiso*. 15: 25-31
- Rimaila, E. 2002. First case of bacterial kidney disease (BKD) in Whitefish (*Coregonus lavaretus*) in Finland. *Bulletin of the European Association of Fish Pathologists*, 22 (6): 403-404
- Roberts M., Maskell D., Novotny P. 1990. Construction and characterization in vivo of *Bordetella pertussis* aroA mutants. *Infection and Immunity*, 58:732-739
- Rodger, H.D. and Drinan, E.M. 1993. Observation of a rickettsia-like organism in Atlantic salmon, *Salmo salar* L., in Ireland. *Journal of Fish Diseases*, 16: 361-369. doi: 10.1111/j.1365-2761.1993.tb00869.x
- Ross, A.J., Rucker, R.R. and Ewing, W.H. 1966. Description of a bacterium associated with redmouth disease of rainbow trout (*Salmo gairdneri*). *Canadian Journal of Microbiology*, 12: 763-770
- Saeed, M.O., Alamoudi, M.M. and Al-Harbi, A.H. 1987. A Pseudomonas associated with disease in cultured rabbit fish *Siganus rivulatus* in the Red Sea. *Diseases in Aquatic Organisms*, 3: 177-180.
- Sağlam, Y.S., Işık, N., Arslan, A. and Erer, H. 2006. Erzurum bölgesindeki gökkuşağı alabalıklarında (*Oncorhynchus mykiss* w. 1792) *Aeromonas hydrophila* ve *Yersinia ruckeri* izolasyonu ve patolojik incelemeler. *Atatürk Üniversitesi Veteriner Bilimleri Dergisi*, 1 (2): 6-10
- Sano, T., Fukuda, M., Furukawa, H., Hosoya, H. and Moriya, Y. 1985. A herpesvirus isolated from carp papilloma in japan. In: A. E. Ellis (Ed.), *Fish and shellfish pathology*, Academic Press, Orlando, FL.
- Santos, Y., Huntly, P.J., Turnbull, A. and Hastein, T.S. 1992. Isolation of *Cytophaga psychrophila* in association with rainbow trout mortalities in the United Kingdom. *Bulletin of the European Association of Fish Pathologists*, 6: 12
- Santos, Y., Pazos, F. and Barja, J.L. 1999. *Flexibacter maritimus*, causal agent of flexibacteriosis in marine fish. G. Oliver (Ed.), *ICES Identification Leaflets for Diseases and Parasites of Fish and Shellfish*, International Council for the Exploration of the Sea, Copenhagen, Denmark: 1-6.
- Sanz, F. 1991. Rainbow trout mortalities associated with a mixed infection with *Citrobacter freundii* and IPN virus. *Bulletin of the European Association of Fish Pathologists*, 11: 222

- Sato, N., Yamane, N. and Kawamura, T. 1982. Systemic *Citrobacter freundii* infection among Sunfish *Mola mola* in matsushima aquarium. Bulletin of the Japanese Society of Fisheries, 48: 1551-1557
- Savaş, H., Yıldırım, Y., Kurtoglu, I.Z., Bascinar, N., Alkan, A., Gürel, M., Ergun, H., Firidin, S., Kutlu, I., Serdar, S. ve Zengin, B. 2006. Ordu ili Perşembe ilçesinde faaliyet gösteren yüzer kafes işletmelerinin çevresel etki ve su ürünleri sağlığı yönünden izlenmesi projesi, Sonuç raporu, SUMAE, 58pp
- Savas, H., Altınok, I., Cakmak, E. and Firidin, S. 2006. Isolation of *Renibacterium salmoninarum* from cultured Black Sea Salmon (*Salmo trutta labrax*): First report in Turkey. Bulletin of the European Association of Fish Pathologists, 26 (6): 238-246
- Savaş, H. and Türe, M. 2007. Bölgemizde doğal ve kültürlü yapılan balıklarda görülen hastalıklar. SUMAE Yunus Araştırma Bülteni, 7: 10-13.
- Schaperclaus, W. 1938. Die schädigungen der deutschen fischerei durch fischparasiten und fischkrankheiten. Allgemeine Fischerei-Zeitung, 41: 256-270
- Schiewe, M.H. and Crosa, J.H. 1981. Molecular characterization of *Vibrio anguillarum* biotype 2. Canadian Journal of Microbiology, 27: 1011-1018
- Sekar, V.T., Santiago, T.C., Vijayan, K.K., Alavandi, S.V., Raj, V.S., Rajan, J.J.S., Sanjuktha, M. and Kalaimani, N. 2008. Involvement of *Enterobacter cloacae* in the mortality of the fish, *Mugil cephalus*. Letters in Applied Microbiology, 46: 667-672
- Şeker, E., Özcan, M., Kilic, A. and Ispir, U. 2005. Malatya'daki bazı balıkçılık işletmelerinden temin edilen gökkusağı alabalığı (*Oncorhynchus mykiss*)'nin bakteriyolojik yonden araştırılması. In XIII. Ulusal Su Ürünleri Sempozyumu. Canakkale.
- Seker, E., Karahan, M., Ispir, U., Cetinkaya, B., Saglam, N. and Sarieyyupoglu, M. 2012. Investigation of *Yersinia ruckeri* infection in rainbow trout (*Oncorhynchus mykiss* walbaum 1792) farms by polymerase chain reaction (PCR) and bacteriological culture. Kafkas Üniversitesi Veteriner Fakültesi Dergisi, 18 (6): 913-916
- Smith, I.W. 1963. The classification of Bacterium salmonicida. Journal of General Microbiology, 33: 263-274.
- Snieszko, S.F. and Bullock, G.L. 1965. Freshwater fish diseases caused by bacteria belonging to the genera aeromonas and pseudomonas. U.S.A. Fish and wildlife service. Fishery Icafilet No.459.
- Soltani, M., Nikbakht, G., Mousavi, H.A.E. and Ahmadzadeh, N. 2008. Epizootic outbreaks of lactococcosis caused by *Lactococcus garviae* in rainbow trout in Iran. Bulletin of the European Association of Fish Pathologists, 28 (5): 207-212
- Starliper, C.E. 2001. Isolation of Serratia liquefaciens as a pathogen of Arctic charr, *Salvelinus alpinus* (L.). Journal of Fish Diseases, 24: 53-56. doi: 10.1046/j.1365-2761.2001.00266.x
- Takahashi, K., Okamoto, N., Kumagai, A., Maita, M., Ikeda, Y. and Rohovec, J.S. 1982. Epizootics of erythrocytic inclusion body syndrome in coho salmon cultured in seawater in Japan. Journal of Aquatic Animal Health, 4: 174-181. doi: 10.1577/1548-8667(1992)004<0174:EOEIBS>2.3.CO;2
- Talaat, A.M., Trucksis, M., Kane, A.S. and Reimschuessel, R. 1999. Pathogenicity of *Mycobacterium fortuitum* and *Mycobacterium smegmatis* to Gold fish, *Carassius auratus*. Veterinary Microbiology, 66: 151-164. doi: 10.1016/S0378-1135(99)00002-4
- Tanrikul, T.T. and Çağırhan, H. 2001. Doğadaki kefal balıklarında görülen pasteurellosis salgını. E.U. Su Ürünleri Dergisi, 18 (3): 509-512
- Tanrikul, T.T., Çağırhan, H. and Toksen, E. 2004. Levreklerden (*Dicentrarchus labrax* L., 1758) izole edilen vibrio türlerinin API 20E yöntemiyle identifikasyonu. Ege Üniversitesi Su Ürünleri Dergisi, 2004 (3-4): 243-247
- Tanrikul, T.T. 2007. Vibriosis as an epizootic disease of rainbow trout (*Oncorhynchus mykiss*) in Turkey. Pakistan Journal of Biological Sciences, 10: 1737-1737
- Tatum, F.M. and Briggs, R.E. 2005. Construction of In-Frame *aroA* Deletion Mutants of *Mannheimia haemolytica*, *Pasteurella multocida*, and *Haemophilus somnus* by Using A New Temperature-Sensitive Plasmid. Applied and Environmental Microbiology, 71: 7196-209. doi: 10.1128/AEM.06362-11
- Tel, O.Y., Irgare, G., Karahan, M. and Keskin, O. 2007. Şanlıurfa Balıklıgöl balıklarından *Aeromonas hydrophila* izolasyonu ve antibiyotik duyarlılıklarının saptanması. Etlik Veteriner Mikrobiyoloji Dergisi, 18 (1): 1-4
- Timur, M. 1983. An outbreak of disease of farmed eel (*Anguilla anguilla*) due to *Aeromonas hydrophila* in Turkey. Histopathological and bacteriological studies. Ankara Üniversitesi Veteriner Fakültesi Dergisi, 30 (3): 361-367
- Timur, G. and Timur, M. 1985. Eğirdir Gölü sudak (*Stizostedion lucioperca* L. 1758) balıklarında yüksek mortaliteye neden olan bakteriyel hemorajik septisemi hastalığı üzerine bir araştırma. Ankara Üniversitesi, Veteriner Fakültesi Dergisi, 32 (1): 33-41
- Timur, G. 1991a. A historical study of a Carp pox (viral epithelioma) disease in Turkey. Bulletin of the European Association of Fish Pathologists, 11 (5): 171
- Timur, G. and Timur, M. 1991b. An outbreak of enteric red mouth disease in farmed rainbow trout (*O.mykiss*) in Turkey. Bulletin of the European Association of Fish Pathologists, 11 (5): 182
- Timur, G., Karataş, S., Çolak, S. and Akaylı, T. 1999. Gökkuşluğu alabalık yavrularında görülen furunkulozis hastalığı üzerine bir çalışma. İstanbul Üniversitesi, Su Ürünleri Dergisi, 461-467,
- Timur, G., Korun, J. and Güvener, R.P. 2003. Lepistes (*Poecilia reticulata*) üretim ünitesindeki anaç ve yavru balıklarda ağır mortalite ile seyreden aeromonad septisemisi üzerine bir çalışma. İstanbul University Journal of Fisheries and Aquatic Science, 15: 1-11
- Timur, G., Timur, M. and Korun, J. 2004. A study on the outbreak of *Flavobacterium psychrophilum* infection in a rainbow trout (*Oncorhynchus mykiss*) hatchery in Turkey. İstanbul University Journal of Fisheries and Aquatic Sciences, 17: 21-27
- Timur, G., Timur, M., Akaylı, T., Korun, J. and Thompson, K.D. 2005. First observation of rickettsia-like organisms in cultured sea bass (*Dicentrarchus labrax*) in Turkey. Bulletin of the European Association of Fish Pathologists, 25 (5): 196-202
- Timur, G., Timur, M., Akaylı, T., Korun, J. and Erkan, M. 2008. The first observation of viral erythrocytic infection associated with pathogenic bacteria and some ectoparasites in cultured Mediterranean Sea bass (*Dicentrarchus labrax* L.) in Turkey. Bulletin of the

- European Association of Fish Pathologists, 28 (2): 58-65
- Timur, G., Akayli, T., Korun, J. and Yardimci, R.E. 2010. A study on bacterial haemorrhagic septicemia in farmed young Russian sturgeon in Turkey (*Acipenser gueldenstaedtii*). Istanbul Universitesi, Su Urunleri Dergisi, 25 (1): 19-27
- Timur, G., Timur, M., Akayli, T., Korun, J. and Erkan, M. 2011. First occurrence of erythrocytic inclusion body syndrome (EIBS) in marine reared rainbow trout (*Oncorhynchus mykiss*) in Turkey. Israeli Journal of Aquaculture-Bamidgeh, 63:1-8
- Timur, G., Yardimci, R.E., Urku, C. and Canak, O. 2011. Marmara Bölgesi kültür gökkuşuğu alabalıklarında lactococcosisin bakteriyolojik ve histopatolojik metotlarla teşhisi. İstanbul Üniversitesi, Su Ürünleri Dergisi, 26 (1): 63-81
- Tokşen, E. 1999. Ege bölgesinde yetiştiriciliği yapılan çipura ve levrek balıklarının solungaçlarında görülen metazoan parazitler ve tedavisi. PhD Thesis, Ege University
- Toranzo, A.E., Baya, A.M., Romalde, J.J. and Hetrick, F.M. 1989. Association of *Aeromonas sobria* with mortalities of adult gizzard shad, *Dorosoma cepedianum lesueur*. Journal of Fish Diseases, 12: 439-448
- Toranzo, A.E. and Barja, J.L. 1990. A review of the taxonomy and seroepizootiology of *Vibrio anguillarum*, with special reference to aquaculture in the Northwest of Spain. Diseases of Aquatic Organisms, 9: 73-82
- Toranzo, A.E., Barreiro, S., Casal, J.F., Figueras, A., Magarinos, B. and Barja, J.L. 1991. Pasteurellosis in cultured Gilthead bream (*Sparus aurata*): First report in Spain. Aquaculture, 99: 1-15
- Toranzo, A.E. and Barja, J.L. 1993. Fry mortality syndrome (FMS) in Spain. Isolation of the causative bacterium *Flexibacter psychrophilus*. Bulletin of the European Association of Fish Pathologists, 13: 30-32
- Toranzo, A.E. 2004. A review of the main bacterial fish diseases in mariculture systems. Aquaculture, 246: 37-61. doi: 10.1016/j.aquaculture.2005.01.002
- Toranzo, E., Magarinos, B. and Romalde, L. 2005. A review of the main bacterial fish diseases in mariculture systems. Aquaculture, 246 (4): 37-61. doi: 10.1016/j.aquaculture.2005.01.002
- TUIK, 2012. Su ürünleri istatistikleri. http://www.tuik.gov.tr/Kitap.do?metod=KitapDetay&KT_ID=13&KITAP_ID=52 (accessed June 15, 2013).
- Türk, N. 2002. Çipura (*Sparus aurata*) ve levrek (*Dicentrarchus labrax*) yavrularından bakteri izolasyonu ve antibiyotiklere duyarlılıklarının belirlenmesi. Bornova Veteriner Kontrol ve Araştırma Enstitüsü Dergisi, 27: 7-14
- Uzbilek, M. and Yildiz, H. 2002. A report on spontaneous diseases in the culture of Grass carp (*Ctenopharyngodon idella* val.1844), Turkey. Turkish Journal of Veterinary and Animal Sciences, 26: 407-410
- Vigneulle, M. and Baudin, F. 1995. *Serratia liquefaciens*: A case report in turbot (*Scophthalmus maximus*) cultured in floating cages in France. Aquaculture, 132: 121-124. doi: 10.1016/0044-8486(94)00375-X
- Wakabayashi, H. and Egusa, S. 1972. Characteristics of a *Pseudomonas* sp. From an epizootic of pond-cultured eels (*Anguilla japonica*). Bulletin of Japanese Society of Science in Fisheries, 38: 577-587
- Wakabayashi, H., Egusa, S. and Fryer, J.L. 1980. Characteristics of filamentous bacteria isolated from a gill disease of salmonids. Canadian Journal of Fisheries and Aquatic Sciences, 37: 1499-1504
- Whalen, J.G., Mully, T.W. and English, J.C. 2007. Spontaneous *Citrobacter freundii* infection in an immunocompetent patient. Archives of Dermatology, 143 (1): 124-125. doi: 10.1001/archderm.143.1.124
- Wichardt, U.P., Johansson, N. and Ljungberg, O. 1989. Occurrence and distribution of *Aeromonas salmonicida* infections on Swedish fish farms, 1951-1987. Journal of Aquatic Animal Health, 1: 187-196. doi: 10.1577/1548-8667(1989)001<0187:OADOAS>2.3.CO;2
- Winton, J.R., Batts, W.N., Nishizawa, T. and Stehr, C.M. 1989. Characterization of the first North America isolates of viral haemorrhagic septicaemia virus. Fish Health Newsletter, 17 (2)
- Woo, P.T.K. and Bruno, D.W. 2011. Fish disease and disorders; viral, bacterial and fungal infections volume 3. CABI Publishing, New York, USA
- Wood, E.M., Snieszko, S.F. and Yasutake, W.T. 1955. Infectious pancreatic necrosis in Brook trout. American Medical Association Archives of Pathology, 60: 26-28
- Yıldırım, S. and Özer, S. 2010. Mersin ili Çağlarca köyündeki gökkuşuğu alabalığı (*Oncorhynchus mykiss*, Walbaum, 1792) kuluçkahanelerinde *Flavobacterium* spp. varlığı. Journal of FisheriesSciences.com, 4(1):112-122.