



Acute Sensitivity Comparison among *Daphnia magna* Straus, 1820 *Daphnia pulex* Leydig, 1860 and *Simocephalus vetulus* Müller, 1776, Exposed to Nine Toxicants

Gustavo Emilio Santos-Medrano^{1,*}, Roberto Rico-Martínez^{1,*} 

¹ Universidad Autónoma de Aguascalientes, Centro de Ciencias Básicas, Departamento de Química. Avenida Universidad 940, Aguascalientes, Ags. México, C.P. 20100.

Article History

Received 05 December 2017
Accepted 11 July 2018
First Online 30 July 2018

Corresponding Author

Tel.: +55.449 9108420
E-mail: rrico@correo.uaa.mx

Keywords

Aquatic Toxicology
Cladocera
Environmental Legislation
Environmental Toxicology
Metal Toxicity

Abstract

A comparison of acute toxicity (LC50 values) among nine toxicants: a) six metals: Cd, Cr, Cu, Hg, Pb, Ti, and b) three organics: benzene, ethyl acetate, and toluene, between the exotic cladoceran species *Daphnia magna* Straus 1820 and two native strains of freshwater cladoceran species: *Daphnia pulex* Leydig, 1860 and *Simocephalus vetulus* (O. F. Müller, 1776) was performed. We hypothesized that the exotic species would be less sensitive than native species. Our hypothesis was fulfilled. *Daphnia magna* was less sensitive than native species to eight of the nine compounds we analyzed. Results suggest that native species are better adjusted to local environmental conditions, and are more reliable as bioindicators of potential effects of toxicants on aquatic biota. Although the use of *D. magna* is recommended because of its ample toxicity database, some researchers propose the use of native species for toxicity tests. Therefore, to propose a native species to be considered as a test organism for official toxicity test for tropical countries, there is a need to increase the database of toxicants and to compare the sensitivities of several classes of toxicants with a model organism whose sensitivity to a broad variety of toxicants is well known, like *D. magna*.

Introduction

Cladocerans (also known as water fleas) are a group of microcrustaceans commonly distributed in freshwater bodies and occurring in a variety of habitats; they play an important role on food webs, both as consumers of phytoplankton and fish prey (Dettmers & Stein, 1992). Within this group, daphnids have been broadly used in aquatic toxicology, they have demonstrated to be sensitive to a large variety of chemicals (Adema, Kuiper, Hanstveit & Canton, 1982; Baudouin & Scoppa, 1974). *Daphnia magna* has been recommended for toxicity evaluation of many chemicals in water (ASTM, 2014; ISO International Standard 6341, 1996; OECD 202, 2004; U.S. EPA, 2002), and has been extensively used in toxicity testing.

However, it is not a native species in many countries, is a Holarctic organism with geographic distribution of cold and temperate environments of North America, Europe and Asia (Weber, 1993). Several countries: Austria, Australia, Belgium, Canada, Denmark, France, Germany, Italy, New Zealand, Netherlands, Norway, Portugal, Spain, Sweden, United States and United Kingdom contemplate *D. magna* in their legislature as model organism to quantify toxicity (Wolska *et al.* 2010). Toxicological information for tropical or subtropical species is very scarce, hence, the countries of these regions have to use international water quality criteria developed for *D. magna* (ASTM, 2014; ISO International Standard 6341, 1996; OECD 202, 2004; U.S. EPA, 2002) which leads to a certain grade of uncertainty. This can be seen in the available data

(acute and chronic) found in the ECOTOX (U.S.EPA, 2017) and ECETOC (2003) databases. For *D. magna*, there are 403 toxicity values in ECOTOX (U.S.EPA, 2017) and 217 in ECETOC (2003), while for *Daphnia pulex* and *Simocephalus vetulus*, there are 87 (U.S.EPA, 2017) and 25 (ECETOC, 2003) and 16 ECOTOX (U.S.EPA, 2017) and 4 (ECETOC, 2003) references, respectively. In Mexico, *D. magna* is an exotic species, and it is unable to survive naturally in Mexican reservoirs. In contrast, there are records of *D. pulex* and *S. vetulus* in the states of Aguascalientes, Campeche, Mexico, Hidalgo, Guanajuato and Yucatan (Elías-Gutiérrez, Gutiérrez, Silva, Granados & Garfias, 2008). At the beginning of the year 2000, the International Development Research Centre (IDCR) started an international research network called WaterTox, with the participation of institutions from Argentina, Canada, Chile, Colombia, Costa Rica, India and Mexico (Forget *et al.* 2000a). The goal of WaterTox was to develop and validate toxicity tests with different organisms, including *D. magna* in acute toxicity tests. During the development of this research, the countries of Argentina, Colombia, Costa Rica and India, reported having problems for the maintenance of cultures of *D. magna* and the consequent loss of the species in addition to considering the realization of these tests with a high cost (Forget *et al.* 2000b). Mohamed (2007) compared the sensitivity to seven chemicals between *Ceriodaphnia rigaudii* (a native species of Trinidad) and the exotic *Daphnia magna* and found that for six toxicants *C. rigaudii* was more sensitive than *D. magna* and concluded that due to this differences in sensitivity perhaps *D. magna* might not be an adequate sentinel species for Trinidad.

The aim of this work was to determine the acute toxicity of nine toxicants (six metals and three organic compounds) and compared the reference values from *D. magna* with those from *D. pulex* and *S. vetulus* to test the hypothesis that native species are more sensitive to toxicants than *D. magna*.

Materials and Methods

The *D. magna* strain is from the EPA and was donated by Dr. Fernando Martinez Jerónimo of the National School of Biological Sciences of IPN, Mexico. The species of *D. pulex* and *S. vetulus* were collected in a pond of the botanic garden of the Autonomous University of Aguascalientes, with geographic coordinates N 21.912, W -102.313. Cladocerans were cultured with EPA medium (96 mg NaHCO₃, 60 mg CaSO₄•2H₂O, 60 mg MgSO₄ and 4 mg KCl per liter of deionized water, U.S.EPA, 2002), which is a synthetic moderately hard water (80-100 CaCO₃ L⁻¹), pH 7.5, temperature 20°C ± 2, and photoperiod of 16: 8, (light: darkness) in a bioclimatic chamber (Revco Scientific), fed with the algae *Pseudokirchneriella subcapitata* (Korshikov) F. Hindák 1990 (UTEX 1648) grown in Bold's Basal Medium (Nichols, 1973). Metals Cd, Cr, Cu, Pb

and Ti were standards of atomic absorption diluted in 3% nitric acid (Sigma). Mercuric chloride was diluted in EPA medium. Benzene, ethyl acetate and toluene were analytical chemistry standards of Sigma Aldrich and was diluted in acetone. Toxicity tests were performed according to EPA methods (U.S.EPA, 2002). Acute test consisted in exposing ten neonates less 24h for each treatment, one negative control and five nominal concentrations. We performed a solvent control for the organic compounds with the highest acetone concentration used in tests to monitor the toxicity due to solvent.

Ten neonates were selected randomly and placed in one of the five replicates 250 ml glass beakers containing 100 ml of test solution per toxicant concentration. Organisms were incubated for 48h at 20°C, photoperiod 16:8 (light: darkness) in absence of food. At the end of incubation dead organisms were counted.

Lethal media concentration (LC50) values were calculated using regression with probit units and the logarithm of each toxicant concentration by the software Statistica 10.0 (Stat Soft Inc., 2001). The NOEC (No observed effect concentration) and LOEC (Lowest observed effect concentration) values were calculated by one-way analysis of variance (ANOVA) and Duncan test. We performed a Spearman correlation by transforming the LC50 values of each species into logarithm, to determine if there is an association between the values of toxicity of *D. magna* compared to *D. pulex* and *S. vetulus*.

Relative sensitivity were calculated by the followed formula (Von der Ohe & Liess, 2004):

$$RS = \log (LC50 D. magna / LC50i)$$

Where: RS= relative sensitivity; LC50 *D. magna* = LC50 value for *D. magna* and LC50 = value for species *i*. A value of zero indicates a sensitivity equal to that of *D. magna*, positive value *D. magna* is less sensitive, negative value *D. magna* is most sensitive.

Results

Table 1 shows the toxicity values achieved for three species of cladocerans used in this work. In terms of LC50 values, *D. magna* was more sensitive to Pb; the native species: a) *D. pulex* was the most sensitive to benzene, Cr, Cu, and toluene; b) *S. vetulus* was the most sensitive to Cd, ethyl acetate, Hg and Ti (Table 1). The order of toxicity of the metals in descending order was, Cu = Hg > Cd > Pb > Cr > Ti to *D. magna*, for *D. pulex*: Hg = Cu > Cd > Pb > Cr > Ti and Hg = Cd = Cu > Pb > Ti > Cr to *S. vetulus*. For the organic compounds the order of toxicity was Ethyl acetate > Toluene > Benzene for the three species. In the case of *D. magna* and *D. pulex* the most toxic metals were Cu and Hg and the least toxic Ti, however for *S. vetulus* the Hg, Cd and Cu presented the highest toxicity and Cr least toxic.

The EPA (U.S.EPA, 2002) establishes a range of coefficient of variation of 21 to 58%, results obtained in our tests are below those values (Table 1).

Metal toxicity for cladoceran group according to the geometric mean of the LC50 (Table 2) was established as follows Hg> Cu> Cd> Pb> Cr> Ti and organic compounds was ethyl acetate > toluene> benzene. In accordance to LC50 values, the difference between the most toxic and less toxic metal was set in this way, *D. magna* 859 fold (Cu-Ti), 2,808 (Hg-Ti) for *D. pulex* and 3,280 (Hg -Cr) for *S. vetulus*. In the case of organic compounds, the differences were 134, 65 and 720 fold (ethyl acetate-benzene) for *D. magna*, *D. pulex* and *S. vetulus* respectively. We compared the mean values of LC50 (Figure 1a) to determine if they show significant differences. There are no differences in LC50 values for Cu and Hg among the three cladoceran species (Figure 1a). The LC50 values for Cr were different among the three species. Metals that did not

show differences between *D. magna* and *D. pulex* were: Cd, Pb, and Ti, and between *D. magna* and *S. vetulus* Pb and Ti were different. Ethyl acetate revealed no significant difference between the three species, only toluene was statistically similar to *D. magna* and *S. vetulus* (Figure 1b). Figure 2a shows that *S. vetulus* was more sensitive than *D. magna* for Cd, Hg, Ti, for organic compounds sensitivity was higher to ethyl acetate and toluene (Figure 2b), *D. pulex* was more sensitive to Cd, Cr, Cu, Hg and the three organic compounds (Figure 2a and 2b) compared to *D. magna*. Titanium was the only toxic without difference in the relative sensitivity among *D. magna* and *D. pulex*.

Table 3 presents different LC50 values of the toxics used in this work, mainly found in ECOTOX (U.S.EPA, 2017) and ECETOC (2003) databases, where it is stated that *D. magna* is the species with the highest number of toxicity tests (287) followed of *D. pulex* (137) and finally *S. vetulus* (5). A Spearman correlation

Table 1. Endpoints of the acute toxicity test performed with three species of cladoceran. The most sensitive LC₅₀ values are highlighted with bold numbers. N = 5

Species	Parameter	Benzene	Cadmium	Chromium	Copper	Ethyl acetate	Lead	Mercuric chloride	Titanium	Toluene
<i>D. magna</i>	LC10	62.01	0.10	2.53	0.006	1.04	0.75	0.017	9.75	33.13
	LC50	207.53	0.20	3.04	0.013	1.54	1.08	0.028	11.17	44.16
	NOEC	50	<0.125	2.0	<0.005	1.0	0.25	<0.015	<10.0	20.0
	LOEC	100	0.125	2.5	0.005	1.5	1.0	0.015	10.0	30.0
	CL	145.64	0.1998	2.9020	0.0120	0.4585	0.9474	0.0258	11.0077	39.6551
		295.80	0.2174	3.2033	0.0140	1.6166	1.2445	0.0315	11.3448	49.1926
	CV%	17.72	3.09	1.96	8.52	4.75	14.26	6.71	0.48	8.27
	R ²	0.81	0.98	0.76	0.94	0.93	0.83	0.76	0.93	0.74
<i>D. pulex</i>	LC10	31.11	0.11	1.16	0.003	1.07	0.56	0.0019	10.71	9.74
	LC50	86.49	0.17	1.43	0.007	1.32	1.09	0.004	11.22	15.45
	NOEC	25	<0.10	0.5	0.001	<1.0	0.1	0.0005	10.0	5
	LOEC	50	0.10	1.0	0.005	1.0	0.5	0.001	11.0	10
	CL	72.2680	0.1670	1.2755	0.0059	1.2855	0.9516	0.0030	11.0509	13.0797
		96.8500	0.1857	1.6062	0.0087	1.3930	1.2676	0.0053	11.3972	18.2767
	CV%	9.06	0.73	3.70	17.14	6.15	3.61	13.17	2.10	21.93
	R ²	0.93	0.95	0.80	0.81	0.91	0.93	0.76	0.78	0.75
<i>S. vetulus</i>	LC10	288.60	0.004	2.21	0.013	0.42	0.75	0.0003	0.51	22.54
	LC50	475.66	0.015	3.28	0.017	0.66	1.34	0.001	2.34	42.89
	NOEC	<200.0	0.001	<2.0	0.010	0.25	0.5	0.0001	<0.2	<20.0
	LOEC	200.0	0.005	2.0	0.015	0.50	1.0	0.0005	0.2	20.0
	CL	422.2795	0.0113	3.0746	0.0160	0.5834	1.1738	0.0008	2.2222	38.7257
		535.9200	0.0208	3.5067	0.0182	0.7603	1.5441	0.0014	3.5620	47.5006
	CV	4.65	5.22	2.39	7.42	4.65	7.12	6.55	5.25	8.37
	R ²	0.83	0.85	0.85	0.81	0.83	0.87	0.85	0.76	0.87

LC₅₀=Lethal concentration where 50% of animals. LC₁₀= Lethal concentration where 10% of animals die. NOEC = No Observed Effect Concentration. LOEC = Lowest Observed Effect Concentration. All values in mg/L. R² = Coefficient of determination of the LC₅₀. CV = Coefficient of variation of LC₅₀. CL = 95% Confidence limits for the LC₅₀ values.

Table 2. Geometric mean of the CL₅₀ toxicity values of three cladoceran species. All values in mg/L

Endpoint	Benzene	Cadmium	Chromium	Copper	Ethyl acetate	Lead	Mercuric chloride	Titanium	Toluene
Geometric mean LC ₅₀	204.384	0.079	2.424	0.011	1.102	1.164	0.004	6.643	30.815

of LC₅₀ values between *D. magna*-*D. pulex* and *D. magna*-*S. vetulus* was performed. The correlation coefficient between *D. magna*-*D. pulex* was 0.88 for metals (figure 3a) and 1.0 for the organic compounds (figure 4a). The correlation coefficient for *D. magna*-*S. vetulus* was 0.59 and 1.0 for metals and organic compounds respectively (Figure 4a and 4b).

Discussion

Our original hypothesis is fulfilled by the experimental data. *Daphnia magna* was less sensitive than the native species (either *D. pulex* or *S. vetulus*) to eight of the nine compounds we analyzed (See Table 1 and Figure 2). Only in the case of lead *D. magna* LC₅₀ absolute value (1.08 mg/L) was lower than that of *D. pulex* (1.09 mg/L) (Table 1), but when we compared the relative sensitivity between both species this value is zero (Figure 2a), which means there is no difference in sensitivity between these two species exposed to lead. These results suggest that native species collected in Mexican reservoirs are better adjusted to the

environmental conditions of Mexico and more reliable as bioindicators of the potential effects of toxicants on Mexican aquatic biota. Although the use of *Daphnia magna* is recommended because of its ample toxicity database, our results showed that it is less sensitive to toxicant exposure than two native species. Some researchers propose the use of native species in toxicity tests: a) Martínez-Jerónimo, Rodríguez-Estrada & Martínez-Jerónimo (2008) proposed *Daphnia exilis* Herrick, 1895 for Mexico, b) Melnikov & de Freitas (2011) proposed *Daphnia similis* Claus, 1876 for Brazil, c) Aparecida, da Silva, Cristina, & Rocha (2014) proposed *Ceriodaphnia silvestrii* Daday 1902 and *Macrothrix flabelligera* Smirnov 1992, for Brazil, and d) Teklu & Van den Brink (2016) proposed *Diaphanosoma brachyurum* (Liévin, 1848) for Ethiopia. With the results obtained in this work we confirmed the difference between the relative sensitivity of *D. magna*/*D. pulex* and *D. magna*/*S. vetulus*, as well as providing toxicity data for *S. vetulus* for different metals and organic compounds that were not found in the databases. Based on this we can suggest the use of




Figure 1. Comparison of LC₅₀ values of *D. magna* against *D. pulex* and *S. vetulus* by ANOVA and Duncan test. Vertical bars represented standard error. The same letter means no significant difference. P<0.05. a) metals, b) organic compounds. Abbreviations: Dm=*D. magna*, Dp=*D. pulex*, Sv=*S. vetulus*.




Figure 2. Relative sensitivity expressed as a logarithmic proportion (LC_{50}) relative to the sensitivity of *D. magna*. a) metals and b) organic compounds.

Table 3. Acute toxicity values for *D. magna*, *D. pulex* and *S. vetulus*, 48h, LC_{50} . All values in mg/L

Toxicant	Species	Range	Mean	n	Reference
Benzene	<i>D. magna</i>	56.65-1130	275.72	18	U.S. EPA (2017) , ECETOC (2003)
	<i>D. pulex</i>	15-345	208.33	3	
Cd	<i>D. magna</i>	0.0009-22.6	0.21	171	U.S. EPA (2017) , ECETOC (2003)
	<i>D. pulex</i>	0.0004-1.04	0.15	82	U.S. EPA (2017) , ECETOC (2003)
	<i>S. vetulus</i>	0.024-0.089	0.056	2	U.S. EPA (2017)
Cr	<i>D. magna</i>	0.02-2.4	0.60	21	U.S. EPA (2017) , ECETOC (2003)
	<i>D. pulex</i>	0.57-2.0	0.57	18	U.S. EPA (2017) , ECETOC (2003)
	<i>S. vetulus</i>	-	0.50	1	U.S. EPA (2017)
Cu	<i>D. magna</i>	0.008-0.12	0.03	57	U.S. EPA (2017) , ECETOC (2003)
	<i>D. pulex</i>	0.006-0.37	0.07	23	U.S. EPA (2017) , ECETOC (2003)
	<i>S. vetulus</i>	-	0.057	1	U.S. EPA (2017)
Pb	<i>D. magna</i>	0.13-4.4	1.0	5	U.S. EPA (2017)
	<i>D. pulex</i>	0.37-9.9	3.54	6	U.S. EPA (2017) , ECETOC (2003)
	<i>S. vetulus</i>	-	4.5	1	U.S. EPA (2017)
Hg	<i>D. magna</i>	0.001-0.02	0.005	13	U.S. EPA (2017)
	<i>D. pulex</i>	0.002-0.024	0.008	5	U.S. EPA (2017)

native species for toxicity tests due to their biological and ecological importance, and that are representative of each region.

The analysis of sensitivity of the three cladoceran species we tested exposed to nine toxicants (six metals and three organics), resulted in *D. pulex* being the most sensitive to two metals (Cr and Cu), and two organics

(benzene and toluene), and with similar sensitivity to Pb than *D. magna*. *S. vetulus* was the most sensitive to three metals (Cd, Hg and Ti), and one organic (ethyl acetate).

Comparing the LC_{50} results of our work with the ECOTOX (U.S.EPA, 2017) and ECETOC (2003) databases (Table 3), *D. magna* remains within the range of




Figure 3. Correlation between acute toxicity values of *D. magna*, *D. pulex* (a) and *S. vetulus* (b) for metals.




Figure 4. Correlation between acute toxicity values of *D. magna*, *D. pulex* (a) and *S. vetulus* (b) for organic compounds.

benzene, Cd, Cu, Pb and Hg, for *D. pulex* benzene, Cd, Cr, Cu, Pb and Hg. Moreover, *D. magna* and *S. vetulus* were less sensitive to Cr; otherwise, *S. vetulus* was more sensitive for Cd, Cu and Pb.

In general, *D. magna* is less sensitive to Cu when comparing these ten values to our results (Table 4). On the other hand, the values between *S. vetulus* and *S. expinosus* are very similar to those obtained in our strain. For *D. pulex*, the values were intermediate between those reported by these authors (Table 4). Our *D. magna* Cd value in three of the four values was less sensitive. The *D. pulex* strain of Elnabarawy *et al.* 1986, show greater resistance than our *D. pulex* strain. Badudouin & Scoppa (1974) determined a lethal median concentration value for Cr of 0.023 mg/L to *D. hyalina*, which differs 138, 65 and 149-fold compared to *D. magna*, *D. pulex* and *S. vetulus* respectively (Table 4). The most sensitive organisms to Pb was *D. hyalina* Leydig 1860 in contrast with *D. pulex* strain of Elnabarawy *et al.* (1986), it was more resistant when compared to the values obtained in this research. The LC50 values of Hg to *D. pulex* and *S. vetulus* of this work are very similar to those found by Badudouin & Scoppa (1974) for *D. hyalina* of 0.005 mg/L and Elnabarawy *et al.* (1986) for *D. pulex* (0.003 mg/L) and *Ceriodaphnia reticulata* (Jurine, 1820) (0.002 mg/L) in contrast to our *D. magna* strain resulted less sensible.

We found differences in the relative sensitivity of *D. magna* compared to *D. pulex* and *S. vetulus* in accordance with Wogram & Liess (2001) and Von der Ohe & Liess (2004), where they assessed the cladocera group against *D. magna*, ensuing the group being more sensitive to metals and organic compounds. The highest differences in relative sensitivity were found

between *D. magna* and *S. vetulus* for Cd, Hg and Ti. *D. magna* was more sensitive solely 27% of total test performed.

Results of the spearman correlation evidence that all three species of cladocerans were responding similar to organic compounds (Figure 4). Moreover, we found differences with action mode of metals (Figure 3) mainly with correlation *D. magna-S. vetulus*.

The LC50 values were evaluated by the Spearman rank correlation coefficient. This comparison provided values of 1.0 between *D. magna-D. pulex* (Fig. 4a) and *D. magna-S. vetulus* (Fig. 4b) for organic compounds. We assume the acute toxic responses of three cladoceran are strongly correlated and the mode of action of chemicals is the same. However, the correlation between *D. magna-D. pulex* (Fig. 3a) was strongest (0.80) than *D. magna-S. vetulus* (0.59) Fig. 3b, we presume, metals have a different mode of intoxicated according to the organism.

Information obtained from the Web of Science (2018) registered 238 publications of *D. magna* on metal acute toxicity during 1984 to 2017. The contribution by country was: United States 62, China 59, Canada 22, Mexico 15, France 14, Portugal 14, Brazil 12, India 12 Netherlands 12, United Kingdom 10, Australia 9, Spain 9, England 6, Germany 6, Japan 6, Turkey 5, Belgium 4, Chile 4, Finland 4, Taiwan 4, Italy 3, Scotland 3, Slovakia 3, South Korea 3, Argentina 2, Egypt 2, Malaysia 2, Nigeria 2, Poland 2 and Vietnam 2. With respect to *D. pulex* the total of publications are 31, compiled in the years 2006 to 2017 and they are distributed in the following way: United States 12, France 5, United Kingdom 4, Mexico 3, Belgium, Canada, England, Germany and China with two works

Table 4. Acute toxicity values for *D. magna*, *D. pulex* and *S. vetulus*, 48h, LC₅₀. All values in mg/L

Toxicant	Species	LC ₅₀ values	References
Cd	<i>D. hyalina</i>	0.055	Baudouin & Scoppa, 1974
	<i>C. reticulata</i>	0.18	Elnabarawy et al.1986
	<i>D. pulex</i>	0.31	
	<i>D. pulex</i>	0.13	Santos-Medrano & Rico-Martínez, 2015
Cr	<i>D. hyalina</i>	0.023	Baudouin & Scoppa, (1974)
	<i>D. hyalina</i>	0.005	Baudouin & Scoppa, (1974)
	<i>D. pulex</i>	0.03	Elnabarawy et al.1986
	<i>Ceriodaphnia reticulata</i>	0.02	
	<i>D. galeata</i> G. O. Sars, 1864	0.02	
Cu	<i>D. pulex</i>	0.003	Bossuyt & Janssen, 2005
	<i>Ceriodaphnia pulchella</i> G. O. Sars, 1862	0.016	
	<i>D. galeata</i>	0.009/0.018	Baudouin & Scoppa, 1974
	<i>D. longispina</i> (O. F. Mueller, 1785)	0.022	
	<i>Simocephalus expinosus</i> (O. F. Müller, 1776)	0.018	
	<i>S. vetulus</i>	0.021	Elnabarawy et al. 1986
<i>D. hyalina</i>	0.60		
Pb	<i>C. reticulata</i>	1.80	Elnabarawy et al. 1986
	<i>D. pulex</i>	2.02	
	<i>D. pulex</i>	1.62	Santos-Medrano & Rico-Martínez, 2015
	<i>D. hyalina</i>	0.005	Baudouin & Scoppa, 1974
Hg	<i>C. reticulata</i>	0.002	Elnabarawy et al. 1986
	<i>D. pulex</i>	0.003	

for each one. The Web of Science database does not contain any publications related to *S. vetulus*.

In this work the acute toxicity tests were developed under the same conditions of temperature, pH, hardness of the test solution, photoperiod and chemical purity, these parameters being indispensable for the quality control of the results obtained from acute toxicity test. According to the USEPA (2002), an important factor for the accuracy of the results is the coefficient of variation (CV). In this research, the CV ranges were 0.48-17.14% and 4.65-21.93 for metals and organic compounds respectively, which indicates the reliability of the results. We report the first acute toxicity values of ethyl acetate and toluene for the three species of cladoceran tested and benzene for *S. vetulus*. This is the first report in which relative sensitivity and the correlation between species for acute toxicity tests with six metals and three organic compounds are included. Some authors (Wogram and Liess, 2001; Sanchez-Bayo, 2006) have reported comparisons of the sensitivity of *D. magna* among other species of cladocerans. However, they did not perform the toxicity tests, they used values of the USEPA database; in these cases, feed data (species and concentration) of the different species of cladocerans was not taken into account, which can reduce the accuracy when comparing the sensitivity between species. On the other hand, Cowgill *et al.* (1984) performed toxicity tests to compare the sensitivity between *D. magna*-*Ceriodaphnia dubia/affinis*, although the feeding of the cultures was with different species of algae, *S. capricornotum* and *A. convolutus* respectively, and this may alter the sensitivity of the organism.

We have a large database for *D. magna*, which includes 2475 toxicants with 48-h exposure tests, and 3184 total toxicants. In contrast, the database for *D. pulex* includes 316 for 48-h tests and 524 total toxicants. In the case of *S. vetulus* includes only 34 for 48-h tests and 49 total toxicants (U.S.EPA, 2017). ECETOC (2003) reports 113 total toxicants for *D. magna*, but only 16 for *D. pulex* and 15 for *S. vetulus*. In the case of ECOTOX (U.S.EPA, 2017) for 48-h tests for the toxicants used in this work, the database includes 157 toxicants for *D. magna*, 97 for *D. pulex*, but only 13 for *S. vetulus*. Therefore, to propose a native species as a model organism for official toxicity test for the Mexican legislation (or for any tropical country), there is an urgent need to increase the database of toxicants. It is necessary to compare the sensitivities of several classes of toxicants with a native model organism whose ecology, genetics, sensitivity to a broad variety of toxicants is well known as is the case of *D. magna*.

Conclusions

The exotic species *D. magna* was the least sensitive species to nine toxicants (six metals and three

organics), when compared with the native species *D. pulex* and *S. vetulus*. Our results suggest that native species collected in Mexican reservoirs are better adjusted to the environmental conditions of Mexico and more reliable as bioindicators of the effects of toxicants on Mexican aquatic biota. There is a need to increase the knowledge on native species (like *D. pulex* and *S. vetulus* used in this contribution), to match information on *Daphnia magna* (especially the ample toxicity database), to propose a native species as a sentinel organism for Mexican reservoirs. This contribution includes the first: a) acute toxicity values of ethyl acetate and toluene for the three species of cladocerans, b) acute toxicity value of benzene for *S. vetulus*, and c) relative sensitivity and correlation between three cladoceran species for acute toxicity tests with six metals and three organic compounds.

References

- Adema, D. M. M., Kuiper, J., Hanstveit, A.O., & Canton J.H. (1982) Consecutive system of tests for assessment of the effects of chemical agents in the aquatic environment. In IUPAC Pesticide Chemistry: *Human Welfare and the Environment*, (Pergamon, Press, Elmsford, New York), 522 pp.
- Aparecida, M.R., da Silva, M. A., Cristina L.S., & Rocha, O. (2014). A comparative study of the acute toxicity of the herbicide atrazine to cladocerans *Daphnia magna*, *Ceriodaphnia silvestrii* and *Macrothrix flabelligera*. *Acta Limnologica Brasiliensia*. 26(1), 1-18. DOI: 10.1590/S2179-975X2014000100002
- ASTM E729-96. (2014). Standard Guide for Conducting Acute Toxicity Tests on Test Materials with Fishes, Macroinvertebrates, and Amphibians, ASTM International, West Conshohocken, PA.
- Baudouin, M.F. & Scoppa, P. (1974). Acute Toxicity of various metals to fresh water zooplankton. *Bulletin of Environmental Contamination and Toxicology*. 12(6), 745-51. <https://doi.org/10.1007/BF01685925>
- Bossuyt, B.T.A., & Janssen, C.R. (2005). Copper toxicity to different field-collected cladoceran species: intra- and inter-species sensitivity. *Environmental Pollution*. 136(1), 145-154. <https://doi.org/10.1016/j.envpol.2004.11.023>
- Cowgill, U. M., Takahashi, I. T. and Applegath, S.L. 1984. A comparison of the effect of four benchmark chemicals on *Daphnia magna* and *Ceriodaphnia dubia-affinis* tested at two different temperatures. *Environmental Toxicology and Chemistry*. 4 (3), 415-422.
- Dettmers, J.M., & Stein R.A. (1992) Food consumption by larval gizzard shad: zooplankton effects and implications for reservoir communities. *Transactions of the American Fisheries Society*. 121(4) 494-507. <http://dx.doi.org/10.1577/1548-8659>.
- Elías-Gutiérrez, M., Suárez, E., Gutiérrez, M., Silva, M., Granados, J. & Garfia, T. (2008). Cladocera y Copépoda de las aguas continentales de México. UNAM, México D.F., 323 pp
- Elabarawy, M.T., Welter, A.N., & Robideau, R.R. (1986). Relative sensitivity of three daphnid species to selected organic and inorganic chemicals. *Environmental*

- Toxicology and Chemistry*. 5(4), 393-398. <http://dx.doi.org/10.1002/etc.5620050409>.
- ECETOC. European Centre for Ecotoxicology and Toxicology of Chemicals. (2003). TR 091: ECETOC Aquatic Toxicity (EAT) database - EAT Database.
- Forget, G., Gagnon, P., Sanchez, W. A. & Dutka, B. J. (2000a). Overview of Methods and Results of the Eight Country International Development Research Centre (IDRC) WaterTox Project. *Environmental Toxicology*. 15(4), 264-276. [http://dx.doi.org/10.1002/1522-7278\(2000\)15:4<264::AID-TOX2>3.0.CO;2-Q](http://dx.doi.org/10.1002/1522-7278(2000)15:4<264::AID-TOX2>3.0.CO;2-Q).
- Forget, G., Sanchez-Bain, A., Diaz-Baez, M. C., Arkhipchuk, V., Pica-Granados, Y., Ronco, A., Beauregard, T., Srivastava, R. C., Blaise, C., Castillo, G., Dutka, B.J. & Castillo, L. E. (2000b). Preliminary Data of a Single-Blind, Multicountry Trial of Six Bioassays for Water Toxicity. Monitoring. *Environmental Toxicology*. 15(5), 362-369. [http://dx.doi.org/10.1002/1522-7278\(2000\)15:5<362::AID-TOX2>3.0.CO;2-Q](http://dx.doi.org/10.1002/1522-7278(2000)15:5<362::AID-TOX2>3.0.CO;2-Q).
- ISO 6341. (1996). Water quality—Determination of the Inhibition of the Mobility of *Daphnia magna* Straus (Cladocera, Crustacea).
- Martínez-Jerónimo F, Rodríguez-Estrada J & Martínez-Jerónimo L. (2008). *Daphnia exilis* Herrick, 1895 (Crustacea:Cladocera). Una especie zooplanctónica potencialmente utilizable como organismo de prueba en bioensayos de toxicidad aguda en ambientes tropicales y subtropicales. *Revista Internacional de Contaminación Ambiental*. 24(4), 153-159.
- Melnikov, P. & de Freitas, T.C.M. (2011). Evaluation of acute Chromium (III) toxicity in relation to *Daphnia similis*. *Journal of Water Resource and Protection*. 3(2), 127-130. doi:10.4236/jwarp.2011.32015.
- Mohamed A. (2007). Comparative sensitivities of the tropical cladoceran, *Ceriodaphnia rigaudii* and the temperate species *Daphnia magna* to seven toxicants. *Toxicological and Environmental Chemistry*. 89(2), 347-352.
- Nichols, H.W. (1973). Growth media-freshwater. In: Stein JR (ed) Handbook of phycological methods. Culture methods and measurements. Cambridge University Press, London, pp 7-24.
- Organization for Economic Cooperation and Development (OECD). (2004). Guideline for Testing of Chemicals. *Daphnia* sp., Acute Immobilisation Test. OECD 202. Paris, France.
- Sánchez-Bayo F. (2006) Comparative acute toxicity of organic pollutants and reference values for crustaceans. I. Branchiopoda, Copepoda and Ostracoda. *Environmental Pollution*. 139 (3), 385–420.
- Santos-Medrano, G.E., & Rico-Martínez, R. (2015). Acute and chronic effects of five metals in a battery of freshwater planktonic organisms. *Fresenius Environmental Bulletin*. 24(12b), 4658-4666.
- Teklu, B.M., Negussie, R. & Van den Brink, P.J. (2016). Sensitivity of Ethiopian aquatic macroinvertebrates to the pesticides endosulfan and diazinon, compared to literature data. *Ecotoxicology*. 25(6), 1226-1233. <https://doi.org/10.1007/s10646-016-1676-0>.
- U.S. Environmental Protection Agency. (2002). Methods for measuring the acute toxicity of effluents and receiving waters to freshwater and marine organisms, 5th ed, EPA-821-R-02-012. US Environmental Protection Agency, Washington.
- U.S. Environmental Protection Agency. (2017). ECOTOX User Guide: ECOTOXicology Knowledgebase System. Version 4.0. Retrieved from <http://www.epa.gov/ecotox/>.
- Von der Ohe, P.C. & Liess, M. (2004). Relative sensitivity distribution of aquatic invertebrates to organic and metal compounds. *Environmental Toxicology and Chemistry*. 23(1), 150-156. <http://dx.doi.org/10.1897/02-577>.
- Web of Science. (2018). Clarivate Analytics. Retrieved from <http://www.webofknowledge.com> (last accessed on February 16th, 2018).
- Weber, C.I. (1993). Methods for measuring the acute toxicity of effluents and receiving waters to freshwater and marine organisms, 4th ed. United States Environmental Protection Agency, Cincinnati, EPA/600/4-90/027F.
- Wogram, J., & Liess, M. (2001). Rank ordering of macroinvertebrates species sensitivity to toxic compounds by comparison with that of *Daphnia magna*. *Bulletin of Environmental Contamination and Toxicology*. 67(3), 360-367. <https://doi.org/10.1007/s001280133>.
- Wolska, L., Kochanowska, A. & Namiesnik, J. (2010). Application of Biotests. In Jacek Namiesnik and Piotr Szefer (Eds.), *Analytical measurements in aquatic environments* (pp. 189-221). Boca Raton, FL. CRC Press, 503 pp.