

# Functional Ingredients from Seafood Processing Wastes: Protein Hydrolysate and Biocalcium

Umesh Patil<sup>1,\*</sup> , Krisana Nilsuwan<sup>2,\*</sup> , Soottawat Benjakul<sup>1,2,\*</sup> 

<sup>1</sup>International Center of Excellence in Seafood Science and Innovation, Faculty of Agro-Industry, Prince of Songkla University, Hat Yai, Songkhla, 90110, Thailand

<sup>2</sup>Kyung Hee University, Department of Food and Nutrition, Seoul 02447, Republic of Korea

## How to Cite

Patil, U., Nilsuwan, K., Benjakul, S. (2024). Functional Ingredients from Seafood Processing Wastes: Protein Hydrolysate and Biocalcium. *Turkish Journal of Fisheries and Aquatic Sciences*, 24(4), TRJFAS25347. <https://doi.org/10.4194/TRJFAS25347>

## Article History

Received 29 September 2023

Accepted 03 January 2024

First Online 12 January 2024

## Corresponding Author

E-mail: krisana.n@psu.ac.th  
soottawat.b@psu.ac.th

## Keywords

Fish protein hydrolysate  
Fish biocalcium  
Production  
Bioactivity and bioavailability  
Applications

## Abstract

The fish processing industry generally generates large amounts of byproducts, including skin, bone, scale, or meat residues, etc. These byproducts contain a wide range of nutritional components such as protein, lipid, mineral, etc. These leftovers can be converted to functional ingredients or nutraceuticals. To produce high value-added functional ingredients, the remaining proteins in fish processing leftovers are hydrolyzed, in which hydrolysate containing different peptides and amino acids can be generated. Frame or backbone as well as scale can be converted to fish biocalcium, which serves as a potential source of calcium and minerals. Nevertheless, potential processing or technology must be adopted to obtain desired products with target bioactivities. This review covers production, characteristics, and bioavailability of fish protein hydrolysate (FPH) and fish biocalcium (BC). This review also focuses on the recent applications of the FPH and BC in drink and foods. Overall, the review points out the better exploitation of fish processing leftovers and simultaneously lower pollution caused by the improper discard or disposal of those wastes to environment. Also, consumers have more choices to consume functional ingredients or nutraceutical from marine resources.

## Introduction

In Asia, the total fisheries and aquaculture production reached a record of 214 million tons in 2020, comprising 178 million tons of aquatic animals and 36 million tons of algae as highlighted in the State of World Fisheries and Aquaculture 2022 report. The increasing global fish production has given rise to a large quantity of raw material utilized for processing. Consequently, significant quantities of fish processing leftover, comprising inedible tissues such as bones/frame, skin/scales, gills, swim bladders, fins, intestines, blood, roes, liver, etc. are generated (Kumar et al., 2018). These

remnants are often disposed into the river, waterways, and oceans, resulting in severe environmental contamination. Over the past few decades, efforts have been made to exploit these leftovers via transforming them into fish meal, fertilizer, or animal feed with a low market value (Patil et al., 2022b). Nonetheless, it is crucial to recognize that these leftovers are rich sources of valuable components such as protein, bioactive peptides, enzymes, minerals, lipid, vitamins, flavorant, pigments, etc. (Kumar et al., 2018). Therefore, it is imperative to convert these wastes into value-added products in order to increase their market value and minimize environmental pollution.

Fish frame or trimming still has some meat retained. These leftovers can be utilized as raw material for protein hydrolysate production. The common method for production of fish protein hydrolysate (FPH) is enzymatic process, which is easily controlled to end up with the designed degree of hydrolysis (DH) (Kristinsson & Rasco, 2000). Also, such a mild condition does not lead to side effects such as the development of toxic compounds, etc. FPH has several functional properties such as emulsification, foaming, oil binding, and water retention, and various bioactive properties including antioxidant, antimicrobial, antihypertensive, etc. (Misir, 2022). To produce high quality FPH, undesirable attributes, especially fishy odor, must be minimized. Also, bitterness is another drawback of FPH, in which debittering could be adopted to increase consumer acceptability (Idowu & Benjakul, 2019). Simultaneously, the bioactivity or nutraceutical properties must be maximized, mainly contributing to the active peptides in FPH. Sequence of amino acid in peptides are also known to determine bioactivity. Due to the hygroscopic in nature of FPH, the handling and storage must be done properly to avoid absorption of moisture from environment (Chiodza & Goosen, 2023). FPH with bioactivities has been widely employed in various processed foods, in order to enhance their intrinsic value and give them "functional food" feature. Moreover, FPH also has been used in pharmaceutical and nutraceutical industries (Rana et al. 2023).

Calcium (Ca) is known to be an important element required for many physiological functions in the human

body, including cofactor in enzymatic reactions, muscle contraction, blood clotting, nerve function, and strengthening teeth and bones (Anderson & Garner, 1996). Dairy products including cheese, milk, calcium fortified products, and dietary supplements are generally the essential sources of calcium (Tunick, 1987). However, milk or dairy products are rarely consumed in some populations, particularly Asian people, owing to lactose intolerance and indigestion (Gaskin & Ilich, 2009). Consequently, calcium-fortified foods or calcium salts can be alternative calcium supplements. Fish calcium showed a comparable bioavailability to calcium from skim milk in both rats (Larsen et al., 2000) and humans (Hansen et al., 1998). Frame or backbone and scale of fish, which are byproducts from fish processing industry, can serve as a potential source of minerals and calcium. Minerals including calcium, phosphorus and hydroxyapatite have been found in fish bones at approximately 60-70% (Kim & Mendis, 2006b). The calcium/phosphorus (Ca/P) mole ratio of hydroxyapatite (HA:  $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$ ) and tricalcium phosphate (TCP:  $\text{Ca}_3(\text{PO}_4)_2$ ) were 1.67 and 1.50, respectively (Piccirillo et al., 2013). HA crystals are localized between gap zones of collagen fibrils. Collagen fibrils align together with HA in bones to provide strengthening effect (Benjakul et al., 2019) (Figure 1). In addition, clinical trial and biological study revealed that fish bone calcium is an excellent source of bioavailable calcium, suggesting that fish bone can serve as the promising source of dietary calcium (Hansen et al., 1998; Idowu et al., 2019a).




Figure 1. The structure of a hydroxyapatite-collagen composite in fish frame/backbone tissues.

**Protein Hydrolysates from Fish Processing Leftover**

Seafood processing by-products can contain up to 60% of proteins (on dry weight basis) (Sasidharan & Venugopal, 2020). This has urged the production of fish protein hydrolysates (FPH) through hydrolysis process to produce bioactive peptides and essential nutrients from these by-products, thereby increasing their value. The schematic presentation of general approach to produce FPH and their applications is shown in Figure 2. The desire for the sustainable utilization of fish processing by-products has spurred the advancement of protein hydrolysis processes and recovery, in which their functionalities can be improved, and application can be achieved (Nirmal et al., 2022).

**Production of Fish Protein Hydrolysate (FPH)**

Hydrolyzing proteins into smaller peptides has been demonstrated to enhance their nutritional and functional properties (Kudo et al., 2009). Protein hydrolysis has been used to transform by-products into more profitable forms (Patil et al., 2023). Head, bone/frame, skin, viscera, and other by-products have potential to be utilized in the production of FPH. These

FPH, composed of relatively short peptides (2–20 amino acids), can be generated by breaking down proteins into peptides either chemical or enzymatic hydrolysis (He et al., 2013). Chemical hydrolysis involves the use of acid or base to facilitate protein hydrolysis. This method is straightforward and advantageous for its cost-effectiveness. However, chemical hydrolysis presents challenges in controlling the molecular weight (MW) of the resulting peptides, and it may yield a significant quantity of salt along with adverse side reactions including decomposition of tryptophan and racemization (Gao et al., 2021).

Alternatively, the enzymatic hydrolysis process has emerged as the preferred method due to fast, safe, and easily controlled methods with mild reaction conditions (Patil et al., 2019). This process has been proven highly effective in producing novel peptides from byproducts, with superior bioactivities and functionality (Kristinsson & Rasco, 2000). Enzymatic hydrolysis employs proteolytic enzymes, both endopeptidases, which cleave the peptide bonds within proteins and exopeptidases, which hydrolyze peptide bonds from either N or C terminal (Clemente, 2000; Raksakulthai & Haard, 2003). Commercially available proteases used for FPH production come from different sources: (a) microbial sources, e.g. flavourzyme, alcalase, neutrase,




Figure 2. The schematic presentation of FPH production and its potential applications.

umamizyme, protamex; (b) plant sources, e.g. bromelain, ficin, papain; and (c) animal sources, e.g. chymotrypsin, trypsin, pepsin (Gao et al., 2021). FPH were produced with the help of commercial microbial enzyme namely Alcalase from *Bacillus licheniformis*, plant enzyme namely papain from the latex of *Carica papaya* (Idowu et al., 2019b) and animal gastrointestinal enzymes including pepsin and trypsin from porcine and bovine sources (Ketnawa et al., 2017). In addition to commercial enzymes, crude proteases from fish viscera were also used to obtain FPH (Nalinanon et al., 2011). Partially purified proteases such as pepsin and trypsin from fish viscera are also utilized for production of FPH (Patil et al., 2022a).

Nevertheless, each protease exhibits a distinct specificity for peptide bonds adjacent to specific amino acid residues, allowing for selective protein hydrolysis to achieve desired properties (Wu et al., 2003). Factors like raw material composition, enzyme type, and hydrolyzing condition including, time, temperature, pH, enzyme-to-substrate ratio, solid-liquid ratio, amount of enzyme, and production conditions (e.g. ultrasonic treatment) directly affect the physicochemical, functional properties and bioactivities of FPH (Misir & Koral, 2019; Patil et al., 2022c). Controlled hydrolysis has been known as a potential means to improve the functional and bioactive properties of proteins (Karami & Akbari-Adergani, 2019).

### Properties of FPH

Protein hydrolysis results in the generation of hydrolysates or peptides possessing altered or improved bioactivities and functional properties when compared to their native proteins (Kitts & Weiler, 2003). FPH possess desirable functional properties (e.g., solubility, emulsifying, foaming, oil, water/oil binding capacities, etc.) (Kristinsson & Rasco, 2000) and several beneficial bio-activities (anti-oxidative, anti-hypertensive, anti-microbial, anti-tumor activities, etc.) with potential applications in foods, nutraceuticals and pharmaceutical products (He et al., 2013; Nirmal et al., 2022).

### Functional Properties

The hydrolysis process plays a vital role in influencing the molecular size, hydrophilicity, and hydrophobicity of FPH (Kristinsson & Rasco, 2000), consequently impacting their functional properties (Karami & Akbari-Adergani, 2019). Solubility is considered as a crucial functional property of FPH. Other functional properties like foaming and emulsifying, are directly linked to solubility (Benjakul et al., 2014). Protein hydrolysis contributes significantly to the enhancement of solubility. This can be ascribed to the breakdown of proteins into smaller soluble peptides, resulting in a higher exposure of charged carboxyl and amino groups (Yin et al., 2008). The high solubility of FPH implies their potential applicability in formulated food

products (Foh et al., 2010; Pires & Batista, 2013). Strong correlation between solubility and degree of hydrolysis (DH) was established (Dinakarkumar et al., 2022). However, an extensive hydrolysis process may negatively impact other functional properties (Benjakul et al., 2014). For example, salmon heads FPH produced by hydrolysis with Alcalase with higher DH exhibited superior solubility. Nonetheless, better properties such as fat absorption capacity, emulsifying capacity, and emulsion stability were obtained when DH was lower (Kristinsson & Rasco, 2000).

Emulsifying properties of FPH are strongly affected by their solubility (Benjakul et al., 2014). Peptides with low molecular weight (LMW) owing for high solubility, which immediately disperse and attach to the interface. Nevertheless, these smaller peptides are less capable of reducing the interfacial tension, compared to larger peptides, primarily because of unfolding and reorganization at the interface (Patil & Benjakul, 2019). Consequently, this causes speedy flocculation and subsequently affects the stability protein layer. The molecular weight (MW) distributions of FPH play an important role in emulsion stabilization. Peptides generated through limited hydrolysis exhibited superior emulsifying and foaming properties compared to those subjected to extensive hydrolysis (Mazorra-Manzano et al., 2012).

FPH with LMW peptide fractions enhanced water holding capacity, since they were highly hydrophilic in nature (Halim et al., 2016). Fat binding capacity can be influenced by the enzyme-substrate specificity, DH and bulk density of proteins (Villamil et al., 2017). Two FPH from *Rastrineobola argentea* had the decreased fat binding ability when the peptide chain length upsurged (Mbatia et al., 2014).

Moreover, functional properties of protein hydrolysate were further modified by formation of different complexes, such as protein-polysaccharide (Xie et al., 2023), protein-polyphenols (Patil et al., 2023), etc. The influences of chitosan and rice protein hydrolysate (CH/RH) ratio and oil-water ratio on the emulsification efficiency, as well as the impact of RH on the formation of CH/RH complex coacervates was investigated by Xie et al. 2023. CH/RH at a 1:1 ratio is more effective in enhancing the interfacial adsorption and emulsification stability than CH or RH alone, especially at 50% oil volume fraction.

### Bioactivities

The bioactivities of FPH or peptides are linked to their structural properties, amino acid composition, and sequences. FPH derived from the enzymatic hydrolysis of fish processing by-products have been demonstrated to be a rich source of bioactive peptides, known for their beneficial effects on consumer health (Bougatef et al., 2010; Ren et al., 2008). Nevertheless, the choice of protease in the enzymatic hydrolysis process plays a crucial role. It can lead to significantly different

characteristics in the resulting peptides, particularly in terms of their bioactivity. The use of multi-step hydrolysis is an effective method to enhance the bioactivities of peptides (Phanturat et al., 2010). The proteases sequence in multi-step hydrolysis also influences the bioactivities of the resultant peptides (Klompong et al., 2007).

Bioactive peptides exhibit a diverse range of biological functions, such as antioxidant, antihypertensive, antimicrobial activities, etc. Among these, the most extensively documented bioactivities associated with FPH are antioxidant and antihypertensive properties (Zamora-Sillero et al., 2018). Some peptides exhibited multiple bioactivities, both antioxidant and antihypertensive activities such as a single peptide from FPH derived from *Merluccius productus* (Samaranayaka et al., 2010). Additionally, peptide fractions obtained from tuna gelatin hydrolysate prepared by Alcalase had both antimicrobial and antioxidant properties (Guillén et al., 2010).

Antioxidant peptides derived from fish processing by-products tend to have a LMW and are predominantly produced through enzymatic hydrolysis. Type of protease is a crucial factor affecting the antioxidant activity. Papain is the most frequently used enzyme for FPH production (Cunha & Pintado, 2022). Studies were primarily focused on the ability of these peptides from FPH to scavenge hydroxyl, peroxy, DPPH and ABTS radicals, with relatively fewer evaluating their protective effects on DNA and Oxygen Radical Absorbance Capacity (ORAC) activity (Cunha & Pintado, 2022). Some researchers have noted that hydrophobic amino acids may enhance the efficiency of antioxidant peptides (Zamora-Sillero et al., 2018). Additionally, different marine sources yield antioxidant peptides with varying capacities to scavenge free radicals. These peptides often contains amino acids like lysine, tyrosine, proline, and histidine (Wang et al., 2008). Antioxidant activities of selected FPHs from several fish species are tabulated in Table 1.

The inhibition of Angiotensin-Converting Enzyme (ACE) has emerged as a primary target in hypertension treatment. Several peptides have ACE-inhibiting activity, acting as natural hypotensive ACE inhibitors due to their high competitive affinity with the ACE active site (Lee et al., 2010). ACE-inhibiting peptides have been isolated from fish processing by-products such as skipjack roe (Intarasirisawat et al., 2013), salmon pectoral fin (Ahn et al., 2012), Atlantic salmon skin (Gu et al., 2011), tuna frame (Lee et al., 2010), sardinelle by-products (Bougatef et al., 2008), yellowfin sole frame (Jung et al., 2006b), Alaska pollock skin (Byun & Kim, 2001) and Tub gurnard skin (Bougatef et al., 2023).

Antimicrobial peptides are also prevalent in FPH, typically containing amino acids less than 50 with MW below 10 kDa (Sinaga et al., 2023). These peptides play a crucial role in natural defenses and not only have the ability to kill microorganisms but also modulate inflammatory responses (Yeaman & Yount, 2003).

Nearly all fish antimicrobial peptides functions against both Gram-positive and Gram-negative strains (Najafian & Babji, 2012). Additionally, Ennaas et al. (2015) demonstrated that FPH from mackerel by-products exhibited antibacterial activity against Gram-negative (*E. coli*) and Gram-positive (*L. innocua*) strains. Hasani et al. (2022) investigated the antioxidant, anti-cancer and antimicrobial properties of hydrolyzed Indian mackerel protein (*Rastrelliger kanagurta*) using two enzymes, alcalase and flavourzyme. It was found that highest antioxidant, anticancer and antimicrobial properties were observed in FPH produced by alcalase for 30 min.

These antimicrobial peptides derived from fish and fish processing byproducts hold great promise for new antibiotic advancement in both pharmaceuticals and food industries (Kim & Wijesekara, 2010).

Furthermore, FPH has a wide range of bioactive properties including antiatherosclerosis, anticancer, anticoagulant, antidiabetic, calcium binding, and anti-inflammatory activities (Cunha & Pintado, 2022; Gao et al., 2021; Zamora-Sillero et al., 2018). The bioactivities of selected FPHs from different fish species are given in Table 2. FPH derived from *Liza abu* has anti-cancer properties against colon cancer cells and Hela cancer cells (Shahosseini et al., 2022). Taroncher et al. (2023) reported that natural antioxidants (vitamins C and E, quercetin, and resveratrol) improved the cell proliferation of FPH in Caco-2 cells. FPH derived from *Trachinotus ovatus* showed significant antidiabetic effects, which alleviated body weight loss, polyphagia, blood glucose elevation and insulin secretion decline in diabetic mice (Wan et al., 2023).

### **Bitterness and Debittering**

Bitterness is a significant challenge that impacts the overall sensory acceptability of FPH. Bitterness in FPH is attributed to the presence of hydrophobic amino acids, including Leucine (Leu), Valine (Val), Tyrosine (Tyr), Isoleucine (Ile), Phenylalanine (Phe), and Tryptophan (Trp). Some factors such as enzyme types, DH, and MW also determine bitterness of FPH (Zhou et al., 2023). The hydrolysis process often exposes the concealed hydrophobic amino acids, contributing to the bitterness (Idowu & Benjakul, 2019). Peptides with bulky hydrophobic groups towards their C-terminal have been identified as the primary contributor to bitterness (Sinthusamran et al., 2020b). The average free energy (Q values) necessary for the transition of amino acid chains from ethanol to water has been used as a bitterness index. Bitter peptides have Q value greater than 1,400 kcal/mole with MW below 6 kDa, while non-bitter peptides exhibit Q value lesser than 1,300 kcal/mole. This practical relationship between average hydrophobicity and bitterness is termed as Q rule. The rule related with average hydrophobicity is implemented for bitterness prediction (Idowu & Benjakul, 2019).

**Table 1.** Antioxidant activities of selected FPHs from different fish species and their isolated peptides.

Fish species and body part used for FPH	Enzymes used for FPH production	Antioxidant activities exhibited	Hydrolysis condition	References
Muscle from Mahseer ( <i>Tor tambroides</i> )	Papain and bromelain	Antioxidant activities and antibacterial activities	Papain and bromelain: pH 7.0 and Temperature 50 °C	Sinaga et al. (2023)
Bighead Carp ( <i>Hypophthalmichthys nobilis</i> )	Flavourzyme	Antioxidant activities	pH 6.0, Temperature 50 °C and Time 60 min	Alahmad et al. (2023)
Salmon ( <i>Salmo salar</i> ) and mackerel ( <i>Scomber scombrus</i> ) heads and backbones	FoodPro PNL	Antioxidant activities	pH 7.0, Temperature 50 °C and Time 60 min	De la Fuente et al. (2023)
Featherback ( <i>Chitala ornata</i> ) skin	Alcalase	Antioxidant activities	pH 7.5, Temperature 55 °C and Time 5 h	Vo et al. (2023)
<i>Liza abu</i> muscle	Alcalase and flavourzyme	DPPH free radical scavenging activity, ferric reducing power, Fe <sup>2+</sup> chelating activity, and ABTS free radical scavenging activity	Alcalase and flavourzyme: pH 7 and pH 8.5 Temperature 58 °C	Shahosseini et al. (2022)
Skipjack tuna ( <i>Katsuwonus pelamis</i> ) roe	Flavourzyme	Antioxidant activities	pH 7.5, Temperature 45 °C and Time 8 h	Wang et al. (2022)
Klunzinger's mullet ( <i>Liza klunzingeri</i> )	Alcalase	Antioxidant activities	pH 8.0, Temperature 50 °C and Time 3 h	Rabiei et al. (2022)
Atlantic codfish ( <i>Gadus morhua</i> ) frames	Alcalase	Antioxidant and antihypertensive activities	Alcalase: pH 8.35 Temperature 56.8 °C Time 3 h	Rodrigues et al. (2021)
Pacific thread herring ( <i>Opisthonema libertate</i> )	Alcalase® 2.4 L and subtilisin A	ABTS, FRAP, and DPPH radical scavenging activity	Alcalase and subtilisin A pH 9.0 and Temperature 55 °C Time 1h	Martínez-Montaño et al. (2021)
Snakehead fish ( <i>Channa striata</i> ) viscera	Papain	DPPH and ABTS radical scavenging activity	Papain: pH 7.0 and Temperature 55 °C	Agustin et al. (2021)
Frame of salmon ( <i>Salmon salar</i> )	Alcalase and papain	DPPH radical scavenging ability, ABTS, ferrous reducing antioxidant power (FRAP), metal chelating activity and oxygen radical antioxidant capacity (ORAC)	Alcalase: pH 8.0 Temperature 60°C Time 15 min Papain: pH 7.0, Temperature 40°C Time 15 min	Idowu et al. (2019)
Muscle of <i>Nemipterus hexodon</i>	Pepsin from skipjack tuna	DPPH radical scavenging activity, ABTS and FRAP	Pepsin extract: pH 2 Temperature 50°C Time 15 min	Nalinanon et al. (2011)
Scale gelatin from <i>Oreochromis niloticus</i>	Alcalase, Pronase E, trypsin and pepsin	DPPH radical scavenging activity, hydroxyl radical scavenging activity and superoxide radical anion scavenging activity	Alcalase, Pronase E & trypsin: pH 8.0 Pepsin: pH 2 Temperature 60°C Time 4 h	Ngo et al. (2010)
Viscera from <i>Parastromateus niger</i>	Pepsin, trypsin and α-chymotrypsin	DPPH radical scavenging activity, FRAP and metal chelating activity	Pepsin: pH 2.5 trypsin & α-chymotrypsin: pH 6.5 Temperature 37°C Time 2 h	Jai Ganesh et al. (2011)
Meat of <i>Misgurnus anguillicaudatus</i>	Papain	DPPH radical scavenging activity, linoleic acid autoxidation inhibition activity, hydroxyl radical scavenging activity, and Cu <sup>2+</sup> ion chelating activity	Papain: pH 7 Temperature 55°C Time 4 h	You et al. (2010)
Muscle from <i>Decapterus maruadsi</i>	Flavourzyme	DPPH radical scavenging activity, FRAP and metal chelating activity	Flavourzyme: pH 8 Temperature 50°C Time 1 h	Thiansilakul et al. (2007)
Body meat of <i>Selaroides leptolepis</i>	Alcalase and flavourzyme	DPPH radical scavenging activity, FRAP, and reducing power activity	Alcalase: pH 8.5 Temperature 60°C Flavourzyme: pH 7 Temperature 50°C DHs: 5, 15 & 25%	Klompong et al. (2007)
Backbones of Tuna	Alcalase, α-chymotrypsin, Neutrase, papain, pepsin and trypsin	DPPH radical scavenging activity, lipid peroxidation inhibition activity, superoxide radical scavenging activity and hydroxyl radical scavenging activity	Alcalase: pH 7, Temperature 50°C Trypsin, α-Chymotrypsin & Neutrase: pH 8.0, Temperature 37°C Pepsin: pH 2.0, Temperature 37°C Papain: pH 6.0, Temperature 37°C	Je et al. (2007)
Protamine derived from Salmon ( <i>Salmon salar</i> )	Pancreatin	DPPH radical scavenging activity, hydroxyl radical scavenging activity and superoxide anion radical scavenging activity	Pancreatin: pH 8.5 Temperature 37°C Time 5h	Wang et al. (2008)



**Table 2.** Bioactivities of selected FPHs from different fish species.

Fish species and body part used for FPH	Bioactivities	Reference
Silverfish ( <i>Trachinotus ovatus</i> ) muscle protein	Antidiabetic	Wan et al. (2023)
Tub Gurnard ( <i>Chelidonichthys lucerna</i> ) Skin	Antihypertensive	Bougatef et al. (2023)
Yellowfin tuna frame ( <i>T. albacares</i> )	Myeloperoxidase inhibitory activity	Cai et al. (2022)
<i>Liza abu</i> muscle protein	Anti-cancer	Shahosseini et al. (2022)
Farm Rainbow Trout	Antihypertensive	Pérez-Escalante, et al. (2022)
Pacific thread herring ( <i>Opisthonema libertate</i> )	Antihypertensive	Martínez-Montaña et al. (2021)
Stickwater	Antiproliferative	Kandyliari et al. (2020)
Meagre ( <i>Argirosomus Regius</i> ) bones, gills and skin	Neuroprotective effect	(Lee et al., 2019)
Pacific hake ( <i>Merluccius productus</i> )	Antidiabetic	Harnedy et al. (2018)
Blue whiting ( <i>Micromesistius poutassou</i> ) muscle	Antibacterial	Wald et al. (2016)
Rainbow trout viscera	Neuroprotective effect	Chai et al. (2016)
Lanternfish ( <i>Benthosema pterotum</i> )	Antiproliferative	Chalamaiah et al. (2015)
Rohu ( <i>Labeo rohita</i> ) roe	Anti-inflammatory	Ahn et al. (2015)
Salmon pectoral fin	Antiproliferative	Song et al. (2014)
Half-fin anchovy ( <i>Setipinna taty</i> )	Anti-atherosclerotic	Parolini et al. (2014)
Norwegian salmon	Anticoagulant	Nasri et al. (2012)
Goby fish muscle protein	Antitumor	Chang et al. (2011)
Tilapia ( <i>Oreochromis mossambicus</i> )	Antihypertensive	Lee et al. (2010)
Tuna frame protein	Antimicrobial	Guillén et al. (2010)
Tuna skin	Antihypertensive	Bougatef et al. (2008)
Sardinelle heads and viscera	Calcium binding	Jung and Kim (2007)
Hoki ( <i>Johnius belengerii</i> ) frame	Antimicrobial	Dong et al. (2002)
Loach ( <i>Misgurnus anguillicaudatus</i> )		

Various methods including liposomal encapsulation, plastein reaction, enzymatic hydrolysis with exopeptidase, chromatographic separation, and extraction with cyclodextrin, activated carbon treatment, alcohol and Maillard reaction have been discovered to reduce or remove the bitterness of FPH (Idowu & Benjakul, 2019; Sharma et al., 2023a; Sharma et al., 2023b; Singh et al., 2020; Sinthusamran et al., 2020b). These approaches effectively reduce bitterness and enhance the overall taste profile. However, it is important to note that these methods may lead to structural changes and potential loss of certain peptides (Sharma et al., 2023b).

### Applications of FPH

Bioactive peptides find applications across diverse industrial sectors including food, cosmetics, and pharmaceuticals. In the food industries, many dairy merchandise fortified with bioactive peptides have been already marketed (Nirmal et al., 2022; Sasidharan & Venugopal, 2020). FPH has been explored for integration into a range of products including fish meat items, cereals, crackers, and desserts. Several brands have successfully introduced products containing these hydrolysates (Chalamaiah et al., 2012). Antioxidant peptides, crucial for retarding lipid oxidation, have gained significant attention in food packaging, as seen in products like sauces, meats, fish, and nuts (Gómez-Estaca et al., 2014). In Thailand, Japan and other countries, various products featuring FPH and peptides are available in the market, advertised as functional foods like powdered soups, dietary supplements, beverages, etc. Over sixty peptides have received FDA

approval for medicinal application, with numerous preclinical studies underway (Rivero-Pino et al., 2020).

### Food Applications

Protein hydrolysates have attracted substantial interest in recent years because of their improved functionalities and high nutritional value resulted from the generation of free amino acids and small peptides through hydrolysis (Benjakul et al., 2014). FPH have been documented as a superior source of protein for human nutrition due to their balanced amino acid composition and high absorption in gastrointestinal tract (Cunha & Pintado, 2022). High solubility of FPH across a wide range of pH renders them highly versatile for various food applications (Dinakarkumar et al., 2022).

FPH has been extensively used in food products. Their excellent interfacial properties make them become valuable emulsifying agents in products such as dressings, margarine, and meat batter. They also contribute to the creation and stabilization of foam-containing products like mousse, meringues, and whipped cream (Nirmal et al., 2022). Khan et al. (2003) employed FPH as a cryoprotectant to inhibit denaturation of proteins in lizardfish surimi during frozen stage. Damodaran (2007) showed the ability of gelatin hydrolysate ability to hinder ice crystallization in ice cream. Additionally, FPH from mince obtained from gutted and headed red hake or frame from red hake produced with Flavozyme at natural pH (6.8) and a fish /water ratio of 5:2 were employed as a natural flavor stock (Imm and Lee 1999). Beverage fortified with FPH from blue whiting (*Micromesistius poutassou*) (Egerton

et al., 2018), instant coffee added with galactose-fish skin gelatin hydrolysate (Karnjanapratum & Benjakul, 2017), and protein-rich ice cream fortified with FPH from saithe (*Pollachius virens*) (Shaviklo et al., 2011) have been documented. Bighead carp (*Hypophthalmichthys nobilis*) hydrolysate was prepared using alcalase and hydrolytic enzymes extracted from *Oncorhynchus mykiss*. The resulting hydrolysate could be used in the development of active packaging as antimicrobial and antioxidant components (Naghdi et al., 2023).

### Pharmaceutical and Nutraceutical Applications


The potential of FPH in lowering the risk of various illnesses has been confirmed, such as reducing plasma cholesterol levels associated with cardiovascular diseases and lowering cell proliferation in cell lines of human breast cancer (Picot et al., 2006). FPH from various fish species demonstrated higher inhibition of cell lines in human breast cancer, MCF-7/6 and MDA-MB-231 (Picot et al., 2006). Furthermore, calcium-binding bioactive peptides obtained from Alaska pollock (*Theragra chalcogramma*) via pepsin hydrolysis and hoki (*Johnius belengerii*) frame offer a potential alternative for individuals with lactose intolerance, with calcium rich foods or calcium added fruit juices serving as dairy substitute (Jung & Kim, 2007). Peptides from FPH have also demonstrated the ability to enhance calcium absorption (Jung et al., 2006a; Kim & Mendis, 2006a). Moreover, clinical studies recommend that the ingestion of FPH can alleviate pain in osteoarthritis patients (Wu et al., 2004). Collagen hydrolysate combined with a calcitonin-rich diet was more effective in reducing breakdown of bone collagen compared to a calcitonin-rich diet alone (Moskowitz, 2000). Fish collagen hydrolysate ingestion has also been augmented with the synthesis of extracellular matrix macromolecules by chondrocytes (Bello & Oesser, 2006). Additionally, fish skin collagen hydrolysates from salmon and trout have been found to influence metabolism and lipid absorption in rat models. (Saito et al., 2009). FPH from Asian seabass skin also showed wound healing properties bone strengthening and skin nourishment effects in *in vitro* cell line studies (Benjakul et al., 2018a; Chotphruethipong et al., 2021). While FPH presents numerous potential benefits, there are currently limited commercial products containing FPH existing for human consumption. Nonetheless, products with health-promoting properties derived from aquatic resources, mainly FPH, hold promising potential in the market.

### Production and Characteristics of Fish Biocalcium

Fish bone and scale, which are important raw material for biocalcium, contain non-collagenous constituents, including meat proteins and lipids, etc. Pretreatments with single or several steps are applied to

diminish the undesirable components before processing to augment the purity of produced fish biocalcium. The first step of pretreatment is the remaining meat removal and thorough cleaning prior to process in further pretreatments (Figure 3) (Benjakul et al., 2017b). Softening and reduction of material size are also important for facilitating the efficacy of fish biocalcium production. Pretreatment with alkaline solution such as sodium hydroxide (NaOH) is normally used for removal of non-collagenous protein, pigments and lipids. Increasing alkaline concentration and pretreatment time are able to augment proteins solubilization or unfolding. NaOH at concentration of 2 M has been used for pretreatment of fish bones and scales (Benjakul et al., 2017b; Nilsuwan et al., 2023). Benjakul et al. (2018b) cleaned the pre-cooked skipjack tuna bone with high pressure water jet at 120 bar for 2 min, followed by treatment with NaOH solution (2 M) at temperature of 50 °C for 30 min with bone/solution ratio of 1:10 (w/v). The aforementioned alkaline pretreatment is able to remove the remaining meat and lipids, leading to the augmenting Ca and P contents in biocalcium powder (Benjakul et al., 2018b). Benjakul et al. (2017a) found that biocalcium powders from pre-cooked tonggol (*Thunnus tonggol*) and yellowfin (*Thunnus albacores*) tuna bones, pretreated with NaOH solution (2 M) at temperature of 50 °C for 30 min with bone/solution ratio of 1:10 (w/v), had Ca contents of 26.76 and 26.73% and P contents of 12.7 and 12.74%, respectively. The Ca:P ratio was nearby 2:1, indicating compatibility to that found in human bones. Idowu et al. (2020) found that biocalcium powder from salmon (*Salmo salar*) frame pretreated with 2 M NaOH for 40 min at 50 °C with a bone/solution ratio of 1:10 (w/v) had 27.32% Ca and 13.22% P with the Ca/P mol ratio of 1.60. The uses of NaOH solutions (0.5 - 1.0 M) and ethanol (40 - 60%, v/v) for processing of bones and skeletons from Indian oil sardine (*Sardinella longiceps*) and small-headed ribbonfish (*Trichiurus savala*) yielded the powders with calcium content ranging from 18.91% to 32.73%. (Logesh et al., 2012). The NaOH solution (2 M, 10 min) can also remove proteins effectively and minimize the loss of collagen as indicated by high hydroxyproline content in biocalcium from Asian sea bass (*Lates calcarifer*) scales (Nilsuwan et al., 2023). In addition, the alkaline pretreatment process can diminish undesirable volatile compounds in biocalcium. Idowu et al. (2019a) documented that lower abundance of volatile compounds such as 2-propanone and 2-butanone was obtained for biocalcium powders from salmon frame after treatment with 2 M NaOH for 40 min at 50 °C with a bone/solution ratio of 1:10 (w/v). However, strong condition should be obviated. Fish frame leftover from filleting process, still contains flesh. This meat rich in protein can be removed by enzymatic hydrolysis (Idowu et al., 2019a). The enzymatic hydrolysis using protease is widely implemented to separate non-collagenous protein from fish frame or trimming (Figure 3) (Idowu et al., 2020; Sinthusamran et al., 2020a). The enzymatic





**Figure 3.** Flowchart for production of biocalcium from pre-cooked and raw frame/backbone

hydrolysis is able to hydrolyze the myofibrillar proteins into hydrolysate solution, which can be separated from the bone solid easily (Idowu et al., 2020).

Regarding fat removal, the solvents are commonly used for treatment of raw materials. Hexane is generally employed for fat removal from biocalcium, which can reduce fat in raw material effectively (Benjakul et al., 2017a; Benjakul et al., 2017b; Idowu et al., 2020). Lipid content of biocalcium powders from pre-cooked tongol tuna bone, pre-cooked yellowfin tuna bone, and salmon frame bone defatted with hexane were 0.26, 0.19 and 0.33%, respectively (Benjakul et al., 2017a; Idowu et al., 2020). Moreover, mixed solvents can also be used for defatting process. Pudtikajorn et al. (2023) documented that biocalcium powder with lipid content of 0.09% was obtained from skipjack tuna eyeball scleral cartilage defatted with the mixture of hexane and isopropanol (1:1) for 6 cycles. The efficiency of defatting process with the hexane and isopropanol mixture (1:1) was 99.63% for lipid reduction (Pudtikajorn et al., 2023).

Whiteness of biocalcium powder is related to customer acceptability. Bleaching using sodium hypochlorite (NaOCl) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) alone or combination can improve the whiteness of biocalcium. *L\**-value of biocalcium powders from pre-cooked tongol and yellowfin tuna bones bleached with 2.5% (v/v) NaOCl for 30 min followed by 2.5% (v/v) H<sub>2</sub>O<sub>2</sub>

for 60 min was 93.92 and 92.52, respectively (Benjakul et al., 2017a). Moreover, the highest *L\**-value (92.26) was observed when H<sub>2</sub>O<sub>2</sub> alone was used as bleaching agent for production of Asian sea bass (*Lates calcarifer*) backbone biocalcium (Wijayanti et al., 2021b).

Grinding is another crucial step for making fish biocalcium powder. Since the bone matrix is strong and compact in structure, heating under high pressure could assist softening of bone. This could facilitate the size reduction process. Softening process can enhance grinding efficiency of raw materials into powders. Wijayanti et al. (2021a) documented that high pressure heating with the aid of autoclave for 90 min lowered hardness of Asian sea bass backbone from 57,000 g force to around 3000 g force. Treated samples could be ground into the powder with ease. Choi et al. (1998) reported that the autoclaving at 40 min resulted in Alaska pollock (*Theragra chalcogramma*) and cod (*Gadus chalcogrammus*) powders having appreciable soluble calcium content. Longer time of autoclave had no impact on the mineral yield and soluble calcium ratio (Choi et al., 1998). Apart from steaming, different acids have been used for treatment of Alaska pollock frame prior to extraction of calcium. In general, coarse ground particles can provide sandiness of mouth feel. Ball milling process has been widely used for production of fish biocalcium from tuna bone, salmon bone and

seabass bone (Benjakul et al., 2017b; Idowu et al., 2020; Wijayanti et al., 2021b). Particle size of bio-calcium lower than 75 µm was generally obtained when the bio-calcium powders from tuna eyeball scleral were sieved using screen having mesh no. 200 (Pudtikajorn et al., 2023).

### Bioavailability of Fish Bio-calcium

The exploitation of fish bone as functional materials is less, though it contains nutraceutical and bioactive molecules. This was due to its low solubility (Kim & Jung, 2012). The solubilization is required for the absorption of calcium in the body. Calcium salt is mostly precipitated in the ileum caused by alkaline conditions, however some calcium ions are present in solution (Kim & Jung, 2012). Typically, calcium is rarely absorbed in the intestine. Gastric contents after calcium is dissolved in the stomach move into the small intestine, where calcium is absorbed. Bicarbonate ( $\text{HCO}^{-3}$ ) is secreted into the intestine to neutralize the contents. The concentration of calcium available for absorption in the jejunum and ileum, which maintain a pH of approximately 7, is therefore expected to be lower than that dissolved in the stomach (Goss et al., 2010). The absorption of calcium is complicated determined by local solubility, the sojourn time in the particular intestinal segment, and the rate of transepithelial movement (Duflos et al., 1995). Malde et al. (2010) reported that young healthy men had absorption of calcium from cod and salmon bones at 21.9 and 22.5%, respectively. Tuna bone powder exhibited higher calcium availability (53.93%) than most calcium salts including calcium carbonate, calcium lactate, calcium citrate and milk powder (Nemati et al., 2016). Based on the the amount of calcium transported across the Caco-2 monolayer, Idowu et al. (2019a) documented that bioavailability of bio-calciums obtained from salmon frames treated with and without alkaline solution were 43.02 and 38.13%, respectively. Hake fish bone (HFB) powder exhibit non cytotoxic effect on the cell line (SaOS-2 cells) (Flammini et al., 2016). HFB powder also had comparable efficacy to commercial calcium tablets on rat bone mineralization (Flammini et al., 2016). Additionally, peptides act as calcium carrier, which enhance calcium absorption at small intestine (Goss et al., 2010). Jung et al. (2006a) found that peptides could bind with calcium and enhance solubility of calcium salts at neutral pH in *In vitro* study. The ovariectomized rats supplemented with calcium binding peptides had higher calcium retention and lower loss of bone mineral (Jung et al., 2006a). Oligophosphopeptides, which are generated via enzymatic hydrolysis of hoki (*Johnius belengerii*) frames by heterogeneous enzyme extracted from the intestine of bluefin tuna, could help solubilization of calcium and improve calcium bioavailability (Jung et al., 2006a).

### Fortification of Bio-calcium in Drink and Foods

Fish bio-calcium has been applied as dietary supplement such as commercial tablets. Fish bio-calcium can also be incorporated or fortified in various foods. Tuna bone peptide (TBP) and tricalcium phosphate (TCP) were added into bakery products i.e. cookies and bread as a calcium supplement (Nemati et al., 2016). A higher *in vitro* bioavailability was found for cookies fortified with TBP, compared to that of bread fortified with TBP. Cookies fortified with TBP and TCP had calcium bioavailability at 38.9% and 39.5%, respectively, while bioavailability of bread containing TBP and TCP were 36.7% and 37.4%, respectively. Cookies and bread containing TBP or TCP had similar overall acceptability scores in the range of 7.76 – 7.80 and 7.30 – 7.48, respectively (Nemati et al., 2016). In addition, Idowu et al. (2019a) prepared bio-calcium powders from salmon (*Salmo salar*) frames via hydrolysis with alcalase prior to treatment with NaOH solution (2 M, 50°C, 40 min). The obtained bio-calcium (BC) was fortified in whole wheat crackers along with protein hydrolysate (PH) at wheat flour substitution level of 16.67%. Fortification of BC and PH powders had a profound impact on compositions of resulting whole wheat cracker. Increases by 17- and 8-fold were obtained for calcium and phosphorus contents, respectively, compared to those found in the control (without BC and PH). This was associated with higher ash content of the BC/PH fortified sample (Idowu et al., 2019a). Furthermore, Karnjanapratum & Benjakul (2018) developed coconut oil based cookie fortified with tuna bone bio-calcium. The coconut oil was used to substitute the shortening at 65% and 100%. Bio-calcium from tuna bone was added into the batter at 8.5%, 12.0% and 15.5% (w/w). Addition of bio-calcium powder at 12.0% had no effect on sensory properties of cookie containing coconut oil as shortening replacer. The developed cookie showed higher calcium and phosphorus contents as well as higher saturated fat, ash, and protein contents but lower energy value, compared to that of cookie without bio-calcium and coconut oil. The significant increases in protein and ash contents in developed cookie were related to high protein (24.26%) and ash (72.20%) of tuna bio-calcium powder (Benjakul et al., 2017b). It is worthnoting that tuna bone bio-calcium powder was effectively used for calcium fortification in cookie. Nevertheless, the replacement of shortening by coconut oil showed higher content of saturated fat with lower mono- and poly-unsaturated fats in cookie. In general, coconut oil is abundant in medium chain saturated fatty acids (MCFAs) such as lauric acid (C12:0) and myristic acid (C14:0) (Karnjanapratum & Benjakul, 2018).

Bio-calcium can be used in surimi products to increase nutritional value and improve surimi gel properties. Wijayanti et al. (2021c) incorporated bio-calcium powder prepared from bone of Asian sea bass (ASBB) in threadfin bream (*Nemipterus* sp.) surimi gel at levels of 0 – 10% (w/w). The addition of 8% (w/w)

ASBB possessed high breaking force and high overall likeness score in resulting surimi gel. The 8% (w/w) ASBB added surimi gel also had higher ash, calcium, and phosphorus content as well as higher compactness and finer microstructure, compared to the control without ASBB (Wijayanti et al., 2021c). Additionally, the addition of biocalcium along with other calcium salts can enhance the improvement of surimi gel properties. Wijayanti et al. (2021d) prepared surimi gel from threadfin bream added with fish bone biocalcium (8%) in combination with calcium chloride (CC) or calcium lactate (CL) at different levels (0.1–0.6%, w/w). The highest breaking force along with the lowest expressible moisture content and higher whiteness were obtained for gel added with 8% biocalcium together with 0.6% CL. The aforementioned gel also had elastic texture as confirmed by the highest  $G'$  and denser microstructure (Wijayanti et al., 2021d). Furthermore, fish tofu produced from threadfin bream surimi fortified with 7.5% (w/w) skipjack tuna eyeball scleral cartilage biocalcium (SCBC) also had a denser microstructure with greater ash and calcium contents than the control without SCBC incorporation (Putdikajorn et al., 2022).

Biocalcium has also been fortified into food emulsion products. Asian sea bass bio-calcium (ASBC) fortified mayonnaise had the enhanced firmness, consistency and cohesiveness with augmenting ASBC levels (0%–10%). Mayonnaise sample added with ASBC showed greater  $G'$ ,  $G''$ , viscosity and shear stress value. The increases in lightness ( $L^*$ ) and total color difference ( $\Delta E^*$ ), but decreases in  $a^*$  and  $b^*$ -values were found in mayonnaise with increasing ASBC amount. Lower creaming index of mayonnaise added with ASBC was noted as indicated by lower droplet size and higher zeta potential. Dense and small droplet sizes were observed for mayonnaise fortified with ASBC at 2.5 with higher protein, ash and calcium content than the control (without ASBC) (Wijayanti et al., 2023). Biocalcium was also added into fish spread. Wijayanti et al. (2022a) prepared the Indian mackerel fish spread incorporated with Asian sea bass bone biocalcium (BC) (0–10%) and potato starch (2–4%). The obtained fish spreads were generally increased in firmness, consistency and cohesiveness as BC level was augmented (Wijayanti et al., 2022a). Shear thinning flow behaviour was generally found in all fish spreads. Fish spread containing 10% BC and 4% starch had higher acceptability without sandy mouth feel. Fish spread had higher ash content with less moisture, protein and lipid contents than those without BC. After *in vitro* gastrointestinal tract system of fish spread fortified with 10% BC, calcium solubility was augmented and higher solubility of calcium was obtained, than that of BC (Wijayanti et al., 2022a).

Nevertheless, biocalcium can also be added into fruit juice. Wijayanti et al. (2022b) prepared the 4% (w/v) soluble Asian sea bass bone bio-calcium (SBC) in 9% (w/v) citric acid prior to fortify into sterilized apple juices. The apple juice samples had varying in pH, turbidity, brown color and calcium content. Lower pH

along with higher turbidity, brown color and calcium content was observed as the levels of SBC was increased. No impact of SBC fortification on likeness score for all quality attributes as well as sandy mouthfeel. Nevertheless, apple juice fortified with excessive SBC possessed lower flavor, taste and overall likeness. Calcium content of apple juice fortified with SBC up to 1% was augmented by 6.13-fold with no detrimental effects on sensory property (Wijayanti et al., 2022b).

## Conclusion

Value-added products, especially functional ingredients or nutraceuticals can be produced from fish processing leftovers, which are generally wasted or used for production of low value products. For better utilization of bone or frame, they can be used for production of protein hydrolysate with bioactivities as well as biocalcium, which has higher availability with absorption. Both protein hydrolysate and biocalcium could be fortified in drink or food products, in which food quality could be improved. More importantly, the functional ingredient could play a major role in health promotion/benefit due to their nutraceutical property.

## Ethical Statement

Ethics approval was not required for this research.

## Funding Information

This study was funded by Prachayacharn Grant from Prince of Songkla University, Hat Yai, Thailand (Grant no. AGR6602079N). This research was also supported by the Postdoctoral Fellowship from Prince of Songkla University to Dr. Umesh Patil.

## Author Contribution

Umesh Patil: Conceptualization, Visualization and Writing-original draft; Krisana Nilsuwan: Conceptualization, Writing-review and editing; Soottawat Benjakul: Supervision, Writing-review and editing. All authors read and approved the final manuscript.

## Conflict of Interest

The authors declare no conflict of interest.

## Acknowledgements

The authors gratefully acknowledge the International Center of Excellence in Seafood Science and Innovation, Faculty of Agro-Industry, Prince of Songkla University for all support.

## References

- Agustin, V., Husni, A., & Putra, M. (2021). Antioxidant activity of protein hydrolysate from snakehead fish (*Channa striata*) viscera obtained by enzymatic process. *IOP Conference Series: Earth and Environmental Science*, 919, 012046. DOI 10.1088/1755-1315/919/1/012046
- Ahn, C.-B., Cho, Y.-S., & Je, J.-Y. (2015). Purification and anti-inflammatory action of tripeptide from salmon pectoral fin byproduct protein hydrolysate. *Food Chemistry*, 168, 151-156. <https://doi.org/10.1016/j.foodchem.2014.05.112>
- Ahn, C.-B., Jeon, Y.-J., Kim, Y.-T., & Je, J.-Y. (2012). Angiotensin I converting enzyme (ACE) inhibitory peptides from salmon byproduct protein hydrolysate by Alcalase hydrolysis. *Process Biochemistry*, 47, 2240-2245. <https://doi.org/10.1016/j.procbio.2012.08.019>
- Alahmad, K., Noman, A., Xia, W., Jiang, Q., Xu, Y. (2023). Influence of the enzymatic hydrolysis using flavourzyme enzyme on functional, secondary structure, and antioxidant characteristics of protein hydrolysates produced from bighead carp (*Hypophthalmichthys nobilis*). *Molecules*, 28, 519. <https://doi.org/10.3390/molecules28020519>
- Anderson, J. J. B., & Garner, S. C. (1996). *Calcium and phosphorous nutrition in health and disease, Introduction*. CRC Press, New York, USA.
- Bello, A. E., & Oesser, S. (2006). Collagen hydrolysate for the treatment of osteoarthritis and other joint disorders: A review of the literature. *Current Medical Research and Opinion*, 22, 2221-2232. <https://doi.org/10.1185/030079906X148373>
- Benjakul, S., Karnjanapratum, S., & Visessanguan, W. (2018a). Hydrolysed collagen from *Lates calcarifer* skin: Its acute toxicity and impact on cell proliferation and collagen production of fibroblasts. *International Journal of Food Science & Technology*, 53, 1871-1879. <https://doi.org/10.1111/ijfs.13772>
- Benjakul, S., Mad-Ali, S., Senphan, T., & Sookchoo, P. (2018b). Characteristics of biocalcium from pre-cooked skipjack tuna bone as affected by different treatments. *Waste and Biomass Valorization*, 9, 1369-1377. <https://doi.org/10.1007/s12649-017-9927-8>
- Benjakul, S., Mad-Ali, S., & Sookchoo, P. (2017a). Characteristics of biocalcium powders from pre-cooked tongol (*Thunnus tonggol*) and yellowfin (*Thunnus albacares*) tuna bones. *Food Biophysics*, 12, 412-421. <https://doi.org/10.1007/s11483-017-9497-0>
- Benjakul, S., Mad-Ali, S., Senphan, T., & Sookchoo, P. (2017b). Biocalcium powder from precooked skipjack tuna bone: Production and its characteristics. *Journal of Food Biochemistry*, 41, e12412. <https://doi.org/10.1111/jfbc.12412>
- Benjakul, S., Sae-leaw, T., & Simpson, B. K. (2019). Byproducts from fish harvesting and processing. In K. B. Simpson, N. A. A. Aryee, & F. Toldrá (Eds.), *Byproducts from Agriculture and Fisheries* (pp. 179-217). John Wiley & Sons Inc. <https://doi.org/https://doi.org/10.1002/9781119383956.ch9>
- Benjakul, S., Yarnpakdee, S., Senphan, T., Halldorsdottir, S. M., & Kristinsson, H. G. (2014). Fish protein hydrolysates: production, bioactivities, and applications. In H. G. Kristinsson (Ed.), *Antioxidants and functional components in aquatic foods* (pp. 237-281). John Wiley & Sons Inc. <https://doi.org/https://doi.org/10.1002/9781118855102.ch9>
- Bougatef, A., Nedjar-Arroume, N., Manni, L., Ravallec, R., Barkia, A., Guillochon, D., & Nasri, M. (2010). Purification and identification of novel antioxidant peptides from enzymatic hydrolysates of sardinelle (*Sardinella aurita*) by-products proteins. *Food Chemistry*, 118, 559-565. <https://doi.org/10.1016/j.foodchem.2009.05.021>
- Bougatef, A., Nedjar-Arroume, N., Ravallec-Plé, R., Leroy, Y., Guillochon, D., Barkia, A., & Nasri, M. (2008). Angiotensin I-converting enzyme (ACE) inhibitory activities of sardinelle (*Sardinella aurita*) by-products protein hydrolysates obtained by treatment with microbial and visceral fish serine proteases. *Food Chemistry*, 111, 350-356. <https://doi.org/10.1016/j.foodchem.2008.03.074>
- Bougatef, H., de la Vega-Fernández, C., Sila, A., Bougatef, A., Martínez-Alvarez, O. (2023). Identification of ACE I-inhibitory peptides released by the hydrolysis of tub gurnard (*Chelidonichthys lucerna*) skin proteins and the impact of their in silico gastrointestinal digestion. *Marine Drugs*, 21, 131. <https://doi.org/10.3390/md21020131>
- Byun, H.-G., & Kim, S.-K. (2001). Purification and characterization of angiotensin I converting enzyme (ACE) inhibitory peptides from Alaska pollack (*Theragra chalcogramma*) skin. *Process Biochemistry*, 36, 1155-1162. [https://doi.org/10.1016/S0032-9592\(00\)00297-1](https://doi.org/10.1016/S0032-9592(00)00297-1)
- Cai, B., Wan, P., Chen, H., Huang, J., Ye, Z., Chen, D., Pan, J. (2022). Purification and identification of novel myeloperoxidase inhibitory antioxidant peptides from tuna (*Thunnus albacares*) protein hydrolysates. *Molecules*, 27, 2681. <https://doi.org/10.3390/molecules27092681>
- Chai, H.-J., Wu, C.-J., Yang, S.-H., Li, T.-L., & Pan, B. S. (2016). Peptides from hydrolysate of lantern fish (*Benthosema pterotum*) proved neuroprotective *in vitro* and *in vivo*. *Journal of Functional Foods*, 24, 438-449. <https://doi.org/10.1016/j.jff.2016.04.009>
- Chalamaiah, M., Hemalatha, R., & Jyothirmayi, T. (2012). Fish protein hydrolysates: proximate composition, amino acid composition, antioxidant activities and applications: A review. *Food Chemistry*, 135, 3020-3038. <https://doi.org/10.1016/j.foodchem.2012.06.100>
- Chalamaiah, M., Jyothirmayi, T., Diwan, P. V., & Dinesh Kumar, B. (2015). Antiproliferative, ACE-inhibitory and functional properties of protein hydrolysates from rohu (*Labeo rohita*) roe (egg) prepared by gastrointestinal proteases. *Journal of Food Science and Technology*, 52, 8300-8307. <https://doi.org/10.1007/s13197-015-1969-y>
- Chang, W.-T., Pan, C.-Y., Rajanbabu, V., Cheng, C.-W., & Chen, J.-Y. (2011). Tilapia (*Oreochromis mossambicus*) antimicrobial peptide, hepcidin 1-5, shows antitumor activity in cancer cells. *Peptides*, 32, 342-352. <https://doi.org/10.1016/j.peptides.2010.11.003>
- Chiodza, K., & Goosen, N. J. (2023). Evaluation of handling and storage stability of spray dried protein hydrolysates from sardine (*Sardina pilchardus*) processing by-products: Effect of enzymatic hydrolysis time, spray drying temperature and maltodextrin concentration. *Food and Bioprocess Processing*, 141, 1-22. <https://doi.org/https://doi.org/10.1016/j.fbp.2023.06.009>
- Choi, J.-D., Kim, D.-S., & Kim, J.-S. (1998). Preparation of calcium-based powder from fish bone and its characteristics. *Applied Biological Chemistry*, 41, 147-152.

- Chotphruethipong, L., Binlath, T., Hutamekalin, P., Aluko, R. E., Tapaamorndech, S., Zhang, B., & Benjakul, S. (2021). Impact of hydrolyzed collagen from defatted sea bass skin on proliferation and differentiation of preosteoblast MC3T3-E1 cells. *Foods*, 10, 1476. <https://doi.org/10.3390/foods10071476>
- Clemente, A. (2000). Enzymatic protein hydrolysates in human nutrition. *Trends in Food Science & Technology*, 11, 254-262. [https://doi.org/10.1016/S0924-2244\(01\)00007-3](https://doi.org/10.1016/S0924-2244(01)00007-3)
- Cunha, S. A., & Pintado, M. E. (2022). Bioactive peptides derived from marine sources: Biological and functional properties. *Trends in Food Science & Technology*, 119, 348-370. <https://doi.org/10.1016/j.tifs.2021.08.017>
- Damodaran, S. (2007). Inhibition of ice crystal growth in ice cream mix by gelatin hydrolysate. *Journal of Agricultural and Food Chemistry*, 55, 10918-10923. <https://doi.org/10.1021/jf0724670>
- de la Fuente, B., Aspevik, T., Barba, F. J., Kousoulaki, K., & Berrada, H. (2023). Mineral bioaccessibility and antioxidant capacity of protein hydrolysates from salmon (*salmo salar*) and mackerel (*scomber scombrus*) backbones and heads. *Marine Drugs*, 21(5), 294. <https://doi.org/10.3390/md21050294>
- Dinakarkumar, Y., Krishnamoorthy, S., Margavelu, G., Ramakrishnan, G., & Chandran, M. (2022). Production and characterization of fish protein hydrolysate: Effective utilization of trawl by-catch. *Food Chemistry Advances*, 1, 100138. <https://doi.org/10.1016/j.focha.2022.100138>
- Dong, X., Xu, H., Huang, K., Liou, Q., & Zhou, J. (2002). The preparation and characterization of an antimicrobial polypeptide from the loach, *Misgurnus anguillicaudatus*. *Protein Expression and Purification*, 26, 235-242. [https://doi.org/10.1016/S1046-5928\(02\)00514-4](https://doi.org/10.1016/S1046-5928(02)00514-4)
- Duflos, C., Bellaton, C., Pansu, D., & Bronner, F. (1995). Calcium solubility, intestinal sojourn time and paracellular permeability codetermine passive calcium absorption in rats. *The Journal of Nutrition*, 125, 2348-2355.
- Egerton, S., Culloty, S., Whooley, J., Stanton, C., & Ross, R. P. (2018). Characterization of protein hydrolysates from blue whiting (*Micromesistius poutassou*) and their application in beverage fortification. *Food Chemistry*, 245, 698-706. <https://doi.org/10.1016/j.foodchem.2017.10.107>
- Ennaas, N., Hammami, R., Beaulieu, L., & Fliss, I. (2015). Purification and characterization of four antibacterial peptides from protamex hydrolysate of Atlantic mackerel (*Scomber scombrus*) by-products. *Biochemical and Biophysical Research Communications*, 462, 195-200. <https://doi.org/10.1016/j.bbrc.2015.04.091>
- Flammini, L., Martuzzi, F., Vivo, V., Ghirri, A., Salomi, E., Bignetti, E., & Barocelli, E. (2016). Hake fish bone as a calcium source for efficient bone mineralization. *International Journal of Food Sciences and Nutrition*, 67, 265-273. <https://doi.org/10.3109/09637486.2016.1150434>
- Foh, M. B. K., Amadou, I., Foh, B. M., Kamara, M. T., & Xia, W. (2010). Functionality and antioxidant properties of tilapia (*Oreochromis niloticus*) as influenced by the degree of hydrolysis. *International Journal of Molecular Sciences*, 11, 1851-1869. <https://doi.org/10.3390/ijms11041851>
- Gao, R., Yu, Q., Shen, Y., Chu, Q., Chen, G., Fen, S., Yang, M., Yuan, L., McClements, D. J., & Sun, Q. (2021). Production, bioactive properties, and potential applications of fish protein hydrolysates: Developments and challenges. *Trends in Food Science & Technology*, 110, 687-699. <https://doi.org/10.1016/j.tifs.2021.02.031>
- Gaskin, D. J., & Ilich, J. Z. (2009). Lactose maldigestion revisited: Diagnosis, prevalence in ethnic minorities, and dietary recommendations to overcome it. *American Journal of Lifestyle Medicine*, 3, 212-218. <https://doi.org/10.1177/1559827609331555>
- Gómez-Estaca, J., López-de-Dicastillo, C., Hernández-Muñoz, P., Catalá, R., & Gavara, R. (2014). Advances in antioxidant active food packaging. *Trends in Food Science & Technology*, 35, 42-51. <https://doi.org/10.1016/j.tifs.2013.10.008>
- Goss, S., Prushko, J., & Bogner, R. (2010). Factors affecting calcium precipitation during neutralisation in a simulated intestinal environment. *Journal of Pharmaceutical Sciences*, 99, 4183-4191. <https://doi.org/https://doi.org/10.1002/jps.22131>
- Gu, R.-Z., Li, C.-Y., Liu, W.-Y., Yi, W.-X., & Cai, M.-Y. (2011). Angiotensin I-converting enzyme inhibitory activity of low-molecular-weight peptides from Atlantic salmon (*Salmo salar* L.) skin. *Food Research International*, 44, 1536-1540. <https://doi.org/10.1016/j.foodres.2011.04.006>
- Guillén, G., López Caballero, M. E., Alemán, A., Lacey, A. L. d., Giménez, B., & Montero García, P. (2010). Antioxidant and antimicrobial peptide fractions from squid and tuna skin gelatin. In E. L. Bihan & N. Koueta (Eds.), *Sea by-products as a real material: New ways of application* (pp. 89-115). Transworld Research Network.
- Halim, N., Yusof, H., & Sarbon, N. (2016). Functional and bioactive properties of fish protein hydrolysates and peptides: A comprehensive review. *Trends in Food Science & Technology*, 51, 24-33. <https://doi.org/10.1016/j.tifs.2016.02.007>
- Hansen, M., Thilsted, S. H., Sandström, B., Kongsbak, K., Larsen, T., Jensen, M., & Sørensen, S. S. (1998). Calcium absorption from small soft-boned fish. *Journal of Trace Elements in Medicine and Biology*, 12, 148-154. [https://doi.org/https://doi.org/10.1016/S0946-672X\(98\)80003-5](https://doi.org/https://doi.org/10.1016/S0946-672X(98)80003-5)
- Harnedy, P. A., Parthasarathy, V., McLaughlin, C. M., O'Keeffe, M. B., Allsopp, P. J., McSorley, E. M., O'Harte, F. P., & FitzGerald, R. J. (2018). Blue whiting (*Micromesistius poutassou*) muscle protein hydrolysate with *in vitro* and *in vivo* antidiabetic properties. *Journal of Functional Foods*, 40, 137-145. <https://doi.org/10.1016/j.jff.2017.10.045>
- He, S., Franco, C., & Zhang, W. (2013). Functions, applications and production of protein hydrolysates from fish processing co-products (FPCP). *Food Research International*, 50, 289-297. <https://doi.org/10.1016/j.foodres.2012.10.031>
- Idowu, A. T., & Benjakul, S. (2019). Bitterness of fish protein hydrolysate and its debittering prospects. *Journal of Food Biochemistry*, 43, e12978. <https://doi.org/10.1111/jfbc.12978>
- Idowu, A. T., Benjakul, S., Sae-Leaw, T., Sookchoo, P., Kishimura, H., Suzuki, N., & Kitani, Y. (2019a). Amino acid composition, volatile compounds and bioavailability of biocalcium powders from salmon frame as affected by pretreatment. *Journal of Aquatic Food Product Technology*, 28, 772-780. <https://doi.org/10.1080/10498850.2019.1639235>
- Idowu, A. T., Benjakul, S., Sinthusamran, S., Sae-leaw, T., Suzuki, N., Kitani, Y., & Sookchoo, P. (2020). Effect of alkaline treatment on characteristics of bio-calcium and hydroxyapatite powders derived from salmon bone. *Applied Sciences (Switzerland)*, 10, 4141.

- <https://doi.org/10.3390/APP10124141>
- Idowu, A. T., Benjakul, S., Sinthusamran, S., Sookchoo, P., & Kishimura, H. (2019b). Protein hydrolysate from salmon frames: Production, characteristics and antioxidative activity. *Journal of Food Biochemistry*, 43, e12734. <https://doi.org/10.1111/jfbc.12734>
- Intarasirisawat, R., Benjakul, S., Wu, J., & Visessanguan, W. (2013). Isolation of antioxidative and ACE inhibitory peptides from protein hydrolysate of skipjack (*Katsuwana pelamis*) roe. *Journal of Functional Foods*, 5, 1854-1862. <https://doi.org/10.1016/j.jff.2013.09.006>
- Jai Ganesh, R., Nazeer, R., & Sampath Kumar, N. (2011). Purification and identification of antioxidant peptide from black pomfret, *Parastromateus niger* (Bloch, 1975) viscera protein hydrolysate. *Food Science and Biotechnology*, 20, 1087-1094. <https://doi.org/10.1007/s10068-011-0147-x>
- Je, J.-Y., Qian, Z.-J., Byun, H.-G., & Kim, S.-K. (2007). Purification and characterization of an antioxidant peptide obtained from tuna backbone protein by enzymatic hydrolysis. *Process Biochemistry*, 42, 840-846. <https://doi.org/10.1016/j.procbio.2007.02.006>
- Jung, W.-K., & Kim, S.-K. (2007). Calcium-binding peptide derived from pepsinolytic hydrolysates of hoki (*Johnius belangerii*) frame. *European Food Research and Technology*, 224, 763-767. <https://doi.org/10.1007/s00217-006-0371-4>
- Jung, W.-K., Lee, B.-J., & Kim, S.-K. (2006a). Fish-bone peptide increases calcium solubility and bioavailability in ovariectomised rats. *British Journal of Nutrition*, 95, 124-128. <https://doi.org/10.1079/BJN20051615>
- Jung, W.-K., Mendis, E., Je, J.-Y., Park, P.-J., Son, B. W., Kim, H. C., Choi, Y. K., & Kim, S.-K. (2006b). Angiotensin I-converting enzyme inhibitory peptide from yellowfin sole (*Limanda aspera*) frame protein and its antihypertensive effect in spontaneously hypertensive rats. *Food Chemistry*, 94, 26-32. <https://doi.org/10.1016/j.foodchem.2004.09.048>
- Kandyliari, A., Golla, J. P., Chen, Y., Papandroulakis, N., Kapsokefalou, M., & Vasiliou, V. (2020). Antiproliferative activity of protein hydrolysates derived from fish by-products on human colon and breast cancer cells. *Proceedings of the Nutrition Society*, 79, E282. <https://doi.org/10.1017/S002966512000230X>
- Karami, Z., & Akbari-Adergani, B. (2019). Bioactive food derived peptides: A review on correlation between structure of bioactive peptides and their functional properties. *Journal of Food Science and Technology*, 56, 535-547. <https://doi.org/10.1007/s13197-018-3549-4>
- Karnjanapratum, S., & Benjakul, S. (2017). Antioxidative and sensory properties of instant coffee fortified with galactose-fish skin gelatin hydrolysate maillard reaction products. *Carpathian Journal of Food Science & Technology*, 9, 90-99.
- Karnjanapratum, S., & Benjakul, S. (2018). Coconut oil based cookies fortified with bio-calcium: Characteristics and nutritional compositions. *Carpathian Journal of Food Science & Technology*, 10, 79-89.
- Ketnawa, S., Benjakul, S., Martínez-Alvarez, O., & Rawdkuen, S. (2017). Fish skin gelatin hydrolysates produced by visceral peptidase and bovine trypsin: Bioactivity and stability. *Food Chemistry*, 215, 383-390. <https://doi.org/10.1016/j.foodchem.2016.07.145>
- Khan, M. A. A., Hossain, M. A., Hara, K., Osatomi, K., Ishihara, T., & Nozaki, Y. (2003). Effect of enzymatic fish-scrap protein hydrolysate on gel-forming ability and denaturation of lizard fish *Saurida wanieso* surimi during frozen storage. *Fisheries Science*, 69, 1271-1280. <https://doi.org/10.1111/j.0919-9268.2003.00755.x>
- Kim, S.-K., & Jung, W.-K. (2012). Chapter 19 - Beneficial effect of teleost fish bone peptide as calcium supplements for bone mineralization. In S.-K. Kim (Ed.), *Advances in Food and Nutrition Research* (pp. 287-295). Academic Press. <https://doi.org/https://doi.org/10.1016/B978-0-12-416003-3.00019-6>
- Kim, S.-K., & Mendis, E. (2006a). Bioactive compounds from marine processing byproducts—a review. *Food Research International*, 39, 383-393. <https://doi.org/10.1016/j.foodres.2005.10.010>
- Kim, S.-K., & Mendis, E. (2006b). Bioactive compounds from marine processing byproducts – A review. *Food Research International*, 39, 383-393. <https://doi.org/https://doi.org/10.1016/j.foodres.2005.10.010>
- Kim, S.-K., & Wijesekara, I. (2010). Development and biological activities of marine-derived bioactive peptides: A review. *Journal of Functional Foods*, 2, 1-9. <https://doi.org/10.1016/j.jff.2010.01.003>
- Kitts, D. D., & Weiler, K. (2003). Bioactive proteins and peptides from food sources. Applications of bioprocesses used in isolation and recovery. *Current Pharmaceutical Design*, 9, 1309-1323. <https://doi.org/10.2174/1381612033454883>
- Klompong, V., Benjakul, S., Kantachote, D., & Shahidi, F. (2007). Antioxidative activity and functional properties of protein hydrolysate of yellow stripe trevally (*Selaroides leptolepis*) as influenced by the degree of hydrolysis and enzyme type. *Food Chemistry*, 102, 1317-1327. <https://doi.org/10.1016/j.foodchem.2006.07.016>
- Kristinsson, H. G., & Rasco, B. A. (2000). Fish protein hydrolysates: Production, biochemical, and functional properties. *Critical Reviews in Food Science and Nutrition*, 40, 43-81. <https://doi.org/10.1080/10408690091189266>
- Kudo, K., Onodera, S., Takeda, Y., Benkeblia, N., & Shiomi, N. (2009). Antioxidative activities of some peptides isolated from hydrolyzed potato protein extract. *Journal of Functional Foods*, 1, 170-176. <https://doi.org/10.1016/j.jff.2009.01.006>
- Kumar, V., Muzaddadi, A. U., Mann, S., Balakrishnan, R., Bembem, K., & Kalnar, Y. (2018). Utilization of fish processing waste: A waste to wealth approach. *Emerging Post-harvest Engineering and Technological Interventions for Enhancing Farmer's Income*, 5, 127-131.
- Larsen, T., Thilsted, S. H., Kongsbak, K., & Hansen, M. (2000). Whole small fish as a rich calcium source. *British Journal of Nutrition*, 83, 191-196. <https://doi.org/10.1017/S0007114500000246>
- Lee, J. K., Li-Chan, E. C., Cheung, I. W., Jeon, Y.-J., Ko, J.-Y., & Byun, H.-G. (2019). Neuroprotective effect of  $\beta$ -secretase inhibitory peptide from Pacific hake (*Merluccius productus*) fish protein hydrolysate. *Current Alzheimer Research*, 16, 1028-1038. <https://doi.org/10.2174/1567205016666191113122046>
- Lee, S.-H., Qian, Z.-J., & Kim, S.-K. (2010). A novel angiotensin I converting enzyme inhibitory peptide from tuna frame protein hydrolysate and its antihypertensive effect in spontaneously hypertensive rats. *Food Chemistry*, 118, 96-102. <https://doi.org/10.1016/j.foodchem.2009.04.086>
- Logesh, A. R., Pravinkumar, M., Raffi, S. M., & Kalaiselvam, M. (2012). Calcium and phosphorus determination in bones of low value fishes, *Sardinella longiceps* (Valenciennes)



- and *Trichiurus savala* (Cuvier), from Parangipettai, Southeast Coast of India. *Asian Pacific Journal of Tropical Disease*, 2, S254-S256.  
[https://doi.org/https://doi.org/10.1016/S2222-1808\(12\)60160-1](https://doi.org/https://doi.org/10.1016/S2222-1808(12)60160-1)
- Malde, M. K., Bügel, S., Kristensen, M., Malde, K., Graff, I. E., & Pedersen, J. I. (2010). Calcium from salmon and cod bone is well absorbed in young healthy men: A double-blinded randomised crossover design. *Nutrition & Metabolism*, 7, 61. <https://doi.org/10.1186/1743-7075-7-61>
- Martínez-Montaño, E., Sarmiento-Machado, R. M., Osuna-Ruiz, I., Benítez-García, I., Pacheco-Aguilar, R., Navarro-Peraza, R. S., Sánchez, M. E. L., Ortiz, A. V., Báez, L. J. G., & Bañuelos-Vargas, I. (2021). Effect of degree of hydrolysis on biochemical properties and biological activities (antioxidant and antihypertensive) of protein hydrolysates from pacific thread herring (*Ophistonema libertate*) stickwater. *Waste and Biomass Valorization*, 13, 1015–1027. <https://doi.org/10.1007/s12649-021-01590-z>
- Mazorra-Manzano, M. A., Pacheco-Aguilar, R., Ramírez-Suárez, J. C., García-Sánchez, G., & Lugo-Sánchez, M. E. (2012). Endogenous proteases in Pacific whiting (*Merluccius productus*) muscle as a processing aid in functional fish protein hydrolysate production. *Food and Bioprocess Technology*, 5, 130-137.  
<https://doi.org/10.1007/s11947-010-0374-9>
- Mbatia, B., Ogonda, L., Muge, E., & Mulaa, F. (2014). Antioxidative and functional properties of *Rastrineobola argentea* (dagaa) fish protein hydrolysate. *Discourse Journal of Agriculture and Food Sciences*, 2, 180-189.
- Misir, G. B. (2022). Novel utilization of fish by-products and wastes: Protein hydrolysates. *Acta Aquatica Turcica*, 18(2), 283-294.  
<https://doi.org/10.22392/actaquatr.1031442>
- Misir, G. B. & Kosal, S. (2019). Effects of ultrasound treatment on structural, chemical and functional properties of protein hydrolysate of rainbow trout (*oncorhynchus mykiss*) by-products. *Italian Journal of Food Science*, 31, 205-223. <https://doi.org/10.14674/IJFS-1218>
- Moskowitz, R. W. (2000). Role of collagen hydrolysate in bone and joint disease. *Seminars in arthritis and rheumatism*, 30, 87-99. <https://doi.org/10.1053/sarh.2000.9622>
- Najafian, L., & Babji, A. (2012). A review of fish-derived antioxidant and antimicrobial peptides: Their production, assessment, and applications. *Peptides*, 33, 178-185.  
<https://doi.org/10.1016/j.peptides.2011.11.013>
- Nalinanon, S., Benjakul, S., Kishimura, H., & Shahidi, F. (2011). Functionalities and antioxidant properties of protein hydrolysates from the muscle of ornate threadfin bream treated with pepsin from skipjack tuna. *Food Chemistry*, 124, 1354-1362.  
<https://doi.org/10.1016/j.foodchem.2010.07.089>
- Nasri, R., Amor, I. B., Bougatef, A., Nedjar-Arroume, N., Dhulster, P., Gargouri, J., Châabouni, M. K., & Nasri, M. (2012). Anticoagulant activities of goby muscle protein hydrolysates. *Food Chemistry*, 133, 835-841.  
<https://doi.org/10.1016/j.foodchem.2012.01.101>
- Nemati, M., Kamilah, H., Huda, N., & Ariffin, F. (2016). *In vitro* calcium availability in bakery products fortified with tuna bone powder as a natural calcium source. *International Journal of Food Sciences and Nutrition*, 67, 535-540.  
<https://doi.org/10.1080/09637486.2016.1179269>
- Ngo, D.-H., Qian, Z.-J., Ryu, B., Park, J. W., & Kim, S.-K. (2010). In vitro antioxidant activity of a peptide isolated from Nile tilapia (*Oreochromis niloticus*) scale gelatin in free radical-mediated oxidative systems. *Journal of Functional Foods*, 2, 107-117.  
<https://doi.org/10.1016/j.jff.2010.02.001>
- Nilsuwan, K., Pomtong, S., Chedosama, A., Sookchoo, P., & Benjakul, S. (2023). Chemical compositions and characteristics of bioactive calcium from Asian sea bass (*Lates calcarifer*) scales as influenced by pretreatment and heating processes. *Foods*, 12, 2695.  
<https://doi.org/10.3390/foods12142695>
- Nirmal, N. P., Santivarangkna, C., Rajput, M. S., Benjakul, S., & Maqsood, S. (2022). Valorization of fish byproducts: Sources to end-product applications of bioactive protein hydrolysate. *Comprehensive Reviews in Food Science and Food Safety*, 21, 1803-1842.  
<https://doi.org/10.1111/1541-4337.12917>
- Parolini, C., Vik, R., Busnelli, M., Bjørndal, B., Holm, S., Brattelid, T., Manzini, S., Ganzetti, G. S., Dellera, F., & Halvorsen, B. (2014). A salmon protein hydrolysate exerts lipid-independent anti-atherosclerotic activity in ApoE-deficient mice. *PLoS One*, 9, e97598.  
<https://doi.org/10.1371/journal.pone.0097598>
- Patil, U., & Benjakul, S. (2019). Comparative study on extraction of virgin coconut oil with the aid of partially purified protease from seabass pyloric caeca and commercial trypsin. *Journal of Food Biochemistry*, 43, e13024. <https://doi.org/10.1111/jfbc.13024>
- Patil, U., Nagarajarao, R. C., Balange, A. K., Zhang, B., & Benjakul, S. (2022a). Proteolytic activity and characteristics of skipjack tuna trypsin loaded alginate-chitosan beads as affected by drying methods and trehalose/glycerol. *International Journal of Food Science & Technology*, 55, 2695-2705.  
<https://doi.org/10.1111/ijfs.16039>
- Patil, U., Nikoo, M., Zhang, B., & Benjakul, S. (2022b). Freeze-dried tuna pepsin powder stabilized by some cryoprotectants: *In vitro* simulated gastric digestion toward different proteins and its storage stability. *Foods*, 11, 2292. <https://doi.org/10.3390/foods11152292>
- Patil, U., Saetang, J., Zhang, B., & Benjakul, S. (2022c). Use of tuna visceral pepsin in combination with trypsin as digestion aid: Enhanced protein hydrolysis and bioavailability. *Foods*, 12, 125.  
<https://doi.org/10.3390/foods12010125>
- Patil, U., Gulzar, S., Ma, L., Zhang, B., & Benjakul, S. (2023). Pickering emulsion stabilized by fish myofibrillar proteins modified with tannic acid, as influenced by different drying methods. *Foods*, 12, 1556.  
<https://doi.org/10.3390/foods12071556>
- Patil, U., Baloch, K. A., Nile, S. H., Kim, J. T., & Benjakul, S. (2023). Trypsin from Pyloric Caeca of Asian Seabass: Purification, Characterization, and Its Use in the Hydrolysis of Acid-Soluble Collagen. *Foods*, 12(15), 2937. <https://doi.org/10.3390/foods12152937>
- Pérez-Escalante, E., Padilla-Zúñiga, S. A., Contreras-López, E., Sebastián-Nicolás, J. L., Pérez-Flores, J. G., Olloqui, E. J., & González-Olivares, L. G. (2022). Antioxidant and Antihypertensive Properties from Muscle Hydrolysates of Farm Rainbow Trout. *Biology and Life Sciences Forum*, 18, 55. <https://doi.org/10.3390/Foods2022-12991>
- Phanturat, P., Benjakul, S., Visessanguan, W., & Roytrakul, S. (2010). Use of pyloric caeca extract from bigeye snapper (*Priacanthus macracanthus*) for the production of gelatin hydrolysate with antioxidative activity. *LWT-Food Science and Technology*, 43, 86-97.  
<https://doi.org/10.1016/j.lwt.2009.06.010>
- Piccirillo, C., Silva, M. F., Pullar, R. C., Braga da Cruz, I., Jorge, R., Pintado, M. M. E., & Castro, P. M. L. (2013). Extraction

- and characterisation of apatite- and tricalcium phosphate-based materials from cod fish bones. *Materials Science and Engineering: C*, 33, 103-110. <https://doi.org/10.1016/j.msec.2012.08.014>
- Picot, L., Bordenave, S., Didelot, S., Fruitier-Arnaudin, I., Sannier, F., Thorkelsson, G., Bergé, J., Guérard, F., Chabeaud, A., & Piot, J. (2006). Antiproliferative activity of fish protein hydrolysates on human breast cancer cell lines. *Process Biochemistry*, 41, 1217-1222. <https://doi.org/10.1016/j.procbio.2005.11.024>
- Pires, C., & Batista, I. (2013). Functional properties of fish protein hydrolysates. In R. P. Gálvez & J. P. Bergé (Eds), *Utilization of fish waste* (pp. 59-75). CRC Press.
- Pudtikajorn, K., Sae-leaw, T., Buamard, N., Zhou, A., Ma, L., & Benjakul, S. (2022). Characterisation of fish tofu fortified with skipjack tuna (*Katsuwonus pelamis*) eyeball scleral cartilage biocalcium. *International Journal of Food Science & Technology*, 57, 6711-6721. <https://doi.org/10.1111/ijfs.16022>
- Pudtikajorn, K., Sae-leaw, T., Yesilsu, A. F., Sookchoo, P., & Benjakul, S. (2023). Process development and characteristics of biocalcium from skipjack tuna (*Katsuwonus pelamis*) eyeball scleral cartilage. *Waste and Biomass Valorization*. <https://doi.org/10.1007/s12649-023-02075-x>
- Rabiei, S., Rezaei, M., Nikoo, M., Khezri, M., Rafieian-Kopai, M., & Anjomshoaa, M. (2022). Antioxidant properties of Klunzinger's mullet (*Liza klunzingeri*) protein hydrolysates prepared with enzymatic hydrolysis using a commercial protease and microbial hydrolysis with *Bacillus licheniformis*. *Food Science and Technology International*, 28(3), 233-246. <https://doi.org/10.1177/10820132211005297>
- Raksakulthai, R., & Haard, N. F. (2003). Exopeptidases and their application to reduce bitterness in food: A review. *Critical Reviews in Food Science and Nutrition*, 43, 401-445. <https://doi.org/10.1080/10408690390826572>
- Rana, S., Singh, A., Surasani, V. K. R., Kapoor, S., Desai, A., & Kumar, S. (2023). Fish processing waste: a novel source of non-conventional functional proteins. *International Journal of Food Science and Technology*, 58(5), 2637-2644. <https://doi.org/10.1111/ijfs.16104>
- Ren, J., Zhao, M., Shi, J., Wang, J., Jiang, Y., Cui, C., Kakuda, Y., & Xue, S. J. (2008). Purification and identification of antioxidant peptides from grass carp muscle hydrolysates by consecutive chromatography and electrospray ionization-mass spectrometry. *Food Chemistry*, 108, 727-736. <https://doi.org/10.1016/j.foodchem.2007.11.010>
- Rivero-Pino, F., Espejo-Carpio, F. J., & Guadix, E. M. (2020). Antidiabetic food-derived peptides for functional feeding: Production, functionality and *in vivo* evidences. *Foods*, 9, 983. <https://doi.org/10.3390/foods9080983>
- Rodrigues, D. P., Calado, R., Ameixa, O. M., Valcarcel, J., & Vázquez, J. A. (2021). Valorisation of Atlantic codfish (*Gadus morhua*) frames from the cure-salting industry as fish protein hydrolysates with *in vitro* bioactive properties. *LWT- Food Science and Technology*, 149, 111840. <https://doi.org/10.1016/j.lwt.2021.111840>
- Saito, M., Kiyose, C., Higuchi, T., Uchida, N., & Suzuki, H. (2009). Effect of collagen hydrolysates from salmon and trout skins on the lipid profile in rats. *Journal of Agricultural and Food Chemistry*, 57, 10477-10482. <https://doi.org/10.1021/jf902355m>
- Samaranayaka, A. G., Kitts, D. D., & Li-Chan, E. C. (2010). Antioxidative and angiotensin-I-converting enzyme inhibitory potential of a Pacific hake (*Merluccius productus*) fish protein hydrolysate subjected to simulated gastrointestinal digestion and Caco-2 cell permeation. *Journal of Agricultural and Food Chemistry*, 58, 1535-1542. <https://doi.org/10.1021/jf9033199>
- Sasidharan, A., & Venugopal, V. (2020). Proteins and co-products from seafood processing discards: Their recovery, functional properties and applications. *Waste and Biomass Valorization*, 11, 5647-5663. <https://doi.org/10.1007/s12649-019-00812-9>
- Shahosseini, S. R., Javadian, S. R., & Safari, R. (2022). Effects of molecular weights-assisted enzymatic hydrolysis on antioxidant and anticancer activities of *liza abu* muscle protein hydrolysates. *International Journal of Peptide Research and Therapeutics*, 28, 72. <https://doi.org/10.1007/s10989-022-10371-8>
- Sharma, K., Nilsuwan, K., Ma, L., & Benjakul, S. (2023a). Effect of liposomal encapsulation and ultrasonication on debittering of protein hydrolysate and plastein from salmon frame. *Foods*, 12, 761. <https://doi.org/10.3390/foods12040761>
- Sharma, K., Nilsuwan, K., Zhang, B., Hong, H., & Benjakul, S. (2023b). Protein hydrolysate from salmon frame debittered by plastein reaction: amino acid composition, characteristics and antioxidant activities. *International Journal of Food Science & Technology*, 58, 154-166. <https://doi.org/10.1111/ijfs.16183>
- Shaviklo, G. R., Thorkelsson, G., Sveinsdottir, K., & Rafipour, F. (2011). Chemical properties and sensory quality of ice cream fortified with fish protein. *Journal of the Science of Food and Agriculture*, 91, 1199-1204. <https://doi.org/10.1002/jsfa.4299>
- Sinaga, M., Zuhra, C. F., Lenny, S., Hardiyanti, R., Zaidar, E., & Haditia, M. (2023). Antibacterial and antioxidant activity of fish protein hydrolysate from mahseer (*Tor tambroides*). *AIP Conference Proceedings*, 2626, 030016. <https://doi.org/10.1063/5.0136087>
- Singh, A., Idowu, A. T., Benjakul, S., Kishimura, H., Aluko, R. E., & Kumagai, Y. (2020). Debittering of salmon (*Salmo salar*) frame protein hydrolysate using 2-butanol in combination with  $\beta$ -cyclodextrin: Impact on some physicochemical characteristics and antioxidant activities. *Food Chemistry*, 321, 126686. <https://doi.org/10.1016/j.foodchem.2020.126686>
- Sinthusamran, S., Benjakul, S., Kijroongrojana, K., Prodpran, T., & Kishimura, H. (2020a). Protein hydrolysates from Pacific white shrimp cephalothorax manufactured with different processes: Compositions, characteristics and antioxidative activity. *Waste and Biomass Valorization*, 11, 1657-1670. <https://doi.org/10.1007/s12649-018-0517-1>
- Sinthusamran, S., Idowu, A. T., Benjakul, S., Prodpran, T., Yesilsu, A. F., & Kishimura, H. (2020b). Effect of proteases and alcohols used for debittering on characteristics and antioxidative activity of protein hydrolysate from salmon frames. *Journal of Food Science and Technology*, 57, 473-483. <https://doi.org/10.1007/s13197-019-04075-z>
- Song, R., Wei, R.-b., Luo, H.-y., & Yang, Z.-s. (2014). Isolation and identification of an antiproliferative peptide derived from heated products of peptic hydrolysates of half-fin anchovy (*Setipinna taty*). *Journal of Functional Foods*, 10, 104-111. <https://doi.org/10.1016/j.jff.2014.06.010>
- Taroncher, M., Rodríguez-Carrasco, Y., Barba, F. J., & Ruiz, M.-J. (2023). Enhancement of the antioxidant effect of natural products on the proliferation of Caco-2 cells produced by fish protein hydrolysates and collagen. *International Journal of Molecular Sciences*, 24, 6871. <https://doi.org/10.3390/ijms24076871>

- Thiansilakul, Y., Benjakul, S., & Shahidi, F. (2007). Antioxidative activity of protein hydrolysate from round scad muscle using alcalase and flavourzyme. *Journal of Food Biochemistry*, 31, 266-287. <https://doi.org/10.1111/j.1745-4514.2007.00111.x>
- Tunick, M. H. (1987). Calcium in dairy products. *Journal of Dairy Science*, 70, 2429-2438. [https://doi.org/https://doi.org/10.3168/jds.S0022-0302\(87\)80305-3](https://doi.org/https://doi.org/10.3168/jds.S0022-0302(87)80305-3)
- Villamil, O., Váquiro, H., & Solanilla, J. F. (2017). Fish viscera protein hydrolysates: Production, potential applications and functional and bioactive properties. *Food Chemistry*, 224, 160-171. <https://doi.org/10.1016/j.foodchem.2016.12.057>
- Vo, T. D., Duong, K. L., Nguyen, A. N., Vo, B. C., Tran, M. C., Tran, H. G., ... & Nguyen, A. T. (2023). Featherback (*Chitala ornata*) skin protein hydrolysate: antioxidant activity, amino acid composition, foaming and emulsifying properties. *Chemical Engineering Transactions*, 106, 877-882.
- Wald, M., Schwarz, K., Rehbein, H., Bußmann, B., & Beermann, C. (2016). Detection of antibacterial activity of an enzymatic hydrolysate generated by processing rainbow trout by-products with trout pepsin. *Food Chemistry*, 205, 221-228. <https://doi.org/10.1016/j.foodchem.2016.03.002>
- Wan, P., Cai, B., Chen, H., Chen, D., Zhao, X., Yuan, H., Huang, J., Chen, X., Luo, L., & Pan, J. (2023). Antidiabetic effects of protein hydrolysates from *Trachinotus ovatus* and identification and screening of peptides with  $\alpha$ -amylase and DPP-IV inhibitory activities. *Current Research in Food Science*, 6, 100446. <https://doi.org/10.1016/j.crfs.2023.100446>
- Wang, J., Wang, Y. M., Li, L. Y., Chi, C. F., & Wang, B. (2022). Twelve antioxidant peptides from protein hydrolysate of skipjack tuna (*Katsuwonus pelamis*) roe prepared by flavourzyme: Purification, sequence identification, and activity evaluation. *Frontiers in Nutrition*, 8, 813780. <https://doi.org/10.3389/fnut.2021.813780>
- Wang, Y., Zhu, F., Han, F., & Wang, H. (2008). Purification and characterization of antioxidative peptides from salmon protamine hydrolysate. *Journal of Food Biochemistry*, 32, 654-671. <https://doi.org/10.1111/j.1745-4514.2008.00190.x>
- Wijayanti, I., Benjakul, S., Chantakun, K., Prodpran, T., & Sookchoo, P. (2022a). Effect of Asian sea bass bio-calcium on textural, rheological, sensorial properties and nutritive value of Indian mackerel fish spread at different levels of potato starch. *International Journal of Food Science & Technology*, 57, 3181-3195. <https://doi.org/https://doi.org/10.1111/ijfs.15651>
- Wijayanti, I., Benjakul, S., Saetang, J., Prodpran, T., & Sookchoo, P. (2022b). Soluble Asian sea bass bone bio-calcium: characteristics, bioavailability across Caco-2 cells and fortification into apple juice. *International Journal of Food Science & Technology*, 57, 5859-5868. <https://doi.org/https://doi.org/10.1111/ijfs.15904>
- Wijayanti, I., Benjakul, S., & Sookchoo, P. (2021a). Effect of high pressure heating on physical and chemical characteristics of Asian sea bass (*Lates calcarifer*) backbone. *Journal of Food Science and Technology*, 58, 3120-3129. <https://doi.org/10.1007/s13197-020-04815-6>
- Wijayanti, I., Benjakul, S., & Sookchoo, P. (2021b). Preheat-treatment and bleaching agents affect characteristics of bio-calcium from Asian sea bass (*Lates calcarifer*) backbone. *Waste and Biomass Valorization*, 12, 3371-3382. <https://doi.org/10.1007/s12649-020-01224-w>
- Wijayanti, I., Prodpran, T., Sookchoo, P., Nirmal, N., Zhang, B., Balange, A., & Benjakul, S. (2023). Textural, rheological and sensorial properties of mayonnaise fortified with Asian sea bass bio-calcium. *Journal of the American Oil Chemists' Society*, 100, 123-140. <https://doi.org/https://doi.org/10.1002/aocs.12649>
- Wijayanti, I., Singh, A., Benjakul, S., & Sookchoo, P. (2021c). Textural, sensory, and chemical characteristic of threadfin bream (*Nemipterus* sp.) surimi gel fortified with bio-calcium from bone of Asian sea bass (*Lates calcarifer*). *Foods*, 10, 976. <https://doi.org/10.3390/foods10050976>
- Wijayanti, I., Singh, A., Prodpran, T., Sookchoo, P., & Benjakul, S. (2021d). Effect of Asian sea bass (*Lates calcarifer*) bio-calcium in combination with different calcium salts on gel properties of threadfin bream surimi. *Journal of Aquatic Food Product Technology*, 30, 1173-1188. <https://doi.org/10.1080/10498850.2021.1975004>
- Wu, H.-C., Chen, H.-M., & Shiau, C.-Y. (2003). Free amino acids and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*). *Food Research International*, 36, 949-957. [https://doi.org/10.1016/S0963-9969\(03\)00104-2](https://doi.org/10.1016/S0963-9969(03)00104-2)
- Wu, J., Fujioka, M., Sugimoto, K., Mu, G., & Ishimi, Y. (2004). Assessment of effectiveness of oral administration of collagen peptide on bone metabolism in growing and mature rats. *Journal of bone and mineral metabolism*, 22, 547-553. <https://doi.org/10.1007/s00774-004-0522-2>
- Yeaman, M. R., & Yount, N. Y. (2003). Mechanisms of antimicrobial peptide action and resistance. *Pharmacological reviews*, 55, 27-55. <https://doi.org/10.1124/pr.55.1.2>
- Xie, H., Ouyang, K., Shi, W., Wang, W., Wang, Y., Xiong, H., & Zhao, Q. (2023). Enhancing the interfacial stability of O/W emulsion by adjusting interactions of chitosan and rice protein hydrolysate. *Food Hydrocolloids*, 137, 108406. <https://doi.org/10.1016/j.foodhyd.2022.108406>
- Yin, S.-W., Tang, C.-H., Cao, J.-S., Hu, E.-K., Wen, Q.-B., & Yang, X.-Q. (2008). Effects of limited enzymatic hydrolysis with trypsin on the functional properties of hemp (*Cannabis sativa* L.) protein isolate. *Food Chemistry*, 106, 1004-1013. <https://doi.org/10.1016/j.foodchem.2007.07.030>
- You, L., Zhao, M., Regenstein, J. M., & Ren, J. (2010). Changes in the antioxidant activity of loach (*Misgurnus anguillicaudatus*) protein hydrolysates during a simulated gastrointestinal digestion. *Food Chemistry*, 120, 810-816. <https://doi.org/10.1016/j.foodchem.2009.11.018>
- Zamora-Sillero, J., Gharsallaoui, A., & Prentice, C. (2018). Peptides from fish by-product protein hydrolysates and its functional properties: An overview. *Marine biotechnology*, 20, 118-130. <https://doi.org/10.1007/s10026-018-9799-3>
- Zhou, Y., Zhang, Y., Hong, H., Luo, Y., Li, B., & Tan, Y. (2023). Mastering the art of taming: Reducing bitterness in fish by-products derived peptides. *Food Research International*, 173, 113241. <https://doi.org/10.1016/j.foodres.2023.113241>