



Effects of Microplastics on the Activity of Digestive-, Antioxidative- and Immune-related Enzymes in Soiny Mullet (*Liza haematocheila* Temminck & Schlegel, 1845) Larvae

Yingying Zhang¹, Shengyuan Zhang^{1,2}, Ke Sun¹, Qihuan Zhang¹, Eakapol Wangkahart³, Zisheng Wang¹, Yanming Sui^{1*} , Zhitao Qi¹ 

¹Jiangsu Key Laboratory of Biochemistry and Biotechnology of Marine Wetland, Yancheng Institute of Technology, Yancheng, Jiangsu Province, China

²College of Biotechnology, Jiangsu University of Science and Technology, Zhenjiang, China

³Laboratory of Fish Immunology and Nutrigenomics, Applied Animal and Aquatic Sciences Research Unit, Division of Fisheries, Faculty of Technology, Mahasarakham University, Maha Sarakham, Thailand

How to Cite

Zhang, Y., Zhang, S., Sun, K., Zhang, Q., Wangkahart, E., Wang, Z., Sui, Y., Qi, Z. (2024). Effects of Microplastics on the Activity of Digestive-, Antioxidative- and Immune-related Enzymes in Soiny Mullet (*Liza haematocheila* Temminck & Schlegel, 1845) Larvae. *Turkish Journal of Fisheries and Aquatic Sciences*, 24(6), TRJFAS24718. <https://doi.org/10.4194/TRJFAS24718>

Article History

Received 09 November 2023

Accepted 07 March 2023

First Online 26 April 2024

Corresponding Author

E-mail: suiyanming@foxmail.com

Keywords

Microplastic

Liza haematocheila

Digestive enzymes

Antioxidant enzymes

Immune enzymes

Abstract

Microplastics (MPs), novel pollutants in the water environment, can cause damage to fish, especially to fish larvae. However, the effects of MPs on fish in estuarine areas with high levels of MPs pollution are unknown. In this study, a 14-day exposure test was conducted to analyze the effects of MPs on soiny mullet larvae by measuring the activities of digestive enzymes, antioxidant enzymes, and immune enzymes in soiny mullet larvae after exposure to MPs of different particle sizes (0.5 μm , 0.1mg/L; 6 μm , 0.1mg/L). Results showed that MPs exposure significantly decreased the activities of pepsin (PPS), alpha-amylase (α -AMS), catalase (CAT), and superoxide dismutase (SOD), and significantly increased the activities of acid phosphatase (ACP) and alkaline phosphatase (ALP), among which the PPS, CAT, SOD, ACP, and ALP were more sensitive to exposure to small particle size MPs (0.5 μm). Overall, exposure to MPs reduced digestion, destroyed the antioxidant system, and triggered an immune response in soiny mullet larvae, especially the small particles of MPs (0.5 μm). The results provide basis for understanding the detriment of microplastic in fish.

Introduction

Plastic products are widely used in all aspects of daily life due to their cheapness and durability (Sui et al., 2020). From the 1950s to 2018, global plastic production has risen from 1.5 million tons to 359 million tons (Dobaradaran et al., 2018). However, the recycling rate of all plastics is only 1% -5%, and the final discharge into the ocean accounts for 10% of the total plastic production (Güven et al., 2017; Mattsson et al., 2017; Tata et al., 2020). Plastics are decomposed into microplastics (MPs) with a diameter of less than 5 mm

under long-term biological, physical and chemical effects (Thompson et al., 2004; Andrady, 2011). At present, MPs pollution in the ocean and rivers has attracted worldwide attention.

Estuaries are generally considered to be the main passageway for MPs to enter the ocean from freshwater environments and it is also a hot spot for investigating MPs pollution in recent years (Zhang et al., 2019). Research in China shows that the average concentration of MPs in Haihe Estuary and Yongdingxinhe Estuary surface water samples were respectively 1485.7 ± 819.9 items/ m^3 and 788.0 ± 464.2 items/ m^3 (Wu et al., 2019).

The abundance of MPs in Jiaojiang, Oujiang, and Minjiang estuaries in Zhejiang ranged from 100 items/m³ to 4100 items/m³ (Zhao et al., 2015). Yangtze Estuary MPs abundance was 4137.3±2461.5 items/m³ (Zhao et al., 2014). The MPs pollution is much more serious at China's Pearl River Estuary, reaching 8902 items/m³ (Yan et al., 2019). Now, the MPs pollution in the estuarine areas is becoming an environmental problem all over the world. However, current studies mainly focus on the pollution degree of MPs in the estuarine areas, limited research studies the effects of MPs on the large of animals in the estuarine areas (e.g. physical or immune status).

The choice of polystyrene for this study stems from its extensive use across various products, making it a significant contributor to the substantial amount of plastic entering our oceans. Its widespread prevalence underscores importance in the context of environmental pollution (Hanachi et al., 2021). MPs can be ingested by a variety of marine animals, including crustaceans (Murray & Cowie, 2011; Devriese et al., 2015), mollusks (Van Cauwenberghe and Janssen, 2014), marine mammals (Fossi et al., 2012; Bravo Rebolledo et al., 2013), zooplankton species (Desforges et al., 2015), marine turtles (Tourinho et al., 2010), sea birds (Bravo Rebolledo et al., 2013), and fish (Pedà et al., 2016; Rummel et al., 2016; Steer et al., 2017). Polystyrene may be excreted in the feces and enter the bloodstream after absorption through the intestinal epithelium and distributed to other tissues in the body. The particles may penetrate cell membranes and enter the fish. Once MPs are absorbed into the digestive system, they can penetrate the anaerobic tissues of the intestine and accumulate in the fatty tissues of the animal (Banaei et al., 2022; Kim et al., 2021). Small size MPs can enter the circulatory system and accumulate in various tissues under certain conditions (Browne et al., 2008). Studies indicate that environmental stress and ingestion of toxic substances can cause oxidative stress and affect fish digestion and immune functions, thereby inducing changes in the activity of these enzymes (Couillard et al., 2009; Blewett & Leonard, 2017; Zhao et al., 2022). The resulting biological consequences can affect the growth and development, reproduction ability, and survival rate of marine animals (Mazurais et al., 2015; Nobre et al., 2015; Sussarellu et al., 2016; Beiras et al., 2018). During the fish larvae stage, organ development and related physiological functions are influenced by many external factors, which determine the ability to cope with the challenges of survival in the following physiological stages (Shahriari Moghadam et al., 2014). The healthy growth of fish larvae is critical to the sustainable development of fish stocks and the stability of ecosystems (Steer et al., 2017). Therefore, it is particularly important to explore the negative effects of MPs on larvae fish.

Soiny mullet (*Liza haematocheila* Temminck & Schlegel, 1845) is an important farmed fish in China (Qi et al., 2022). As migratory fish, soiny mullet larvae live in

the estuaries with dramatic environmental changes, such as MPs pollution, N₂O pollution, metal pollution, and ocean acidification (Soletchnik et al., 2007; Murray et al., 2015; Wijesiriet al., 2019). Compared with adult fish, fish larvae are more sensitive to the changes of the external environment and easier to be observed (Yang et al., 2020). Therefore, by taking fish larvae as the test object, we aimed to observe the impact of external stimuli on the body in detail. As filter-feeding fish, soiny mullet larvae can inadvertently eat toxic MPs. Furthermore, this study aims to explore the effect of exposure to polystyrene (PS) MPs with different particle sizes (6 µm and 0.5 µm) for 14 d on the soiny mullet larvae under laboratory conditions. We stimulated the extreme environments by using the abundance of 0.1 mg/L MPs in both treatment groups. The activity of digestive enzymes, antioxidant enzymes, and immune enzymes, including pepsin (PPS), alpha-amylase (α-AMS), catalase (CAT), superoxide dismutase (SOD), acid phosphatase (ACP), and alkaline phosphatase (ALP) were analyzed, providing the theoretical basis for further research on the effects of MPs on fish, especially larvae fish.

Materials and Methods

Soiny Mullet Preparation

A total of 1,000 healthy soiny mullet larvae (average body length 2.67±0.16 cm, average body weight 0.14±0.01g) were provided by a local fish farm in Sheyangtown, Yancheng city, Jiangsu province, China. In the laboratory, the fish were reared for at least one week in artificial seawater simulating pristine water conditions (temperature=25°C, salinity=15‰, pH=8.1, dissolved oxygen>7.0 mg/L) to acclimate. The fish were fed with brine shrimp (*Artemia salina* Linnaeus, 1758) twice daily at the rate of 2% of the fish's body weight. Excrement and uneaten food were removed by suction on a daily basis.

MPs Exposure

The monodisperse PS microspheres were purchased from BaseLine ChromTech Research Centre (Tianjin, China). Based on the size of the fish being studied and the size of the microplastics that are more common in the environment in previous studies, 0.5 µm (2.5 w/v, 10 ml, 0.5±0.012 µm, Figure 1) and 6 µm (2.5 w/v, 10 ml, 6±0.031 µm, Figure 2) MPs were chosen in this study and added to the seawater (Huang et al., 2020; Le et al., 2018). To simulate extreme MPs pollution in the water environment, the concentration of MPs in the experimental group was set to be 0.1 mg/L (Yan et al., 2020). During the experimental period, the aerators and air stones were used to maintain the suspension of MPs.

Three treated groups were set up in this study, which was named as blank control group (CTRL), small particle size MPs (0.5 µm) treated group (SMPs), and

large particle size MPs (6 μm) treated group (LMPs), respectively. Each treated group contained 3 replicated glass tanks (20 fish per tank). During the 14 d test period, the artificial seawater was completely replaced daily to to guarantee the constant concentration of microplastics.

Sample Collection

All the following experimental methods were under the National Institutes of Health Guide for the Care and Use of Laboratory Animals of China.

At the 3rd, 7th, and 14th day post exposure (dpe), 3 fish were randomly selected from each tank for sampling. Before sampling, the soiny mullet larvae were euthanized through anesthetic overdose with tricaine methane-sulfonate (MS-222) (Sigma, USA). Sterile scissors and tweezers to cut off the head and tail of the larvae, and the remains were transferred the remainder into a cryovial. In order to handle the samples in a uniform manner to ensure the reliability of the experiments and the accuracy of the results, the cryovials were quickly frozen in liquid nitrogen and transferred to -80°C for storage until the biochemical analyses began. The FTIR analysis was used to confirm that microplastic has been removed by the exposed larvae.

Enzymes Activities Assay

Each sample was homogenized in normal saline (0.9%) in a 1:10 weight: volume ratio. After high-speed centrifugation, the supernatant was absorbed for the measurement of different enzyme activities. Mixed them well, incubated at 37°C for 20 minutes, and read at 450 nm. The amount of enzyme corresponding to 50% SOD inhibition in this reaction system is one SOD activity unit (U) and the unit of enzyme is U/mgprot. The protein concentration of each sample was determined by Coomassie brilliant blue method. Afterward, the activities of PPS, α -AMS, CAT, SOD, ACP, and ALP were measured on a microplate reader, respectively. All enzymes activities kits in this study were purchased from Nanjing Jiancheng Bioengineering Institute (Nanjing, China). We strictly followed the operation steps and calculation formula of the kit's instructions to determine the activities of each enzyme.

Statistical Analyses

The data in the study were tested for homogeneity of variances and normal distribution first. The data significant differences were analyzed using one-way or multiple-way analysis of variance (ANOVA) followed by Tukey's post hoc test to evaluate the differences in the




Figure 1. Microscopic image of 0.5 μm MPs.




Figure 2. Microscopic image of 6 μm MPs.

influence of MPs size and MPs exposure time on test indexes. The data difference was considered significant and had statistical significance when $P < 0.05$. The statistical analysis of the data was carried out by the SPSS 24.0 statistical software (Chicago, USA). All results were expressed as mean \pm standard deviation (SD).

Results

Effects of MPs on the Activities of Digestive Enzymes of Soiny Mullet Larvae

The effect of MPs exposure on the PPS activities of soiny mullet was shown in Figure 3. The MPs exposure reduced the activity of PPS, especially in SMPs. However, the enzyme activities of the two treated

groups remained stable throughout the experimental period. The PPS activity of LMPs and SMPs were significantly lower than that of CTRL ($P < 0.05$). In addition, the PPS activity of SMPs was significantly lower than that of LMPs ($P < 0.05$) (Figure 3).

The effect of MPs exposure on the α -AMS activities of soiny mullet was shown in Figure 4. At 3 dpe, the MPs exposure had no significant effect on the α -AMS activity ($P > 0.05$). At 7 dpe, MPs exposure did not show no significant effect on the α -AMS activity of SMPs ($P > 0.05$), either, but significantly inhibited the the α -AMS activity in LMPs ($P < 0.05$). At 14 dpe, the MPs exposure significantly reduced the α -AMS activity in SMPs ($P < 0.05$), and also significantly inhibited the α -AMS activity in LMPs, which was lower than these in the CTRL and SMPs ($P < 0.05$) (Figure 4).




Figure 3. Effects of exposure to MPs with different particle sizes on PPS activity of soiny mullet (*Liza haematocheila*) larvae (n=9). Data were presented as mean \pm SD. Different letters indicate significant differences ($P < 0.05$). Conversely, there is no significant difference between the same letters.




Figure 4. Effects of exposure to MPs with different particle sizes on α -AMS activity of soiny mullet (*Liza haematocheila*) larvae (n=9). Data were presented as mean \pm SD. Different letters indicate significant differences ($P < 0.05$). Conversely, there is no significant difference between the same letters.

Effects of MPs on the Activities of Antioxidative Enzymes of Soiny Mullet Larvae

The effect of MPs exposure on the SOD activities of soiny mullet was shown in Figure 5. At 3 dpe, the SOD activity in SMPs was significantly lower than that in CTRL (P<0.05). However, no significant difference was observed for the SOD activity between the LMPs and CTRL (P>0.05). At 7 and 14 dpe, the MPs exposure significantly reduced the SOD activity in LMPs and SMPs (P<0.05)(Figure 5). The SOD activity reached lowest values in the LMPs, compared with that of CTRL and SMPs (P<0.05).

The MPs exposure slightly increased the CAT activity in LMPs at 3 dpe, but shared no significant change with other groups (P>0.05). However, the CAT

activity in SMPs was inhibited and significantly lower than that in the LMPs and CTRL (P<0.05). At 7 dpe and 14 dpe, the CAT activities of SMPs and LMPs were significantly lower than that of CTRL (P<0.05) (Figure 6).

Effects MPs on the Activities of Immune Enzymes of Soiny Mullet Larvae

The effect of MPs exposure on the ACP activities of soiny mullet was shown in Figure 7. MPs exposure significantly induced the ACP activities throughout the test period, compared with that of CTRL (P<0.05). As shown in Figure 7, the ACP activity in SMPs was significantly higher than that of LMPs (P<0.05), indicating that the ACP activity was mostly affected by the small particle size MPs.




Figure 5. Effects of exposure to MPs with different particle sizes on SOD activity of soiny mullet (*Liza haematocheila*) larvae (n=9). Data were presented as mean ± SD. Different letters indicate significant differences (P<0.05). Conversely, there is no significant difference between the same letters.




Figure 6. Effects of exposure to MPs with different particle sizes on CAT activity of soiny mullet (*Liza haematocheila*) larvae (n=9). Data were presented as mean ± SD. Different letters indicate significant differences (P<0.05). Conversely, there is no significant difference between the same letters.

Similarly, the MPs exposure increased the ALP activities of soiny mullet larvae throughout the trial period (Figure 8). At 3 dpe, the ALP activities in LMPs and SMPs were significantly higher than that of CTRL ($P < 0.05$). At 7 dpe, the ALP activity in LMPs shared similar level as that in CTRL ($P > 0.05$), while that in the SMPs was remained higher than that in the CTRL ($P < 0.05$). At 14 dpe, the ALP activities in LMPs and SMPs were significantly higher than that in CTRL ($P < 0.05$) (Figure 8).

Discussion

Effects of MPs on the Activities of Digestive Enzymes of Soiny Mullet Larvae

Digestive enzymes play important roles in the digestion and absorption of nutrients of animals

(Ahmadifaret et al., 2020). Various studies have shown that MPs challenge can negatively affect the activities of digestive enzymes of several fish species, such as orange-spotted grouper (*Epinephelus coioides* Hamilton, 1822) (Wang et al., 2022a), guppy (*Poecilia reticulata* reticulata W. Peters, 1859) (Huang et al., 2020), and large yellow croaker (*Larimichthys crocea* Richardson, 1846) (Gu et al., 2020). PPS and α -AMS are two important digestive enzymes involved in the digestion and absorption of proteins and carbohydrates. PPS is a digestive protease, secreted by the host cell, that functions to break down proteins in food into small peptide fragments. α -AMS is one of the extracellular enzymes that play an important role in the degradation of starch in living organisms. It acts on the α -1,4 glycosidic bond of the starch chain to break down starch into dextrans and reducing sugars. Our study found that MPs exposure inhibited the activities of PPS and α -AMS




Figure 7. Effects of exposure to MPs with different particle sizes on ACP activity of soiny mullet (*Liza haematocheila*) larvae (n=9). Data were presented as mean \pm SD. Different letters indicate significant differences ($P < 0.05$). Conversely, there is no significant difference between the same letters.




Figure 8. Effects of exposure to MPs with different particle sizes on AKP activity of soiny mullet (*Liza haematocheila*) larvae (n=9). Data were presented as mean \pm SD. Different letters indicate significant differences ($P < 0.05$). Conversely, there is no significant difference between the same letters.

in soiny mullet. Similar results were also observed in discus fish (*Symphysodon aequifasciatus* Pellegrin, 1904) after MPs (20µg/L; 200 µg/L) challenge for 28 days (Zhang et al., 2022). The reason for the activity inhibiting by the MPs exposure might be the accumulation of MPs in the gut, which causes gut blockage to produce a feeling of satiety and reduce food intake (Lu et al., 2016). When there were fewer proteins and carbohydrates in the food, there was less need for these enzymes that break down proteins and carbohydrates in the fish, which affected the activity of digestive enzymes. In addition, we found that the PPS activity was more strongly responsive to small MPs (0.5µm), while the α-AMS activity was more strongly responsive to large MPs (6µm). We speculated that MPs with different particle sizes could have various effects on different flora in the gut, which cause different microbiota dysbiosis and finally resulting in inconsistent levels of digestion and absorption of different nutrients by the body, and different digestive enzymes showing different activities, which was previously reported in the studies of Liu et al. (2019), Huang et al. (2020), Huang et al. (2022a) and Huang et al. (2022b).

Effects of MPs on the Activities of Antioxidative Enzymes of Soiny Mullet Larvae

The detoxification process in animals is the result of the joint action of multiple antioxidant defense enzymes. SOD and CAT are antioxidant enzymes involved in scavenging superoxide radicals. SOD firstly converts the O_2^- into H_2O_2 to maintain the normal physiological activities of cells and the body and prevent excessive radicals from damaging macromolecules such as proteins, lipids, and DNA (Geracitano et al., 2002). Then, the CAT splits H_2O_2 into H_2O and O_2 . When facing the stress of toxic substances in the environment, the body will induce an increasing activity of antioxidant enzymes to adapt to the environment change (Ajitha et al., 2021). Studies showed that 90-day exposure to food rich with MPs increased the SOD and CAT activities and induced oxidative stress in gilthead seabream (*Sparus aurata* Linnaeus, 1758) (Capó et al., 2021). In the present study, MPs exposure induced a significant reduction in the SOD and CAT activities of soiny mullet. Similarly, exposure to MPs (0.5 µm, 100µg/L; 0.5 µm, 1000µg/L; 5 µm, 100µg/L; 5 µm, 1000µg/L) for 21 days decreased the SOD and CAT activities and caused oxidative damage of loach (*Paramisgurnus dabryanus* Dabry de Thiersant, 1872) (Wang et al., 2022b). A study in common carp showed a reduction in SOD activity after an early rise in target organs, following MP exposure because ROS produced in tissues could not be removed instantly (Xia et al., 2020). However, Banaei (2022) found that exposure to MPs (175, 350, 700, 1400 µg/L) for 30 days, the activity of SOD was increased, while the activity of CAT was decreased, and oxidative stress was induced in common carp (*Cyprinus carpio* Linnaeus, 1758). High concentration and weak antioxidant

capacity of soiny mullet larvae might be the reasons for these different results. In addition, we found that the SOD and CAT activities in SMPs were lower than those in LMPs, which might have been caused by the small particles of MPs (0.5 µm), which are more likely to accumulate in tissues and thus exhibit stronger cytotoxicity as previously explained by Choi & Hu (2008), Jeong et al. (2016) and Lu et al. (2016), leading to reactive oxygen species (ROS) production by NADPH oxidase, SOD converts superoxide radicals to hydrogen peroxide, and CAT converts hydrogen peroxide to water and oxygen as the prime protection mechanism against oxidative stress (Sahabuddin, et al., 2023).

Effects MPs on the Activities of Immune Enzymes of Soiny Mullet Larvae

Fish innate immunity is first defense line against pathogens or stress. ACP and ALP are important innate immune related enzymes (NavinChandran et al., 2014; Suvetha et al., 2015). ACP is a marker enzyme of cellular lysosomes and involves in the inflammatory response (Lallès, 2019; He et al., 2021), commonly found in macrophages and neutrophils, which are key components of the fish immune system. ALP is associated with damage to cell membranes, it is involved in a variety of physiological processes, including phosphate metabolism and transport. In the immune response, ALP can be involved in cell signalling and cell activation processes. ALP is also associated with damage to cell membranes (Banaei et al., 2019). Both ACP and ALP involve in the cellular phosphorylation and dephosphorylation, as well as reflecting the metabolic rate of toxic substances in the body (Ellis et al., 2011; Lu et al., 2019). Therefore, ACP and ALP are usually used as biomarkers of environmental pollution (Wen et al., 2018). In the present study, we found that MPs exposure significantly increased the activities of ACP and ALP in soiny mullet. Similar results were also observed in other animals. The study of Liu et al. (2019) revealed that 14 days of MPs exposure (5 µm, 0.4 mg/L) significantly increased the activities of ACP and ALP and activated the immune system in Chinese mitten crab. Wen (2018) also found that MPs exposure (50µg/L; 500µg/L) for 14 days increased the activities of ACP and ALP and induced the immune responses in the tested fish larvae. Our result are in agreement with Pitt et al. (2018) and indicated that the innate immunity of animals is triggered to adapt to MPs in the environment, When fish were exposed to pathogens or inflammation, activation of the immune system increased the number of these immune cells, which may lead to elevated enzyme activity.

Conclusions

The effects of different size MPs on the activities of digestive enzymes, antioxidant enzymes and immune enzymes of soiny mullet larvae were investigated. To the best of our knowledge, this is the first report about the

effect of MPs on the soiny mullet larvae. Our results demonstrated that the digestion and antioxidant system of soiny mullet larvae were inhibited by MPs exposures. And, the activities of immune enzymes were induced. Both sizes of MPs were toxic to soiny mullet larvae, and the small particles of MPs (0.5 µm) showed to be more toxic compared with the large particles of MPs (6 µm). These results provide solide basis for understanding the detriment of MPs in larvae fish. Notedly, the detoxification mechanism of microplastic on larvae fish needs further investigation.

Ethical Statement

Not applicable

Funding Information

This study was supported by the Major Projects of Natural Science Research for University and Colleges in Jiangsu Province (Grant No. 21KJA240001) and partial by the Key Research and Development Program of Jiangsu Province (Grant No. BE2018348) and the Projects for the High-quality Development of Fishery Industry of Yancheng City (Grant No. YCSCYJ20210014). Q. Z was supported financially by the projects for 'six talents' of Jiangsu Province (Grant No. NY-115).

Author Contribution

Zhitao Qi: Supervision, Writing- Reviewing and Editing. Yingying Zhang; Shengyuan Zhang: Enzymes activities analysis. Yanming Sui: microplastics preparation and writing. Ke Sun; Qihuan Zhang: data analysis. Eakapol Wangkahart; Zisheng Wang: Supervision and writing.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements

Not applicable

References

- Ahmadifar, E., Sadegh, T. H., Dawood, M. A., Dadar, M., & Sheikhzadeh, N., (2020). The effects of dietary *Pediococcus pentosaceus* on growth performance, hemato-immunological parameters and digestive enzyme activities of common carp (*Cyprinus carpio*). *Aquaculture*, 516, 734656. <https://doi.org/10.1016/j.aquaculture.2019.734656>
- Ajitha, V., Sreevidya, C. P., Sarasan, M., Park, J. C., Mohandas, A., Singh, I. S. B., Puthumana, J., & Lee, J. S., (2021). Effects of zinc and mercury on ROS-mediated oxidative stress-induced physiological impairments and antioxidant responses in the microalga *Chlorella vulgaris*. *Environmental Science and Pollution Research*, 28(25), 32475-32492. <https://doi.org/10.1007/s11356-021-12950-6>
- Andrady, A. L., (2011). Microplastics in the marine environment. *Marine Pollution Bulletin*, 62(8), 1596-1605. <https://doi.org/10.1016/j.marpolbul.2011.05.030>
- Banaee, M., Soltanian, S., Sureda, A., Gholamhosseini, A., Hagi, B. N., Akhlaghi, M., Akhlaghi, M., & Derikvandy, A., (2019). Evaluation of single and combined effects of cadmium and micro-plastic particles on biochemical and immunological parameters of common carp (*Cyprinus carpio*). *Chemosphere*, 236, 124335. <https://doi.org/10.1016/j.chemosphere.2019.07.066>
- Banaei, M., Forouzanfar, M., & Jafarinaia, M., (2022). Toxic effects of polyethylene microplastics on transcriptional changes, biochemical response, and oxidative stress in common carp (*Cyprinus carpio*). *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology*, 261, 109423. <https://doi.org/10.1016/j.cbpc.2022.109423>
- Beiras, R., Bellas, J., Cachot, J., Cormier, B., Cousin, X., Engwall, M., Gambardella, C., Garaventa, F., Keiter, S., Le Bihanic, F., López-Ibáñez, S., Piazza, V., Rial, D., Tato, T., & Vidal-Liñán, L., (2018). Ingestion and contact with polyethylene microplastics does not cause acute toxicity on marine zooplankton. *Journal of Hazardous Materials*, 360, 452-460. <https://doi.org/10.1016/j.jhazmat.2018.07.101>
- Blewett, T. A., & Leonard, E. M., (2017). Mechanisms of nickel toxicity to fish and invertebrates in marine and estuarine waters. *Environmental Pollution*, 223, 311-322. <https://doi.org/10.1016/j.envpol.2017.01.028>
- Bravo Rebolledo, E. L., Van Franeker, J. A., Jansen, O. E., & Brasseur, S. M., (2013). Plastic ingestion by harbour seals (*Phoca vitulina*) in The Netherlands. *Marine Pollution Bulletin*, 67(1-2), 200-202. <https://doi.org/10.1016/j.marpolbul.2012.11.035>
- Browne, M. A., Dissanayake, A., Galloway, T. S., Lowe, D. M., & Thompson, R. C., (2008). Ingested microscopic plastic translocates to the circulatory system of the mussel, *Mytilus edulis* (L.). *Environmental Science & Technology*, 42(13), 5026-5031. <https://doi.org/10.1021/es800249a>
- Capó, X., Company, J. J., Alomar, C., Compa, M., Sureda, A., Grau, A., Hansjosten, B., López-Vázquez, J., Quintana, J. B., Rodil, R., & Deudero, S., (2021). Long-term exposure to virgin and seawater exposed microplastic enriched-diet causes liver oxidative stress and inflammation in gilthead seabream *Sparus aurata*, Linnaeus 1758. *Science of the Total Environment*, 767, 144976. <https://doi.org/10.1016/j.scitotenv.2021.144976>
- Choi, O., & Hu, Z., (2008). Size dependent and reactive oxygen species related nanosilver toxicity to nitrifying bacteria. *Environmental Science & Technology*, 42(12), 4583-4588. <https://doi.org/10.1021/es703238h>
- Couillard, C. M., Laplatte, B., & Pelletier, E., (2009). A fish bioassay to evaluate the toxicity associated with the ingestion of benzo[a]pyrene-contaminated benthic prey. *Environmental Toxicology and Chemistry*, 28(4), 772-781. <https://doi.org/10.1897/08-092R.1>
- Desforges, J. P., Galbraith, M., & Ross, P. S., (2015). Ingestion of microplastics by zooplankton in the Northeast Pacific Ocean. *Archives of Environmental Contamination and Toxicology*, 69(3), 320-330. <https://doi.org/10.1007/s00244-015-0172-5>
- Devriese, L. I., van der Meulen, M. D., Maes, T., Bekaert, K., Paul-Pont, I., Frère, L., Robbens, J., & Vethaak, A. D.,

- (2015). Microplastic contamination in brown shrimp (*Crangon crangon*, Linnaeus 1758) from coastal waters of the Southern North Sea and Channel area. *Marine Pollution Bulletin*, 98(1-2), 179-187. <https://doi.org/10.1016/j.marpolbul.2015.06.051>
- Dobaradaran, S., Schmidt, T. C., Nabipour, I., Khajeahmadi, N., Tajbakhsh, S., Saeedi, R., Javad Mohammadi, M., Keshtkar, M., Khorsand, M., & Faraji Ghasemi, F., (2018). Characterization of plastic debris and association of metals with microplastics in coastline sediment along the Persian Gulf. *Waste Management*, 78, 649-658. <https://doi.org/10.1016/j.wasman.2018.06.037>
- Ellis, R. P., Parry, H., Spicer, J. I., Hutchinson, T. H., Pipe, R. K., & Widdicombe, S., (2011). Immunological function in marine invertebrates: responses to environmental perturbation. *Fish & Shellfish Immunology*, 30(6), 1209-1222. <https://doi.org/10.1016/j.fs.2011.03.017>
- Fossi, M. C., Panti, C., Guerranti, C., Coppola, D., Giannetti, M., Marsili, L., & Minutoli, R., (2012). Are baleen whales exposed to the threat of microplastics? A case study of the Mediterranean fin whale (*Balaenoptera physalus*). *Marine Pollution Bulletin*, 64(11), 2374-2379. <https://doi.org/10.1016/j.marpolbul.2012.08.013>
- Geracitano, L., Monserrat, J. M., & Bianchini, A., (2002). Physiological and antioxidant enzyme responses to acute and chronic exposure of *Laonereis acuta* (Polychaeta, Nereididae) to copper. *Journal of Experimental Marine Biology and Ecology*, 277(2), 145-156. [https://doi.org/10.1016/S0022-0981\(02\)00306-4](https://doi.org/10.1016/S0022-0981(02)00306-4)
- Gu, H., Wang, S., Wang, X., Yu, X., Hu, M., Huang, W., & Wang, Y., (2020). Nanoplastics impair the intestinal health of the juvenile large yellow croaker *Larimichthys crocea*. *Journal of Hazardous Materials*, 397, 122773. <https://doi.org/10.1016/j.jhazmat.2020.122773>
- Güven, O., Gökdağ, K., Jovanović, B., & Kideys, A. E., (2017). Microplastic litter composition of the Turkish territorial waters of the Mediterranean Sea, and its occurrence in the gastrointestinal tract of fish. *Environmental Pollution*, 223, 286-294. <https://doi.org/10.1016/j.envpol.2017.01.025>
- Hanachi, P., Karbalaeei, S., & Yu, S. J., (2021). Combined polystyrene microplastics and chlorpyrifos decrease levels of nutritional parameters in muscle of rainbow trout (*Oncorhynchus mykiss*). *Environmental Science and Pollution Research*, 28(45), 64908-64920. <https://doi.org/10.1007/s11356-021-15536-4>
- He, R. Z., Li, Z. C., Li, S. Y., & Li, A. X., (2021). Development of an immersion challenge model for *Streptococcus agalactiae* in Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 531, 735877. <https://doi.org/10.1016/j.aquaculture.2020.735877>
- Huang, J. N., Wen, B., Zhu, J. G., Zhang, Y. S., Gao, J. Z., & Chen, Z. Z., (2020). Exposure to microplastics impairs digestive performance, stimulates immune response and induces microbiota dysbiosis in the gut of juvenile guppy (*Poecilia reticulata*). *Science of the Total Environment*, 733, 138929. <https://doi.org/10.1016/j.scitotenv.2020.138929>
- Huang, J. N., Wen, B., Xu, L., Ma, H. C., Li, X. X., Gao, J. Z., & Chen, Z. Z., (2022a). Micro/nano-plastics cause neurobehavioral toxicity in discus fish (*Symphysodon aequifasciatus*): Insight from brain-gut-microbiota axis. *Journal of Hazardous Materials*, 421, 126830. <https://doi.org/10.1016/j.jhazmat.2021.126830>
- Huang, J. N., Zhang, Y., Xu, L., He, K. X., Wen, B., Yang, P. W., Ding, J. Y., Li, J. Z., Ma, H. C., Gao, J. Z., & Chen, Z. Z., (2022b). Microplastics: A tissue-specific threat to microbial community and biomarkers of discus fish (*Symphysodon aequifasciatus*). *Journal of Hazardous Materials*, 424, 127751. <https://doi.org/10.1016/j.jhazmat.2021.127751>
- Kim, J. H., Yu, Y. B., & Choi, J. H., (2021). Toxic effects on bioaccumulation, hematological parameters, oxidative stress, immune responses and neurotoxicity in fish exposed to microplastics: A review. *Journal of Hazardous Materials*, 413, 125423. <https://doi.org/10.1016/j.jhazmat.2021.125423>
- Jeong, C. B., Won, E. J., Kang, H. M., Lee, M. C., Hwang, D. S., Hwang, U. K., Zhou, B., Souissi, S., Lee, S. J., & Lee, J. S., (2016). Microplastic size-dependent toxicity, oxidative stress induction, and p-JNK and p-p38 activation in the monogonont rotifer (*Brachionus koreanus*). *Environmental Science & Technology*, 50(16), 8849-8857. <https://doi.org/10.1021/acs.est.6b01441>
- Lallès, J. P., (2019). Biology, environmental and nutritional modulation of skin mucus alkaline phosphatase in fish: A review. *Fish & Shellfish Immunology*, 89, 179-186. <https://doi.org/10.1016/j.fsi.2019.03.053>
- Le, L. L., Liu, M. T., Song, Y., Lu, S. B., Hu, J. N., Cao, C. J., Xie, B., Shi, H. H., & He, D. F., (2018). Polystyrene (nano) microplastics cause size-dependent neurotoxicity, oxidative damage and other adverse effects in *Caenorhabditis elegans*. *Environmental Science-Nano*, 5(8), 2009-2020. <https://doi.org/10.1039/c8en00412a>
- Liu, Z., Yu, P., Cai, M., Wu, D., Zhang, M., Chen, M., & Zhao, Y., (2019). Effects of microplastics on the innate immunity and intestinal microflora of juvenile *Eriocheir sinensis*. *Science of the Total Environment*, 685, 836-846. <https://doi.org/10.1016/j.scitotenv.2019.06.265>
- Lu, Y., Zhang, Y., Deng, Y., Jiang, W., Zhao, Y., Geng, J., Ding, L., & Ren, H., (2016). Uptake and accumulation of polystyrene microplastics in zebrafish (*Danio rerio*) and toxic effects in liver. *Environmental Science & Technology*, 50(7), 4054-4060. <https://doi.org/10.1021/acs.est.6b00183>
- Lu, J., Qi, C., Limbu, S. M., Han, F., Yang, L., Wang, X., Qin, J. G., & Chen, L., (2019). Dietary mannan oligosaccharide (MOS) improves growth performance, antioxidant capacity, non-specific immunity and intestinal histology of juvenile Chinese mitten crabs (*Eriocheir sinensis*). *Aquaculture*, 510, 337-346. <https://doi.org/10.1016/j.aquaculture.2019.05.048>
- Mattsson, K., Johnson, E. V., Malmendal, A., Linse, S., Hansson, L. A., & Cedervall, T., (2017). Brain damage and behavioural disorders in fish induced by plastic nanoparticles delivered through the food chain. *Scientific Reports*, 7(1), 11452. <https://doi.org/10.1038/s41598-017-10813-0>
- Mazurais, D., Ernande, B., Quazuguel, P., Severe, A., Huelvan, C., Madec, L., Mouchel, O., Soudant, P., Robbens, J., Huvet, A., & Zambonino-Infante, J., (2015). Evaluation of the impact of polyethylene microbeads ingestion in European sea bass (*Dicentrarchus labrax*) larvae. *Marine Environmental Research*, 112, 78-85. <https://doi.org/10.1016/j.marenvres.2015.09.009>
- Murray, F., & Cowie, P. R. (2011). Plastic contamination in the decapod crustacean *Nephrops norvegicus* (Linnaeus, 1758). *Marine Pollution Bulletin*, 62(6), 1207-1217. <https://doi.org/10.1016/j.marpolbul.2011.03.032>
- Murray, R. H., Erler, D. V., & Eyre, B. D., (2015). Nitrous oxide

- fluxes in estuarine environments: response to global change. *Global Change Biology*, 21(9), 3219-3245. <https://doi.org/10.1111/gcb.12923>
- NavinChandran, M., Iyapparaj, P., Moovendhan, S., Ramasubburayan, R., Prakash, S., Immanuel, G., & Palavesam, A., (2014). Influence of probiotic bacterium *Bacillus cereus* isolated from the gut of wild shrimp *Penaeus monodon* in turn as a potent growth promoter and immune enhancer in *P. monodon*. *Fish & Shellfish Immunology*, 36(1), 38-45.
- Nobre, C. R., Santana, M., Maluf, A., Cortez, F. S., Cesar, A., Pereira, C., & Turra, A., (2015). Assessment of microplastic toxicity to embryonic development of the sea urchin *Lytechinus variegatus* (Echinodermata: Echinoidea). *Marine Pollution Bulletin*, 92(1-2), 99-104. <https://doi.org/10.1016/j.marpolbul.2014.12.050>
- Pedà, C., Caccamo, L., Fossi, M. C., Gai, F., Andaloro, F., Genovese, L., Perdichizzi, A., Romeo, T., & Maricchiolo, G., (2016). Intestinal alterations in European sea bass *Dicentrarchus labrax* (Linnaeus, 1758) exposed to microplastics: Preliminary results. *Environmental Pollution*, 212, 251-256. <https://doi.org/10.1016/j.envpol.2016.01.083>
- Pitt, J. A., Kozal, J. S., Jayasundara, N., Massarsky, A., Trevisan, R., Geitner, N., Wiesner, M., Levin, E. D., & Di Giulio, R. T., (2018). Uptake, tissue distribution, and toxicity of polystyrene nanoparticles in developing zebrafish (*Danio rerio*). *Aquatic Toxicology*, 194, 185-194. <https://doi.org/10.1016/j.aquatox.2017.11.017>
- Qi, Z., Pi, X., Xu, Y., Zhang, Q., Wangkahart, E., Meng, F., & Wang, Z., (2022). Molecular characterization of the evolutionary conserved signaling intermediate in Toll pathways (ECSIT) of soiny mullet (*Liza haematocheila*). *Fish & Shellfish Immunology*, 130, 79-85. <https://doi.org/10.1016/j.fsi.2022.09.009>
- Rummel, C. D., Löder, M. G., Fricke, N. F., Lang, T., Griebeler, E. M., Janke, M., & Gerdt, G., (2016). Plastic ingestion by pelagic and demersal fish from the North Sea and Baltic Sea. *Marine Pollution Bulletin*, 102(1), 134-141. <https://doi.org/10.1016/j.marpolbul.2015.11.043>
- Sahabuddin, E. S., Noreen, A., Daabo, H. M. A., Kandeel, M., Saleh, M. M., Al-qaim, Z. H., Jawad, M. A., Sivaraman, R., Fenjan, M. N., Mustafa, Y. F., Heidary, A., Abarghouei, S., & Norbakhsh, M., (2023). Microplastic and oil pollutant agglomerates synergistically intensify toxicity in the marine fish, Asian seabass, Lates calcalifer. *Environmental Toxicology and Pharmacology*, 98, 104059. <https://doi:10.1016/j.etap.2022.104059>
- Shahriari Moghadam, M., Abtahi, B., Rezaei, S., & Rahdari, A., (2014). Early ontogenetic development of digestive system in *Schizothorax zarudnyi* Nikolskii, 1897 (Actinopterygii: Cyprinidae) larvae. *Italian Journal of Zoology*, 81(2), 194-203. <https://doi.org/10.1080/11250003.2014.903304>
- Soletchnik, P., Ropert, M., Mazurié, J., Fleury, P. G., & Le Coz, F., (2007). Relationships between oyster mortality patterns and environmental data from monitoring databases along the coasts of France. *Aquaculture*, 271(1-4), 384-400. <https://doi.org/10.1016/j.aquaculture.2007.02.049>
- Steer, M., Cole, M., Thompson, R. C., & Lindeque, P. K., (2017). Microplastic ingestion in fish larvae in the western English Channel. *Environmental Pollution*, 226, 250-259. <https://doi.org/10.1016/j.envpol.2017.03.062>
- Sui, Q., Zhang, L., Xia, B., Chen, B., Sun, X., Zhu, L., Wang, R., & Qu, K., (2020). Spatiotemporal distribution, source identification and inventory of microplastics in surface sediments from Sanggou Bay, China. *Science of the Total Environment*, 723, 138064. <https://doi.org/10.1016/j.scitotenv.2020.138064>
- Sussarellu, R., Suquet, M., Thomas, Y., Lambert, C., Fabioux, C., Pernet, M. E., Le Goïc, N., Quillien, V., Mingant, C., Epelboin, Y., Corporeau, C., Guyomarch, J., Robbens, J., Paul-Pont, I., Soudant, P., & Huvet, A., (2016). Oyster reproduction is affected by exposure to polystyrene microplastics. *Proceedings of the National Academy of Sciences of the United States of America*, 113(9), 2430-2435. <https://doi.org/10.1073/pnas.1519019113>
- Suvetha, L., Saravanan, M., Hur, J. H., Ramesh, M., & Krishnapriya, K., (2015). Acute and sublethal intoxication of deltamethrin in an Indian major carp, *Labeo rohita*: Hormonal and enzymological responses. *The Journal of Basic & Applied Zoology*, 72, 58-65. <https://doi.org/10.1016/j.jobaz.2015.04.005>
- Tata, T., Belabed, B. E., Bououdina, M., & Bellucci, S., (2020). Occurrence and characterization of surface sediment microplastics and litter from North African coasts of Mediterranean Sea: Preliminary research and first evidence. *Science of the Total Environment*, 713, 136664. <https://doi.org/10.1016/j.scitotenv.2020.136664>
- Thompson, R. C., Olsen, Y., Mitchell, R. P., Davis, A., Rowland, S. J., John, A. W., McGonigle, D., & Russell, A. E., (2004). Lost at sea: where is all the plastic?. *Science*, 304(5672), 838. <https://doi.org/10.1126/science.1094559>
- Tourinho, P. S., Ivar do Sul, J. A., & Fillmann, G., (2010). Is marine debris ingestion still a problem for the coastal marine biota of southern Brazil?. *Marine Pollution Bulletin*, 60(3), 396-401. <https://doi.org/10.1016/j.marpolbul.2009.10.013>
- Van Cauwenberghe, L., & Janssen, C. R., (2014). Microplastics in bivalves cultured for human consumption. *Environmental Pollution*, 193, 65-70. <https://doi.org/10.1016/j.envpol.2014.06.010>
- Wang, Q., Huang, F., Liang, K., Niu, W., Duan, X., Jia, X., Wu, X., Xu, P., & Zhou, L., (2022a). Polystyrene nanoplastics affect digestive function and growth in juvenile groupers. *Science of the Total Environment*, 808, 152098. <https://doi.org/10.1016/j.scitotenv.2021.152098>
- Wang, X., Jian, S., Zhang, S., Wu, D., Wang, J., Gao, M., Sheng, J., & Hong, Y., (2022b). Enrichment of polystyrene microplastics induces histological damage, oxidative stress, Keap1-Nrf2 signaling pathway-related gene expression in loach juveniles (*Paramisgurnus dabryanus*). *Ecotoxicology and Environmental Safety*, 237, 113540. <https://doi.org/10.1016/j.ecoenv.2022.113540>
- Wen, B., Jin, S. R., Chen, Z. Z., Gao, J. Z., Liu, Y. N., Liu, J. H., & Feng, X. S., (2018). Single and combined effects of microplastics and cadmium on the cadmium accumulation, antioxidant defence and innate immunity of the discus fish (*Symphysodon aequifasciatus*). *Environmental Pollution*, 243, 462-471. <https://doi.org/10.1016/j.envpol.2018.09.029>
- Wijesiri, B., Liu, A., He, B., Yang, B., Zhao, X., Ayoko, G., & Goonetilleke, A., (2019). Behaviour of metals in an urban river and the pollution of estuarine environment. *Water Research*, 64, 114911. <https://doi.org/10.1016/j.watres.2019.114911>

- Wu, N., Zhang, Y., Zhang, X., Zhao, Z., He, J., Li, W., Ma, Y., & Niu, Z., (2019). Occurrence and distribution of microplastics in the surface water and sediment of two typical estuaries in Bohai Bay, China. *Environmental Science: Processes & Impacts*, 21(7), 1143-1152. <https://doi.org/10.1039/c9em00148d>
- Xia, X., Sun, M., Zhou, M., Chang, Z., & Li, L., (2020). Polyvinyl chloride microplastics induce growth inhibition and oxidative stress in *Cyprinus carpio* var. larvae. *Science of the Total Environment*, 716, 136479. <https://doi.org/10.1016/j.scitotenv.2019.136479>
- Xie, M., Xu, P., Zhou, W., Xu, X., Li, H., He, W., Yue, W., Zhang, L., Ding, D., & Suo, A., (2022). Impacts of conventional and biodegradable microplastics on juvenile *Lates calcarifer*: Bioaccumulation, antioxidant response, microbiome, and proteome alteration. *Marine Pollution Bulletin*, 179, 113744. <https://doi.org/10.1016/j.marpolbul.2022.113744>
- Yan, M., Nie, H., Xu, K., He, Y., Hu, Y., Huang, Y., & Wang, J., (2019). Microplastic abundance, distribution and composition in the Pearl River along Guangzhou city and Pearl River estuary, China. *Chemosphere*, 217, 879-886. <https://doi.org/10.1016/j.chemosphere.2018.11.093>
- Yan, W., Hamid, N., Deng, S., Jia, P. P., & Pei, D. S., (2020). Individual and combined toxicogenetic effects of microplastics and heavy metals (Cd, Pb, and Zn) perturb gut microbiota homeostasis and gonadal development in marine medaka (*Oryzias melastigma*). *Journal of Hazardous Materials*, 397, 122795. <https://doi.org/10.1016/j.jhazmat.2020.122795>
- Yang, H., Xiong, H., Mi, K., Xue, W., Wei, W., & Zhang, Y., (2020). Toxicity comparison of nano-sized and micron-sized microplastics to goldfish *Carassius auratus* Larvae. *Journal of Hazardous Materials*, 388, 122058. <https://doi.org/10.1016/j.jhazmat.2020.122058>
- Zhang, J., Zhang, C., Deng, Y., Wang, R., Ma, E., Wang, J., Bai, J., Wu, J., & Zhou, Y., (2019). Microplastics in the surface water of small-scale estuaries in Shanghai. *Marine Pollution Bulletin*, 149, 110569. <https://doi.org/10.1016/j.marpolbul.2019.110569>
- Zhao, S., Zhu, L., Wang, T., & Li, D., (2014). Suspended microplastics in the surface water of the Yangtze Estuary System, China: first observations on occurrence, distribution. *Marine Pollution Bulletin*, 86(1-2), 562-568. <https://doi.org/10.1016/j.marpolbul.2014.06.032>
- Zhao, S., Zhu, L., & Li, D., (2015). Microplastic in three urban estuaries, China. *Environmental Pollution*, 206, 597-604. <https://doi.org/10.1016/j.envpol.2015.08.027>
- Zhao, F., Guo, M., Zhang, M., Duan, M., Zheng, J., Liu, Y., & Qiu, L., (2022). Sub-lethal concentration of metamifop exposure impair gut health of zebrafish (*Danio rerio*). *Chemosphere*, 303, 135081. <https://doi.org/10.1016/j.chemosphere.2022.135081>