




# Development and Validation of 31 SNP Markers of Topmouth Culter, *Culter alburnus* by Whole-Genome Resequencing

Huan Wang<sup>1</sup> , Huaxing Zhou<sup>1</sup>, Yuting Hu<sup>1</sup>, Jun Ling<sup>1</sup> , He Jiang<sup>1</sup>, Guoqing Duan<sup>1,\*</sup> 

<sup>1</sup>Anhui Academy of Agricultural Sciences, Fisheries Research Institution, Anhui Key Laboratory of Aquaculture and Stock Enhancement, Hefei, China, 230031 Institute of Fisheries Science, Anhui Academy of Agricultural Sciences, Nongke South Road, Hefei, Anhui, China.

## How to Cite

Wang, H., Zhou, H., Hu, Y., Ling, J., Jiang, H., Duan G. (2025). Development and Validation of 31 SNP Markers of Topmouth Culter, *Culter alburnus* by Whole-Genome Resequencing. *Turkish Journal of Fisheries and Aquatic Sciences*, 25(4), TRJFAS25574. <https://doi.org/10.4194/TRJFAS25574>

## Article History

Received 11 February 2024

Accepted 08 November 2024

First Online 12 December 2024

## Corresponding Author

E-mail: duangq2010@126.com

## Keywords

Topmouth culter,  
SNP markers,  
Population genetics,  
Whole genome resequencing

## Abstract

*Culter alburnus* is one of the most important commercial freshwater fishes and widely distributed in China. In this study, a total of 31 novel single nucleotide polymorphism (SNP) markers were developed based on whole-genome resequencing. The analysis of the 31 SNP genotypes in 60 fish showed that the observed heterozygosity and expected heterozygosity ranged from 0.283 to 0.650 and 0.280 to 0.504, respectively. The minor allele frequency ranged from 0.167 to 0.458, and the polymorphism information content ranged from 0.239 to 0.375. Among these SNPs, only one SNP were found to deviate from Hardy-Weinberg equilibrium significantly ( $P < 0.01$ ). These SNP markers could be a valuable tool for population conservation in this species.

## Introduction

The topmouth culter (*Culter alburnus*) belongs to the order Cypriniformes, family Xenocyprididae and inhabits the major rivers, lakes, and reservoirs in China (Li et al., 2018). It is the most important commercial freshwater fishes, among the species of the genus *Culter*, *C. alburnus* has the longest body size. Because *C. alburnus* has the advantage of rapid growth, low oxygen tolerance, and disease resistance, hybridization breeding with *Megalobrama amblycephala* became important (National Reg no: GS-02-001-2020). However, wild *C. alburnus* populations are declining and their germplasm resources have been threatened in recent years because of the overfishing and habitat changes caused by environmental pollution and water-

related construction (Qi et al., 2013). This has necessitated population genetic investigations on *C. alburnus* to protect its wild resources.

Previous studies have performed valuable genetic evaluations of this species using mitochondrial markers, amplified fragment length polymorphism (AFLP) markers, and simple sequence repeat (SSR) markers (Wang et al., 2007; Chen et al., 2009; Liu et al., 2013; Sun et al., 2021). Wild *C. alburnus* in China has low genetic diversity, and the significant genetic differentiation, in addition, the genetic diversity of wild populations is lower than that of cultured populations (Qi et al., 2013; Qi et al., 2015). The historical effective population size of *C. alburnus* has been found to correlate with the uplift of the Tibetan Plateau, as indicated by demographic history reconstruction through whole-genome

resequencing (Jiang et al., 2023). Single nucleotide polymorphisms (SNPs) are characterized by a wide genomic distribution, high polymorphism levels, and good reproducibility, and have been used in molecular ecology, evolution, and conservation genetics (Vignal et al. 2002; Shan et al., 2022). SNP markers have significant advantages, such as highly abundance, wide distribution, site-specificity, and low error-detection rate, over SSR markers (Zhou et al., 2022). In this study new SNP loci of *C. alburnus* investigated for utilization on genetic evaluation of this specie.

## Materials and methods

In 2023, cultured *C. alburnus* samples were randomly selected from the Fishery Research Institute of the Anhui Academy of Agricultural Sciences, Hefei, Anhui, China. The parents of these cultured populations were wild populations originating from the Yangtze and Huaihe Rivers, which are two different water systems in China. Genomic DNA was extracted from the muscle tissue samples collected from 150 individuals using the TIANamp Genomic DNA Kit (DP304, Tiangen, Beijing, China), according to the manufacturer's instructions. DNA quality and concentration were determined using NanoDrop 2100 (Thermo Scientific, DE, USA) and 1.0% agarose-gel electrophoresis. Ninety DNA samples were randomly selected from the 150 samples and sequenced using an Illumina HiSeq 4000 platform. SNP detection was performed by gatk4 (<https://gatk.broadinstitute.org/hc/en-us/articles/360035890471-Hard-filtering-germline-short-variants>), and the SNP filter expression parameters were set as follows: QD<2.0, MQ<40.0, SOR>5.0, MQRankSum<-12.5, FS>60.0, QUAL<30 (DePristo et al., 2011). After quality control and strict filtration, 1060.57 Gb of clean data and 9561,979 putative SNPs were obtained. A total of 31 SNPs with high heterozygosity were selected to analyze the polymorphisms in the remaining 60 samples using PCR. A 20 µL PCR reaction volume contained 10 µL of 2×Master Mix, 0.5 µL each of the forward and reverse primers (10 mmol/L), 1 µL of DNA template (20 ng), and 8 µL of dd H<sub>2</sub>O. The PCR parameters were as follows: 95°C for 5 min; followed by 32 cycles of 94°C for 30 s, annealing temperature for 40 s, and 72°C for 50 s; and final extension at 72°C for 10 min.

The amplification products were sequenced using an ABI PRISM 3730 automated sequencer (Applied Biosystems, Foster City, CA, USA), and SNP genotypes were tested and analyzed using DNAMAN V6.0. The observed number of alleles ( $N_a$ ), observed heterozygosity ( $H_o$ ), expected heterozygosity ( $H_e$ ), minor allele frequency ( $MAF$ ), and deviations from the Hardy–Weinberg equilibrium ( $P_{HWE}$ ) were calculated using the chi-square test with POPGENE 3.2. Polymorphism information content ( $PIC$ ) was estimated from allele frequencies using Cervus 3.0 (Kalinowski et al., 2007).

## Results and Discussion

A total of 31 SNP markers were polymorphic and bi-allelic and were amplified successfully in the 60 individuals of *C. alburnus* (Table 1). The  $H_o$ ,  $H_e$ ,  $MAF$  and  $PIC$  of the SNPs were 0.283–0.650, 0.280–0.504, 0.167–0.458, and 0.239–0.375, respectively. The mean  $H_o$ ,  $H_e$  and  $PIC$  were 0.394, 0.362, and 0.292, respectively. These results indicate that cultured *C. alburnus* has a lower genetic diversity. In a previous study, the wild population of *C. alburnus* from Dianshan Lake (Shanghai, China) showed high genetic diversity measured using 11 microsatellite DNA markers; with mean  $H_o$  and  $H_e$  of 0.6288 and 0.7946, respectively (Chen et al., 2009). In addition, among the present SNPs, six loci deviated significantly from Hardy–Weinberg equilibrium (HWE) after Bonferroni adjustments ( $P<0.05$ ). Several factors, including random mating deviations, mutations, selection, genetic drift, and migration, can disrupt HWE (Zhang et al., 2024). In the current study involving cultured *C. alburnus*, deviations from HWE may be attributed to a lack of heterozygosity stemming from historical founder effects, artificial selection, and the impact of inbreeding (Chen et al., 2017). A study revealed that French rainbow trout (*Oncorhynchus mykiss*) exhibit a relatively high level of inbreeding, which is likely attributable in part to a founder effect and to “sweepstakes reproductive success.” (D’Ambrosio et al., 2019). These results suggested that the number of breeding parents should be increased and genetic exchanges among the populations should be promoted to improve the genetic diversity of cultured *C. alburnus* during the subsequent breeding processes. The SNPs identified in this study will be valuable tools for population conservation and management of *C. alburnus*.

## Ethical Statement

The experimental protocol was established, according to the ethical guidelines of the Basel Declaration and was approved by the Experimental Animal Welfare and Ethical of Anhui Academy of Agricultural Sciences (NO. AAAS 2024-06).

## Funding Information

This work was financially supported by the Young Talent Program of Anhui Academy of Agricultural Sciences (2023–2027) and the Wuhu City Science and Technology Planning Project (No. 2022ly15).

## Author Contribution

Huan Wang performed the experiment, designed, and analyzed the data, and wrote the original manuscript. Huaxing Zhou and Yuting Hu collected the materials, measured parameters and analyzed the data. Jun Ling and He Jiang collected the materials and

**Table 1.** Summary information of the 31 SNP markers developed for topmouth culter *Culter alburnus*

Locus	Primer sequences (5'-3')	Mutation type	Amplification length (bp)	<i>Ho</i>	<i>He</i>	<i>MAF</i>	<i>P<sub>HWE</sub></i>	<i>PIC</i>
JAOTNS010003905	F:AGATCTCGCACAGGTCGATT R:AGGTCAGCGCAGATGTTTCT	A/G	124	0.367	0.323	0.200	0.282	0.269
JAOTNS010003744	F:AACAGCACAGGGATTCTGTC R:ATGACATCCCAGCATCAGTG	A/G	138	0.333	0.484	0.400	0.015*	0.365
JAOTNS010003713	F:ACCTCTGCATCTGCCACTG R:CAGGAGCAGTCTCCATGTCA	T/C	146	0.533	0.495	0.433	0.548	0.371
JAOTNS010003696	F:TTCCACCACTTCATTACCA R:TGAGTGCTCGACGTAACG	T/A	157	0.417	0.352	0.225	0.146	0.288
JAOTNS010003632	F:TTTTGCTGGGACCTGTATCC R:CATATCCTTATGGATGCGCC	T/C	181	0.500	0.424	0.300	0.157	0.332
JAOTNS010003585	F:TCAATCTACAGGCAGGGGTC R:CGCAAAGTCTTGATTGCCA	T/C	251	0.383	0.313	0.192	0.073	0.262
JAOTNS010003549	F:CTCTACCGGTGAGATGGCTC R:ACAGGATCACTGCCCTTCA	A/T	182	0.333	0.280	0.167	0.132	0.239
JAOTNS010003508	F:AACCACTGCAGCCTGACTCT R:TCAGTTTGTGCGTGAGCAAT	A/G	177	0.350	0.291	0.175	0.110	0.247
JAOTNS010003484	F:GGACCACCGTAGTGTTCGTT R:TGCTCCACTGTGATTTCCA	A/T	124	0.300	0.323	0.200	0.579	0.269
JAOTNS010003033	F:ATGTCGACGCTCACACAGAC R:GTGTTGAAAAGTGGGGCGT	A/G	234	0.417	0.333	0.208	0.047*	0.275
JAOTNS010002888	F:GAGAGGATGACAGAGTCGCC R:AACGGGAGTTCAGAGTGGTG	T/C	243	0.333	0.280	0.167	0.132	0.239
JAOTNS010002678	F:ACCTCCTGAACCTCCATGTG R:TGGTACAAGCAAGCAACAC	A/G	233	0.467	0.361	0.233	0.021*	0.294
JAOTNS010002589	F:ATTCGGACATGCCCTAAGTG R:AGCCTCTCAGGCACCATCTA	T/C	172	0.367	0.504	0.383	0.034*	0.375
JAOTNS010002489	F:GGAGGAACAACAACAATGA R:GACCAACAGTGAAGGAACGC	A/G	200	0.383	0.352	0.225	0.479	0.288
JAOTNS010002366	F:TCAGAGTTCGATAGGTGCTGA R:AGCCTGCTCTCCGTGTCTA	A/G	310	0.383	0.333	0.208	0.230	0.275
JAOTNS010002188	F:AGTGTCTTGGCACCAAAAGG R:GAACCAAAAAGCTCCCTTCC	A/G	218	0.383	0.313	0.192	0.073	0.262
JAOTNS010002055	F:TTTCAAACGTTTCCACTCCC R:GTTTTGAGCGCTGTTGAGTG	A/G	221	0.383	0.313	0.192	0.073	0.262
JAOTNS010001952	F:TTTCAAACGTTTCCACTCCC R:GTTTTGAGCGCTGTTGAGTG	A/G	219	0.400	0.323	0.200	0.059	0.269
JAOTNS010001890	F:GCCTGAAAGTCATAGGTGGC R:GATCTCGCCTTCACTGGAG	T/C	283	0.400	0.323	0.200	0.059	0.269
JAOTNS010001555	F:AGCTGACAGGAGGGAACAGA R:TACTTGGGTCTGTAACCGGG	G/T	170	0.283	0.493	0.425	0.001*	0.369
JAOTNS010001422	F:GTGGAACATCGTGCATTG R:TCCACAATGAAAGCTGAGGA	T/C	157	0.383	0.370	0.242	0.770	0.299
JAOTNS010001200	F:CAGACAGGCTGCATCCAATA R:TACAGTGACCTAAGGCATCA	A/G	310	0.417	0.370	0.242	0.317	0.299
JAOTNS010001167	F:ACATCCATTACAAAACCGGC R:TTTAGATCAGGTTTCAAGCGGG	A/G	202	0.333	0.342	0.217	0.837	0.282
JAOTNS010000844	F:CTCACTGAGCCATCGGATT R:CGCCTTACTACCCACCTCTG	A/G	180	0.650	0.501	0.458	0.020*	0.373
JAOTNS010000619	F:TACAAATGACTGCGGGTCAA R:AGCCACATGGATCACCTTTC	A/G	183	0.383	0.333	0.208	0.230	0.275
JAOTNS010000510	F:GTGATGGACTGGGGTTTT R:CATCCGGAGAGAGACCGTAT	A/G	154	0.333	0.323	0.200	0.795	0.269
JAOTNS010000455	F:ATGATCATGTGGGTGGCTCT R:TTCAGACCGTTGTGCATCTG	A/G	202	0.367	0.323	0.200	0.282	0.269
JAOTNS010000388	F:CTGCGCTACATGCATTCT R:ATGGAACCCAGCCTAAATCC	A/T	137	0.383	0.370	0.242	0.770	0.299
JAOTNS010000204	F:GAGGTGATTTCTGCATGGGT R:ACGGCTTGGCAGTACAATCT	G/T	172	0.450	0.386	0.258	0.196	0.310
JAOTNS010000034	F:CTCACGAATGTGAACACGGT R:CGACACCTTGCCTCATTAT	A/T	194	0.417	0.333	0.208	0.250	0.275
JAOTNS010000005	F:GAAGCCCTGACTTTGATGGA R:TGCTAAAATGCCAGAGGAC	G/T	193	0.383	0.352	0.225	0.479	0.288
Mean±SD	/	/	195.710±48.614	0.394±0.071	0.362±0.067	0.246±0.083	/	0.292±0.040

Note: *Ho* represents the observed heterozygosity, *He* represents the expected heterozygosity, *MAF* represents minor allele frequency, *P<sub>HWE</sub>* represents the results for Hardy–Weinberg equilibrium, \**P*<0.05, *PIC* represents polymorphism information content.

corrected English grammar. Guoqing Duan designed, edited and corrected the manuscript. All authors read and approved the final manuscript.

### Conflict of Interest

The authors declare no conflict of interest.

### Acknowledgements

We would like to thank Anhui topmouth culter breeding centers for the helping of sample collection.

### References

- Chen, S.Q., Liu, Q.G., Ren, S.J., & Gong, X.L. (2009). Polymorphic microsatellite loci isolated from the topmouth culter (*Culter alburnus* Basilewsky). *Conservation Genetics Resources*, 1, 337. <https://doi.org/10.1007/s12686-009-9078-8>
- Chen, B., Cole, J.W., & Grond-Ginsbach, C. (2017). Departure from Hardy Weinberg Equilibrium and Genotyping Error. *Frontiers in Genetics*, 8, 167. <https://doi.org/10.3389/fgene.2017.00167>.
- D'Ambrosio, J., Phocas, F., Haffray, P., Bestin, A., Brard-Fudulea, S., Poncet, C., Quillet, E., Dechamp, N., Fraslin, C., Charles, M., Dupont-Nivet, & M. (2019). Genome-wide estimates of genetic diversity, inbreeding and effective size of experimental and commercial rainbow trout lines undergoing selective breeding. *Genetics Selection Evolution*, 51, 26. <https://doi.org/10.1186/s12711-019-0468-4>
- DePristo, M.A., Banks, E., Poplin, R., Garimella, K.V., Maguire, J.R., Hartl, C., et al. (2011). A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nature Genetics*, 43, 491-498. <https://doi.org/10.1038/ng.806>
- Jiang, H., Qian, Y., Zhang, Z., Meng, M., Deng, Y., Wang, G., He, S., & Yang, L. (2023). Chromosome-level genome assembly and whole-genome resequencing of topmouth culter (*Culter alburnus*) provide insights into the intraspecific variation of its semi-buoyant and adhesive eggs. *Molecular Ecology Resources*, 23, 1841-1852. <https://doi.org/10.1111/1755-0998.13845>
- Kalinowski, S.T., Taper, M.L., & Marshall, T.C. (2007). Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology*, 16(5), 1099-1106. <https://doi.org/10.1111/j.1365294X.2007.03089.x>
- Li, T., Jia, Y., Gao, J., Zhao, J., Jiang, W., Gu, Z., & Yang, R. (2018). Optimization of shock conditions to induce gynogenesis in topmouth culter (*Culter alburnus*). *Aquaculture*, 490, 344-349. <https://doi.org/10.1016/j.aquaculture.2018.03.005>
- Liu, Y., & Yang, J. (2013). The complete mitochondrial genome sequence of Xingkai topmouth culter (*Culter alburnus*). *Mitochondrial DNA*, 25(6), 451-453. <https://doi.org/10.3109/19401736.2013.809430>
- Qi, P.Z., Qin, J.H., & Xie, C.X. (2015). Determination of genetic diversity of wild and cultured topmouth culter (*Culter alburnus*) inhabiting China using mitochondrial DNA and microsatellites. *Biochemical Systematics and Ecology*, 61, 232-239. <https://doi.org/10.1016/j.bse.2015.06.023>
- Qi, P.Z., Xie, C.X., Guo, B.Y., Wu, C.W., Lu, S.M., & Duan, Y.J. (2013). Development of new polymorphic microsatellite markers in topmouth culter (*Culter alburnus*) and determination of their applicability in *Culter mongolicus*. *Genetics and Molecular Research*, 12(2), 1761-1765. <http://dx.doi.org/10.4238/2013.May.21.7>
- Shan, S., Kan, X., Gong, J., Shao, C., & Wang, Y. (2022). Development and characterization of 36 SNP markers for *Hynobius yiwuensis*. *Conservation Genetic Resources*, 14, 15-18. <https://doi.org/10.1007/s12686-021-01233-4>
- Sun, N., Zhu, D., Li, Q., Wang, G., Chen, J., Zheng, F., Li, P., & Sun, Y. (2021). Genetic diversity analysis of Topmouth Culter (*Culter alburnus*) based on microsatellites and D-loop sequences. *Environmental Biology of Fishes*, 104, 213-228. <https://doi.org/10.1007/s10641-021-01062-2>
- Vignal, A., Milan, D., SanCristobal, M., & Eggen, A. (2002). A review on SNP and other types of molecular markers and their use in animal genetics. *Genetics Selection Evolution*, 34, 275-305. <https://doi.org/10.1186/1297-9686-34-3-275>
- Wang, W., Chen, L., Yang, P., Hou, L., He, C., Gu, Z., & Liu, Z. (2007). Assessing genetic diversity of populations of topmouth culter (*Culter alburnus*) in China using AFLP markers. *Biochemical Systematics and Ecology*, 35, 662-669. <https://doi.org/10.1016/j.bse.2007.04.008>
- Zhang, L., Sun, G., Azhanhan, E., Li, Y., Shi, W., Zou, D., Zhang, H., Ding, J., Kessler, M., Wang, M., & Yang, W. (2024). Genetic diversity assessment for reintroduced Przewalski's horse (*Equus ferus*) based on newly developed SSR markers. *Conservation Genetic Resources*, 16, 89-101. <https://doi.org/10.1007/s12686-023-01330-6>
- Zhou, H., Duan, G., Pan, T., Wang, H., Jiang, H., Ling, J., Hu, Y., & Yang, M. (2022). Development a Set of 46 SNP Markers for the Yellow Catfish (*Tachysurus fulvidraco*). *Turkish Journal of Fisheries and Aquatic Sciences*, 22(12), TRJFAS20976. <https://doi.org/10.4194/TRJFAS20976>