

Effects of a Myrrh (*Commiphora myrrha*) Essential Oil Supplemented Diet on Haemato-Biochemical Parameters, Expression of Tissue-Specific Immune- and Stress-Related Genes, and Resistance of *Cyprinus carpio* to *Aeromonas hydrophila* Infection

Ümit Acar¹, Önder Yıldırım^{2,*}, Esin Baba², Fahriye Zemheri Navruz³, Mustafa Hacisa²,
Sevdan Yılmaz⁴

¹Çanakkale Onsekiz Mart University, Bayramiç Vocational School, Department of Forestry, Çanakkale, Türkiye.

²Muğla Sıtkı Koçman University, Faculty of Fisheries, Department of Aquaculture, 48000 Muğla, Türkiye.

³Bartın University, Faculty of Science, Department of Molecular Biology and Genetics, Türkiye.

⁴Çanakkale Onsekiz Mart University, Faculty of Marine Science and Technology, Department of Aquaculture, Çanakkale, Türkiye.

How to Cite

Acar, Ü., Yıldırım, Ö., Baba, E., Zemheri Navruz, F., Hacisa, M., Yılmaz, S. (2025). Effects of a Myrrh (*Commiphora myrrha*) Essential Oil Supplemented Diet on Haemato-Biochemical Parameters, Expression of Tissue-Specific Immune- and Stress-Related Genes, and Resistance of *Cyprinus carpio* to *Aeromonas hydrophila* Infection. *Turkish Journal of Fisheries and Aquatic Sciences*, 25(8), TRJFAS27525. <https://doi.org/10.4194/TRJFAS27525>

Article History

Received 13 December 2024

Accepted 20 February 2025

First Online 26 February 2025

Corresponding Author

E-mail: onderyildirim@mu.edu.tr

Keywords

Myrrh essential oil
Hematological parameters mRNA
expression
Common carp
Immunomodulatory effects

Abstract

Plant-based additives or botanicals, have garnered considerable interest in the aquaculture industry for their multifaceted benefits in fish diets. In this study, a 30 days short term feeding experiment was designed to determine the effects of Myrrh (*Commiphora myrrha*) essential oil on hematological, biochemical parameters and tissue specific mRNA expression responses of common carp (*Cyprinus carpio*). Fish were experimentally infected with *Aeromonas hydrophila* and monitored for 20 days after been divided in four groups: a control group without *C. myrrha* and three experimental groups supplemented with 0.25%, 0.50%, and 1.0% *C. myrrha* essential oil (CMO 0.25, CMO 0.50, and CMO 1.0, respectively). Results revealed a significant increase in RBC count, hemoglobin, and hematocrit values compared to the control diet. Positive effects of *C. myrrha* supplementation have been observed also in serum biochemistry parameters. mRNA transcripts of *il-1 β* , *tnf- α* , *il-8*, *sod*, *cat*, *gpx* and *gst* genes significantly increased in spleen and liver tissue of fish fed with *C. myrrha* supplemented diets. After the challenge with *Aeromonas hydrophila*, the survival rates were 60%, and 83.33%, respectively, suggesting enhanced fish resistance in the CMO 0.50 and CMO 1.0 groups compared to the control. The results suggest the immunomodulatory roles of *C. myrrha* essential oil.

Introduction

Driven by increasing global demand for aquafood and sustainable food production, aquaculture has rapidly expanded as a key agricultural sector. Among the widely cultivated freshwater species, the common carp (*Cyprinus carpio*) is particularly valued for its adaptability to diverse climatic conditions and its significant contribution to the industry (Eddy and Underhill, 1974). In addition, common carp is the third most frequently introduced species globally and the most commercially significant freshwater fish species in the world (Saika and Das, 2009). Nonetheless, the rapid growth of common carp aquaculture has resulted in the emergence of numerous diseases and health

complications within this culture. Bacterial infections, in particular "motile aeromonas septicemia" caused by *Aeromonas hydrophila*, have significantly affected the culture of common carp in river aquaculture and resulted in significant mortality as well as financial losses for fish farmers (Austin and Adams, 1996). *A. hydrophila* is a facultatively anaerobic, motile, gram-negative, oxidase-positive bacilli or coccobacilli rod (Dias *et al.*, 2012; Semwall *et al.*, 2023). In fish, it is an opportunistic pathogen and causes hemorrhagic septicemia, exophthalmia, infectious abnormal dropsy, fin, hemorrhagic septicemia, exophthalmia, infectious abnormal dropsy, and fin and tail rot (Austin and Austin, 2007; Zhang *et al.*, 2020).

Antibiotics are extensively used to control bacterial diseases in aquaculture. Nevertheless, their long-term use leads to the rise of antibiotic-resistant bacteria and other negative consequences such as the permanence of antibiotic residues in both fish and the environment (Cabello, 2006). Thus, the development of innovative methods for disease control in common carp aquaculture is critically needed.

Essential oils (EOs) may have the potential to be used as an alternative to antibiotics in aquaculture (Romero *et al.*, 2012; Cunha *et al.*, 2018). EOs obtained from plant raw materials are lipophilic, volatile, fragrant, liquid natural multicomponent systems (Edris, 2007) obtained from the flowers, leaves, seeds, roots, peel, wood, herbs, and fruits parts of the plant. EOs are mixtures of low molecular weight substances with a wide variation in their chemical properties (Hussain *et al.*, 2008). More than 3,000 different chemicals with a wide variety of chemical structures have been identified in EOs (Cunha *et al.*, 2018). EOs are generally classified as terpene hydrocarbons, simple and terpene alcohols, aldehydes, phenols, ketones, esters, ethers, peroxides, oxides, organic acids, furans, lactones, coumarins (Swamy *et al.*, 2016). EO compounds have a wide spectrum of inhibitors to the growth of Gram-negative and Gram-positive bacteria (Remmal *et al.*, 1993). Some studies have determined that the mechanism of action of EOs depends on their major functional groups. Due to their antimicrobial properties, these oils have the ability to constitute alternative prophylactic and therapeutic agents in aquaculture (Romero *et al.*, 2012). Previous research has highlighted the benefits of herbal medicine, citing its potential to replace antibiotics in the treatment of disease, availability locally, biodegradability, and lack of adverse effects. Nowadays, the effect of essential oils or extracts on aquaculture pathogens has been reported in some studies (Wu *et al.*, 2010; Wu *et al.*, 2013; Dotta *et al.*, 2014; Tang *et al.*, 2014; Zahran *et al.*, 2014; Gabriel *et al.*, 2015; Acar *et al.*, 2015; Baba *et al.*, 2016; Hashimoto *et al.*, 2016; Fazio *et al.*, 2022; Habib *et al.*, 2024; Naz *et al.*, 2024; Ujan *et al.*, 2025). There are various studies showing that different essential oils in carp, such as *Zataria multiflora* (Soltani *et al.*, 2010), blue gum (Sheikhzadeh *et al.*, 2009), oregano (Abdel-Latif *et al.*, 2020), *Thymus vulgaris* (Ghafarifarsani *et al.*, 2022), increase the immune system and disease resistance. In many studies where plant essential oils were used as feed additives, it was determined that the application produced successful results in cultivation. A number of beneficial effects of essential oils were documented by these earlier researchers, including improved growth performance, feed utilization, and disease resistance; they also strengthened the immune system, prevented stress, and decreased mortality. No studies have examined the use and effects of myrrh (*Commiphora myrrha*) essential oil in fish feed.

Myrrh is an important medicinal plant of the Burseraceae family. It has more than 150 species of

plant spread across subtropical and tropical regions, particularly north-east Africa, southern Arabia, and India (Demissew, 1993). Myrrh and myrrh derivatives have long been used as a raw material in the traditional medicines of India, China, Rome, Greece and Babylon (Shen and Lou, 2008). Myrrh is derived from an Arabic term that translates to 'bitter.' One especially valued type of myrrh is referred to as *karam* or Turkish myrrh (Encyclopedia, 2024). Myrrh is composed of several components, which consist of volatile oil ranging from 2 to 8%, resin making up 23 to 40%, gum accounting for 40 to 60%, and bitter compounds that comprise 10 to 25% (Chen *et al.*, 2013).

Myrrh has been used traditionally for the treatment of many diseases, such as mouth ulcers, aches, stomach disorders, fractures, inflammatory diseases, and microbial infections since ancient times (Su *et al.*, 2011; Abdelsalam *et al.*, 2022). *C. myrrha* is utilized for its antiseptic, astringent, anthelmintic, carminative, emmenagogue, and expectorant properties. The primary components of myrrh are terpenoids (monoterpenoids, sesquiterpenoids, and volatile/essential oils), diterpenoids, triterpenoids, and steroids (Batiha *et al.*, 2023). Studies have illustrated its diverse biological activities, including anti-inflammatory, antioxidant, antimicrobial, anti-viral, neuroprotective, antidiabetic, anticancer, analgesic, and antiparasitic properties (Gadir and Ahmet., 2014; Madia *et al.*, 2021; Abdelsalam *et al.*, 2022; Batiha *et al.*, 2023) immune response (Noyal *et al.*, 2018), hematology-serum biochemistry (Mitsumoto *et al.*, 2021) and leukocytes proliferation (Haffor, 2009).

Considering the introduction presented above, this study is specifically designed to investigate the effects of myrrh essential oil on the health components of carp, including the expression levels of immune and antioxidant genes, hematological and serum biochemical responses, and resistance to *A. hydrophila*.

Materials and Methods

This study was approved by the Animal Ethics Committee of Muğla Sıtkı Koçman University (Muğla,

Türkiye, Approval Number: 2024/02-2). Myrrh essential oil was chosen due to its valued for its antimicrobial, anti-inflammatory, and antioxidant properties, supporting immune function and overall health. Myrrh essential oil was purchased from a local herbal medicine store in İzmir, Turkey. Four diets with different concentrations of CMO (0.25%, 0.50%, and 1%) along with a control diet without CMO were prepared. The details of the dietary formulations and ingredient compositions are provided in Table 1.

Crude ash (%), crude protein (CP; %), and crude lipid (CL; %) of the test diets were determined using the AOAC procedures. The experimental fish were supplied by the Central Fisheries Research Institute in Antalya, Türkiye. Upon arrival at the laboratory, 180 clinically healthy juvenile common carp weighing 14.77 ± 0.21 g

were given a 14-day adaptation period before the start of the feeding experiment. Subsequently, they were randomly divided into 12 experimental glass tanks with triplicate (15 fish/tank), with an equal number of fish in each tank. The feeding experiment was carried out at a temperature of $22\pm 0.5^\circ\text{C}$, following a 12-hour light to 12-hour dark photoperiod. Throughout the feeding experiment, water conditions were maintained at a pH of 7.6 ± 0.1 and a dissolved oxygen level above 5.5 mg/L. The fish were fed twice daily, in the morning and afternoon, with the amount of food given being 2% of their body weight per day. The experimental diets were administered to the fish for a period of 30 days (Ghelichpour et al., 2021). No mortality was recorded during the course of the feeding experiment. The phytonutrients found in the CMO supplement were investigated using the GC-MS analysis method. Chromatograms of the CAO were characterized and then identified after matching with the spectra of the Wiley W9N11 library.

At the ending of the experimental period, the fish were not fed for 24 hours. 15 fish from each experimental group (5 fish from each aquarium) were randomly selected and anesthetized with 50 $\mu\text{L/L}$ of clove oil to collect blood samples (Woody et al., 2002). In order to prevent contamination, the caudal region of the fish was meticulously cleaned with 70% ethanol. Syringes were employed to capture blood samples in the subsequent phase. Following this, the samples were divided into two categories for analysis: lavender-top tubes with anticoagulant (K3EDTA) for hematological analysis and red-top (SST™ II) advance serum separator tubes for biochemical analysis. Finally, the serum separator containers were centrifuged at 4000 g for 10 minutes to separate the serum for biochemical analysis. Serum samples were subsequently stored at -80°C for subsequent analysis. Liver and splenic tissues were extracted from the fish subsequent to blood collection. Until the analysis, the samples were maintained at -80°C .

Hemoglobin (HGB) content was measured using the cyanmethemoglobin technique after observing red blood cells (RBCs) under a light microscope. The hematocrit (HCT) measurement was conducted by employing a capillary HCT tube subsequent to centrifugation per the procedures outlined by (Hesser, 1960). Commercial test kits from the Bioanalytic Diagnostic Industry in Germany were used for biochemical analysis. The spectrophotometer (OptizenPOP UV/VIS) was utilized to measure various parameters including glucose (GLU), total protein (TPROT), cholesterol (CHOL), albumin (ALB), globulin (GLO), triglyceride (TRIG), alkaline phosphatase (ALP), lactate dehydrogenase (LDH), glutamic pyruvic transaminase (GPT), and glutamic-oxaloacetic transaminase (GOT) (Yilmaz and Ergün, 2012).

Thirty fish from both the control and experimental groups were used to investigate the resistance of the common carp to *A. hydrophila*. Following a period of 30 days during which the fish were given food and had their blood samples taken, they were then injected with 0.1 mL of a suspension containing 1.5×10^6 colony-forming units per milliliter of *A. hydrophila*, using the intraperitoneal route. The suspension was prepared in phosphate buffered saline. The fish were constantly examined for any obvious symptoms of sickness, such as odd behavior, and deceased fish were carefully removed from the tanks to minimize stress. Death occurred in all of the groups within 20 days after infection. The infection was confirmed by re-isolating the germs from the deceased fish. Bacterial identification was achieved by using traditional biochemical methods (Austin and Austin, 2007) and the API 20 Strep kit (Biomerieux, France).

The liver and spleen tissues were collected from 6 fish per group, with 2 fish from each tank, for the purpose of RNA isolation. The GeneJET RNA Purification Kit, made by Thermo Fisher Scientific in the United States, was used in our study. Quantification of RNA and study of RNA integrity was performed using a Multiskan

Table 1. Formulations and chemical composition of experimental diets.

Ingredient	Control	CMO _{0.25}	CMO _{0.50}	CMO _{1.0}
Fish meal	25.00	25.00	25.00	25.00
Soybean meal	35.00	35.00	35.00	35.00
Wheat meal	12.00	12.00	12.00	12.00
Fish oil	5.00	5.00	5.00	5.00
Mineral/vitamin premix ^{a,b}	4.00	4.00	4.00	4.00
Starch	19.00	18.75	18.50	18.00
Myrrh essential oil	0	0.25	0.50	1.00
Chemical composition				
Protein	36.62	36.84	36.76	36.19
Lipid	7.51	7.34	7.56	7.23
Ash	5.49	5.60	5.47	5.72

^aVitamin Mix: Vit. A, 18000 IU; Vit. D3, 2500 IU; Vit. E, 250 mg/kg; Vit. K3, 12 mg/kg; Vit. B1, 25 mg; Vit. B2, 50 mg; Vit. B3, 270 mg; Vit. B6, 20 mg; Vit. B12, 0.06 mg; Vit. C, 200 mg; Folic acid, 10 mg; Calcium d-pantothenate, 50 mg; Biotin, 1 mg; Inositol, 120 mg; Choline chloride, 2000 mg.
^bMineral Mix: Fe, 75.3 mg; Cu, 12.2 mg; Mn, 206 mg; Zn, 85 mg; I, 3 mg; Se, 0.350 mg; Co, 1 mg

Go instrument manufactured by Thermo Fisher Scientific in the United States. The purified RNA was mixed with RNase-free DNase I (Thermo Fisher Scientific, USA) and then analyzed using Real-time PCR. Specific primer pairs were used to amplify SOD, CAT, GpX, GST, IL-1 β , IL-8, TNF- α , and β actin. The cDNA synthesis process included reverse-transcription of 1 μ g of total RNA from each sample using the RevertAid™ H Minus First Strand cDNA Synthesis Kit (Thermo Fisher Scientific, USA). The cDNA was stored at a temperature of -20°C to guarantee its long-term preservation. The FastPCR 6.0 program (Kalendar et al., 2009) was used to generate primers (Table 2) that were meant to cover at least one exon boundary.

The data were reported as mean values with standard deviation (SD) and analyzed using SPSS 17.0 software (SPSS Inc., Chicago, USA). The study used one-way analysis of variance (ANOVA) followed by the Tukey's test to see whether there were significant differences in the measured parameters across the treatments. The analyses were conducted with a significance threshold of 0.05.

Results

Bioactive Components of CMO

The primary functional bioactive components in CMO used in the current investigation were also assessed. The GC-MS chromatograms showed 46 peaks representing different metabolites, as shown in Table 3.

The phyto-ingredients that were found in the highest quantities were cis-Ocimene (24.49%), Eucalyptol (34.77%), linalool (13.83%), alpha-Terpineol (4.16%), and Linalyl acetate (5.87%) (Table 3).

Hematological and Serum Biochemical Parameters

Based on the results reported in Table 4, the inclusion of dietary CMO at a dose range of 0.25-1% led to a substantial and measurable increase ($P < 0.05$) in the red blood cell count, Hgb concentration, and Hct value of common carp.

Significant changes were seen in the assessed biochemical variables across all experimental groups. When comparing the control group to the CMO1.0 group, it was noticed that the CMO1.0 group had the highest TPROT levels ($P < 0.05$). However, the TPROT values in other groups did not exhibit a statistically significant difference compared to the controls ($P > 0.05$). The control group exhibited a substantial ($P < 0.05$) rise in CHOL and LDH levels compared to the groups supplemented with CMO. No significant variations were seen in the other serum biochemical measures, including GLU, TRIG, ALP, and AST levels, between the experimental groups ($P > 0.05$).

Gene Expression Profile

Dietary supplementation with CMO over a period of 30 days elicited notable changes in gene expression levels within the spleen and liver tissues of common carp, particularly affecting genes associated with anti-oxidative responses and immune function (Figure 1).

Specifically, both spleen and liver tissues exhibited a significant upregulation in the mRNA expression levels of SOD, CAT and GpX in groups supplemented with CMO at concentrations of 0.5% and 1%. Concurrently, there was a similar upregulation observed in pro-inflammatory cytokines such as IL-1 β , IL-8, and TNF- α within both spleen and liver tissues, compared to the control group.

Table 2. Primer nucleotide sequences used to conduct this research.

Gene		Primer sequence	Product size (bp)	GenBank No
β -Actin	F	CTGGTATCGTGATGGACTCT	204	M24113
	R	CAGAGCTTCTCCTTGATGTC		
SOD	F	GATAGTGACAGACACGTCGGGA	180	XM_019111527.1
	R	AAGACTTTCGTCATTGCCTCC		
CAT	F	GAGCACGTAGGGTTCAAGTGC	162	NC_031745.1
	R	TGCGCTTATCTCTGTCTGCCA		
GpX	F	TGTCCTTGATGGGTGATCCCA	144	GQ376155.1
	R	CAATGTCGCTGGTGAGGAACC		
GST	F	CCTCGCTGAAAGAGCTTCAC	138	XM_019117245.1
	R	ATACTGGGACGGTCTTCAGC		
TNF- α	F	GTGTCTACAGAAACCCTGGA	109	AJ311800
	R	AGTAAATGCCGTCAGTAGGA		
IL-1 β	F	TTACAGTAAGACCAGCCTGA	89	AJ245635
	R	AGGCTCGTCACTAGTTTGT		
IL-8	F	GTCTTAGAGGACTGGGTGTA	120	AB470924.1
	R	ACAGTGTGAGCTTGGAGGGA		

*The real-time PCR was performed using the Maxima SYBR Green/ROX qPCR Master Mix (Thermo Fisher Scientific, USA) according to the manufacturer's instructions. The cDNA used was 1 μ g. The thermal cycling conditions consisted of 95°C for 10 minutes, followed by 40 cycles of 95°C for 15 seconds, 55 cycles of 95°C for 30 seconds, and an extension step at 72°C for 10 minutes. The target genes included β -actin, SOD, CAT, GpX, GST, IL-1 β , IL-8, and TNF- α . Electrophoresis was performed on the PCR and RNA product using a 2% w/v agarose gel electrophoresis technique, using a 2 μ g/ml SYBR Safe DNA Gel Stain (Thermo Fisher Scientific, USA). The length of PCR and RNA products was assessed by administering and checking a DNA ladder Gene Ruler 50 bp Marker Plus (Thermo Fisher Scientific, USA) alongside the samples. The findings were assessed using real-time PCR using the CFX Connect™ RealTime PCR Detection System, manufactured in the United States.

Table 3. Chemical composition of *Commiphora myrrha* essential oil.

Peak#	R.Time	Area%	Name
1	6.020	0.01	4-Octen-3-one (CAS)
2	8.985	0.16	Tricyclene
3	9.242	0.59	Thujene <alpha->
4	9.510	24.49	cis-Ocimene
5	10.012	0.55	Camphene
6	11.046	0.08	Sabinene
7	11.126	1.01	Pinene <beta->
8	11.820	0.44	Myrcene
9	12.051	0.47	Butanoate <butyl->
10	12.272	0.14	Phellandrene <alpha->
11	12.499	0.63	DELTA.3-Carene
12	12.774	0.14	Terpinene <alpha->
13	13.107	1.42	Cymene <para->
14	13.392	34.77	Eucalyptol
15	13.708	0.15	trans-.beta.-Ocimene
16	14.479	0.28	.gamma.-Terpinene
17	15.048	0.23	Linalool oxide <trans->
18	15.670	0.31	Linalool oxide <cis->
19	16.010	0.04	alpha.-Pinene oxide
20	16.237	13.83	Linalool
21	16.385	0.06	Limonene oxide <cis->
22	16.679	0.04	Oct-1-en-3-yl acetate
23	17.496	0.16	Dihydrolinalool
24	17.811	0.92	Camphor
25	17.901	0.16	BETA. TERPINEOL
26	18.211	0.04	Cyclohexanone, 5-methyl-2-(1-methylethyl)-, (2R-cis)- (CAS)
27	18.347	0.29	Isoborneol
28	18.635	0.07	p-Menth-8-en-1-ol, stereoisomer
29	18.707	0.32	Bicyclo[2.2.1]heptan-2-ol, 1,7,7-trimethyl-, (1S-endo)-
30	18.790	0.06	Lavandulol
31	19.146	0.68	3-Cyclohexen-1-ol, 4-methyl-1-(1-methylethyl)-, (R)-
32	19.494	0.04	Benzenemethanol, 4-(1-methylethyl)- (CAS)
33	19.693	4.16	.alpha.-Terpineol
34	19.942	0.86	Terpineol <gamma->
35	21.092	0.24	Nerol (CAS)
36	22.102	5.87	Linalyl acetate
37	23.173	0.12	Anisole <para-allyl->
38	23.363	0.24	Lavandulyl acetate
39	25.423	1.30	Terpinyl acetate <alpha->
40	25.490	0.16	Terpineol <gamma->
41	25.926	0.59	Neryl acetate
42	26.586	1.89	Geranyl acetate
43	27.318	0.42	Benzene, 1,2-dimethoxy-4-(2-propenyl)- (CAS)
44	27.838	0.72	Caryophyllene
45	28.969	0.67	alpha.-Humulene (CAS)
46	33.097	0.20	Caryophyllene oxide
		100.00	

Table 4. Hematological and Serum Biochemical Parameters of fish fed with experimental diets

Parameter	Control	CMO _{0.25}	CMO _{0.50}	CMO _{1.0}
RBC (×10 ⁶ per mm ³)	1.93±0.05 ^b	2.09±0.07 ^b	2.73±0.16 ^a	2.68±0.19 ^a
HGB (g/dL)	9.38±0.16 ^d	10.55±0.42 ^c	11.65±0.49 ^b	13.31±0.41 ^a
HCT (%)	26.14±0.62 ^c	28.00±0.55 ^c	30.48±0.70 ^b	33.17±1.87 ^a
TPROT(g/dL)	9.80±1.95 ^b	9.83±0.96 ^b	10.98±3.09 ^{ab}	13.27±1.39 ^a
GLU(mg/dL)	49.83±9.15 ^a	49.71±11.64 ^a	49.40±9.55 ^a	50.54±4.43 ^a
CHOL(mg/dL)	50.27±13.02 ^a	35.67±6.93 ^b	25.33±6.19 ^b	31.60±5.67 ^b
TRIG(mg/dL)	21.19±5.49 ^a	23.14±6.50 ^a	23.14±7.46 ^a	21.86±5.14 ^a
ALP (U L ⁻¹)	45.00±8.01 ^a	28.63±4.17 ^b	27.37±9.44 ^b	51.00±4.81 ^a
AST (U L ⁻¹)	47.82±12.77 ^a	35.11±6.93 ^a	40.83±2.20 ^a	40.26±8.37 ^a
LDH (U L ⁻¹)	165.5±44.5 ^a	113.8±29.9 ^{ab}	130.3±25.6 ^{ab}	109.03±23.67 ^b

*The data is shown as the mean value plus or minus the standard deviation, with a sample size of 15. Values with distinct letters in the same row are deemed statistically significant within groups (P<0.05). Acronyms: RBC is for red blood cells, HGB stands for hemoglobin, HCT stands for hematocrit, TPROT stands for total protein, GLU stands for glucose, and CHOL stands for cholesterol. TRIG: Triglycerides; AST stands for aspartate aminotransferase, ALT stands for alanine aminotransferase, and LDH stands for lactate dehydrogenase.

Bacterial Pathogen Challenge

Common carp’s resistance to *Aeromonas hydrophila* after a 20-day observation period, survival rates recorded in control, CMO0.25, CMO0.50, and CMO1 groups 50, 43.33, 60, and 83.33%, respectively (Table 5).

According to Kaplan–Meier survivorship curves, it was noticed that the highest survival rates were found in CMO1 (P<0.05; Figure 2).

Discussion

Limitation of the use of antibiotics is crucial to prevent the proliferation of antibiotic-resistant bacteria and maintain the health of aquatic environments in aquaculture. For this purpose, incorporating natural additives can be an effective strategy for improving the well-being and resilience of cultured fish (Mathew *et al.*, 2022). Reducing dependence on antibiotics and

embracing natural remedies such as herbal extracts and essential oils is in line with sustainable aquaculture practices. These additives not only stimulate appetite, promote growth, and enhance immune responses but also provide a safer alternative to synthetic chemicals. Furthermore, focusing on the immunostimulatory and antioxidant properties of essential oils derived from medicinal plants can contribute to healthier aquatic ecosystems and more resilient fish populations (Zhang *et al.*, 2020; Farag *et al.*, 2023). This study assesses, for the first time, the impact of adding *Commiphora myrrha* essential oil to the diet on physiological reactions, antioxidant levels, enhancement of immune function, and increased resilience of common carp against *Aeromonas hydrophila* infections.

Monitoring parameters such as the count of red blood cells (RBC), levels of hemoglobin (Hb), and levels of hematocrit (Hct) may offer important information about the health of fish, including their ability to transport oxygen and the possibility of hemolytic




Figure 1. Gene expression responses in spleen (A-C) and liver (B-D) tissue of common carp fed diets supplemented with different levels of CMO. Values are presented as means ± S.D. (n=6) (P<0.05).

Table 5. Mortality rate (MR; %), survival rate (SR; %), and relative percent survival (RPS) of common carp fed with diets containing different levels of CMO for 30 days and then experimentally challenged with *A. hydrophila* infection and observed for 20 days.

Experimental groups	Number of challenged fish	Mortality rate (MR;%)	Survival rate (SR;%)	RPS
Control	30	50.0 ^a	50.0 ^c	-
CMO _{0.25}	30	56.66 ^a	43.33 ^c	-13.33
CMO _{0.50}	30	40.0 ^b	60.0 ^b	20.0
CMO ₁	30	16.66 ^c	83.33 ^a	66.66

*Values with different superscript letters in the same column are considered statistically different within groups (P<0.05). MR (%) = (No. of fish died after challenge / Total No. of challenged fish) × 100. RPS (%) = [1 – (Mortalities of CMO-supplied groups / Mortalities of Control group)] × 100.

processes (Clauss *et al.*, 2008; Fazio *et al.*, 2013). Examination of qualitative and quantitative variations in these parameters may serve as indicators for different illnesses, such as anemia or dehydration, and assist the researchers in accurately diagnosing and treating fish. Moreover, these characteristics aid in assessing the efficacy of the fish's hematological system and its capacity to facilitate oxygen transportation and eliminate carbon dioxide, which are crucial functions for survival and general well-being (Abdel-Tawwab *et al.*, 2006). In the present study, RBC count, Hct, and Hbg values were significantly increased in all groups of fish fed with CMO-supplemented diets compared to the control group. The findings indicate that adding CMO to common carp fish's diet effectively decreases hemolysis. This can be due to the enhancement of antioxidant capacity, which safeguards red blood cells from hemolysis (Gaudet *et al.*, 1975). Nevertheless, eucalyptol, the main constituent of CMO used in the current study, seems to alleviate such harm, resulting in a significant enhancement in hematology when compared to the control group. The potential reason for this protective effect may be the reduction of oxidative stress as for previous studies reporting that eucalyptol has potent inhibitory effects on pro-oxidant-induced oxidative stress [Cho, 2012]. As far as we know, this is the first documentation of CMO in diets for common carp. Consistent with these results, gibel carp (*Carassius gibelio*) that were given diets enhanced with thyme essential oil showed improvements in hematological parameters, including higher levels of red blood cells (RBCs), hematocrit (Hct), and hemoglobin (Hb) [(Zadmajid *et al.*, 2017). Moreover, research has shown

that adding herbal supplements to the diet might improve hematological parameters in different species of teleost fish (Acar *et al.*, 2015; Rashidian *et al.*, 2020).

Serum biochemical parameters are essential biological markers that aid in evaluating the impact of feed additives on the health of fish. These metrics provide essential data on the metabolic processes, organ function, and general physiological condition of fish [Harikrishnan *et al.*, 2012]. This study focused on analyzing many important serum biochemical markers, including TPROT, Glu, Chol, Trig, ALP, AST, and LDH. TPROT has a role in the health of fish (Magnadóttir, 2006; Yilmaz *et al.*, 2024) and indicates the nutritional and health state of fish (El-Houseiny *et al.*, 2022). The findings of our research show that the CMO1.0 group had the highest levels of TPROT compared to the control group. In another study, higher blood protein levels were associated with improved fish survival rates and enhanced immunological effectiveness (Simajuntak *et al.*, 2018). These results were in line with the findings of (Souza *et al.*, 2020), who discovered that adding Cochin grass (*Cymbopogon flexuosus*) EO to diets might raise TPROT levels in the blood of Nile tilapia (*Oreochromis niloticus*). Furthermore, addition of 2 ml of Panax ginseng essential oil to the diets of Nile tilapia was reported to significantly increase the TPROT level (Yilmaz *et al.*, 2024). In this study, GLU, note to be a stress indicator in fish (Martínez-Porchas *et al.*, 2009), did not varied between all treatment groups. In contrast, other studies have reported that dietary supplementation with herbal additives can reduce serum GLU levels in fish (Ghafarifarsani *et al.*, 2022; Adeli *et al.*, 2021). However, based on the current




Figure 2. Kaplan–Meier survivorship curves (cumulative survival [%] over time [h]) for common carp after challenge with *Aeromonas hydrophila*; the fish were fed with CMO supplemented diets.

results, it can be concluded that supplementation with CMO did not induce any stress in the fish. This observation suggests that the inclusion of CMO in the diet did not lead to an increase in stress levels as reflected by serum GLU levels. CHOL is essential for the structure of biomembranes, especially in the formation of the outer layer of serum lipoproteins (Dorojan *et al.*, 2015). In addition, CHOL acts as a precursor for the production of steroid hormones (Pourmozaffar *et al.*, 2019), which play a crucial role in numerous physiological processes in organisms. However, TRIG operates largely as an energy storage form in cells and serves as the major source of metabolic and cellular energy (Pourmozaffar *et al.*, 2019). They play a crucial role in energy metabolism and are activated when the body needs more energy. We noticed a substantial reduction in cholesterol levels in those fed with a meal supplemented with CMO and no significant changes in TRIG. Several studies have proven the efficacy of herbal oils and essential oils in reducing cholesterol levels. Similarly, Monterey cypress (*Cupressus macrocarpa* Hartw) leaf essential oil was shown to lower blood cholesterol levels in common carp (Kesbiç *et al.*, 2020). Also, ginger oil was proven to have a similar effect on cholesterol concentrations in the same fish species (Immanuel *et al.*, 2009). Some researchers reported that these lower levels of cholesterol in the blood may be attributed to the presence of phytosterols (Rinchar *et al.*, 2003). Additionally, the suppression of cholesterol production by essential oils derived from plants may also contribute to the decrease in cholesterol levels (Ngugi *et al.*, 2017). Monitoring AST and ALP levels in fish serum is crucial for assessing liver health and detecting potential liver problems, especially when fish are exposed to external stressors or environmental challenges (Yilmaz *et al.*, 2024). Changes in these enzyme levels can provide valuable insights into the impact of stress, diseases, or dietary factors on liver function and overall fish health. Elevated levels of these enzymes in fish serum can indicate liver damage or dysfunction, as these enzymes are primarily found in hepatocytes (liver cells) and are released into circulation when hepatocytes are damaged (Ghelichpour *et al.*, 2021). The current findings suggest hepatoprotective properties of CMO, which may be attributed to its enhanced antioxidant capacity in the liver of the fish. In the study conducted by (Shourbela *et al.*, 2021), Nile tilapia fed with diets containing oregano essential oil did not exhibit any signs of toxicity, stress, inflammatory diseases, or other undesirable effects. Aquaculture researchers are particularly interested in the ability of fish to withstand damaging stress, especially in aquaculture settings. One possible method to mitigate this stress is to use herbal sources that are abundant in physiologically active compounds (Khalil *et al.*, 2022).

The enzymatic antioxidant protectors such as superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPX), and glutathione S-transferase (GST) play a significant role in the detoxification process and

in counteracting the adverse effects of reactive oxygen species (ROS) and free radicals (Oberley *et al.*, 1988). These enzymes work synergistically to mitigate oxidative stress by reducing the accumulation of harmful oxidants. In this study, the increase in Myrrh essential oil dosage in the experimental diets led to enhanced expression of antioxidant enzymes such as SOD, CAT, GPX, and GST in the spleen and liver of common carp. This indicates that higher doses of the oil strengthened the fish's defense mechanisms against stress and/or pathogens. The results obtained in the present study may be associated with myrrh essential oils' antioxidant abilities. Similarly, palm fruit extracts (Hoseinifar *et al.*, 2017), *Thymus vulgaris* extract (Hoseini *et al.*, 2019), *Santalum album* essential oil (Mansour *et al.*, 2024), *Lavandula angustifolia* extract (Yousefi *et al.*, 2020) could increase SOD, CAT, GPX and/or GST expression in fish tissue.

In recent years, gene expression analysis has become increasingly important in assessing the status of fish immunity using reliable technological techniques. Interleukin-8 (IL-8) functions as a crucial chemokine in fish, mediating the activation and recruitment of neutrophils to sites of inflammation, and directing immune cell migration, thus playing a pivotal role in regulating immune responses and enhancing the defense against bacterial pathogens (Ahmadifar *et al.*, 2021). Interleukin-1 beta (IL-1 β) acts as a central pro-inflammatory cytokine in fish, primarily produced by macrophages and monocytes, it not only mediates inflammatory responses by enhancing the production of other cytokines but also initiates crucial immune processes necessary for combating infections (Dawood, 2021). Tumor Necrosis Factor alpha (TNF- α) is a key cytokine that regulates acute inflammatory responses and immune cell functions in fish, significantly impacting systemic inflammation and playing a central role in both inflammation and apoptosis during pathogenic attacks (Ahmadifar *et al.*, 2021; Dawood, 2021). The present study determined that the IL-8, IL-1 β , and TNF- α gene expression levels in the spleen and/or liver were significantly higher in the group supplemented with all myrrh essential oil supplemented groups compared to the control group. Moreover, as the concentration of Myrrh essential oil in the diet increased, there was a proportional rise in Relative Percent Survival (RPS) values, suggesting a dose-dependent reduction in mortality from *Aeromonas hydrophila* infection. Similar to our results, Spirulina platensis (*Arthrospira platensis*) and Sage (*Salvia officinalis*) supplementation significantly increased the expression of IL-1 β and TNF- α in head kidney tissue in *Oreochromis niloticus*, and protecting against *Pseudomonas aeruginosa*. Trans-cinnamic acid was reported to increase the head kidney expression levels of immune-related genes, including TNF- α , IL-8, IL-1 β , and other cytokines, in rainbow trout (*Oncorhynchus mykiss*), enhancing their disease resistance against *Yersinia ruckeri* (Yilmaz and Sergün, 2018).

Conclusions

The pharmacological properties of myrrh are due to a variety of phytochemicals, including terpenoids (monoterpenoids, sesquiterpenoids and essential oils), diterpenoids, triterpenoids and steroids (Hanus et al., 2005; Marcotullio et al., 2009; Batiha et al., 2023). Addition of CMO to the diets of common carp, especially at concentrations of 0.5% or 1.0%, has evidently improved their antioxidant status and immune responses, thereby enhancing their resistance against the pathogen *Aeromonas hydrophila*. This study confirms the potential of Myrrh essential oil as a natural additive to improve disease resistance in aquaculture. As this research represents the first investigation into the culture of various aquatic species, particularly carp, it could provide valuable insights into fish health, blood parameters, and gene expression. However, further studies are required to assess growth performance, feed utilization, and the anti-parasitic effects of myrrh in fish. In particular, long-term trials and investigations involving other fish species are recommended.

Ethical Statement

This study was approved by the Animal Ethics Committee of Muğla Sıtkı Koçman University (Muğla, Türkiye, Approval Number: 2024/02-2)

Funding Information

This research received no external funding.

Author Contribution

Conceptualization, Ü.A. Ö.Y.; methodology, Ü.A., E.B. and F.Z.N.; validation, Ü.A. and S.Y.; formal analysis, Ü.A., E.B. M.H. and F.Z.N.; data curation, Ü.A., E.B. and F.Z.N.; writing—original draft preparation, Ü.A., Ö.Y. and S.Y.; writing—review and editing, Ü.A. , Ö.Y. All authors have read and agreed to the published version of the manuscript

Conflict of Interest

The author(s) declare that they have no known competing financial or non-financial, professional, or personal conflicts that could have appeared to influence the work reported in this paper.

References

- Abdel-Latif, H. M. R., Abdel-Tawwab, M., Khafaga, A. F., & Dawood, M. A. O. (2020). Dietary oregano essential oil improved the growth performance via enhancing the intestinal morphometry and hepato-renal functions of common carp (*Cyprinus carpio* L.) fingerlings. *Aquaculture*, 526, 735432. <https://doi.org/10.1016/j.aquaculture.2020.735432>
- Abdelsalam, A. H., & Ila, H. B. (2022). In vitro cytogenotoxic and mutagenic effects of *Commiphora myrrha* essential oil. *Drug and Chemical Toxicology*, 45(6), 2718–2726. <https://doi.org/10.1080/01480545.2021.1984518>
- Abdel-Tawwab, M., Khattab, Y. A., Ahmad, M. H., & Shalaby, A. M. (2006). Compensatory growth, feed utilization, whole-body composition, and hematological changes in starved juvenile Nile tilapia, *Oreochromis niloticus* (L.). *Journal of Applied Aquaculture*, 18(3), 17–36. https://doi.org/10.1300/J028v18n03_02
- Acar, Ü., Kesbiç, O. S., Yılmaz, S., Gültepe, N., & Türker, A. (2015). Evaluation of the effects of essential oil extracted from sweet orange peel (*Citrus sinensis*) on growth rate of tilapia (*Oreochromis mossambicus*) and possible disease resistance against *Streptococcus iniae*. *Aquaculture*, 437, 282–286. <https://doi.org/10.1016/j.aquaculture.2014.12.015>
- Adeli, A., & Shamloofar, M., & Akrami, R. (2021). Dietary effect of Lemon Verbena (*Aloysia triphylla*) extract on growth performance, haematological, biochemical, and non-specific immunity, and stocking density challenge of rainbow trout juveniles (*Oncorhynchus mykiss*). *Journal of Applied Animal Research*, 49(1), 382–390. <https://doi.org/10.1080/09712119.2021.1990069>
- Ahmadifar, E., Pourmohammadi Fallah, H., Yousefi, M., Dawood, M. A. O., Hoseinifar, S. H., Adineh, H., Yılmaz, S., Paolucci, M., & Doan, H. V. (2021). The gene regulatory roles of herbal extracts on the growth, immune system, and reproduction of fish. *Animals*, 11(8), 2167. <https://doi.org/10.3390/ani11082167>
- Austin, B., & Adams, C. (1996). Fish pathogens. In B. Austin, M. Altwegg, P. J. Gosling, & S. Joseph (Eds.), *The genus Aeromonas* (pp. 197–243). John Wiley and Sons.
- Austin, B., & Austin, D. A. (2007). *Bacterial fish pathogens: Diseases of farmed and wild fish* (4th ed.). Springer-Verlag.
- Baba, E., Acar, Ü., Öntaş, C., Kesbiç, O. S., & Yılmaz, S. (2016). Evaluation of Citrus limon peels essential oil on growth performance, immune response of Mozambique tilapia *Oreochromis mossambicus* challenged with *Edwardsiella tarda*. *Aquaculture*, 465, 15–16. <https://doi.org/10.1016/j.aquaculture.2016.08.023>
- Batiha, G. E., Wasef, L., Teibo, J. O., Shaheen, H. M., Zakariya, A. M., Akinfe, O. A., Teibo, T. K. A., Al-Kuraishy, H. M., Al-Garbee, A. I., Alexiou, A., & Papadakis, M. (2023). *Commiphora myrrha*: A phytochemical and pharmacological update. *Naunyn-Schmiedeberg's Archives of Pharmacology*, 396(3), 405–420. <https://doi.org/10.1007/s00210-022-02325-0>
- Cabello, F. C. (2006). Heavy use of prophylactic antibiotics in aquaculture: A growing problem for human and animal health and for the environment. *Environmental Microbiology*, 8(7), 1137–1144. <https://doi.org/10.1111/j.1462-2920.2006.01054.x>
- Chen, Y., Zhou, C., Ge, Z., Liu, Y., Liu, Y., Feng, W., Li, S., Chen, G., Wei, T. (2013). Composition and potential anticancer activities of essential oils obtained from myrrh and frankincense. *Oncol Lett.* Oct;6(4):1140-1146. doi: 10.3892/ol.2013.1520.
- Cho, K. H. (2012). 1,8-cineole protected human lipoproteins from modification by oxidation and glycation and exhibited serum lipid-lowering and anti-inflammatory activity in zebrafish. *BMB Reports*, 45(10), 565–570. <https://doi.org/10.5483/BMBRep.2012.45.10.044>
- Clauss, T. M., Dove, A. D., & Arnold, J. E. (2008). Hematologic disorders of fish. *Veterinary Clinics of North America: Exotic Animal Practice*, 11(3), 445–462.

- <https://doi.org/10.1016/j.cvex.2008.03.007>
- Cunha, J. A., Heinzmann, B. M., & Baldisserotto, B. (2018). The effects of essential oils and their major compounds on fish bacterial pathogens: A review. *Journal of Applied Microbiology*, 125(2), 328–344. <https://doi.org/10.1111/jam.13911>
- Dawood, M. A. (2021). Nutritional immunity of fish intestines: Important insights for sustainable aquaculture. *Reviews in Aquaculture*, 13(1), 642–663. <https://doi.org/10.1111/raq.12492>
- Demissew, S. (1993). A description of some essential oil bearing plants in Ethiopia and their indigenous uses. *Journal of Essential Oil Research*, 5(5), 465–479. <https://doi.org/10.1080/10412905.1993.9698266>
- Dias, C., Mota, V., Martinez-Murcia, A., & Saavedra, M. J. (2012). Antimicrobial resistance patterns of *Aeromonas* spp. isolated from ornamental fish. *Journal of Aquaculture Research & Development*, 3, 1000131. <https://doi.org/10.4172/2155-9546.1000131>
- Dorojan, O. G., Varlan, V., Cristea, M., Crețu, L., Dediu, A. I., Docan, M. T., & Coadă, M. T. (2015). The effect of thyme (*Thymus vulgaris*) and vitamin E on the welfare of juvenile *Acipenser stellatus*, reared in a recirculating aquaculture system. *AAFL Bioflux*, 8(2), 150–158.
- Dotta, G., de Andrade, J. I., Tavares Gonçalves, E. L., Brum, A., Mattos, J. J., Maraschin, M., & Martins, M. L. (2014). Leukocyte phagocytosis and lysozyme activity in Nile tilapia fed supplemented diet with natural extracts of propolis and *Aloe barbadensis*. *Fish & Shellfish Immunology*, 39(2), 280–284. <https://doi.org/10.1016/j.fsi.2014.05.020>
- Eddy, S., & Underhill, J. C. (1974). *Northern fishes with special reference to the upper Mississippi Valley* (3rd ed.). University of Minnesota Press.
- Edris, A. E. (2007). Pharmaceutical and therapeutic potentials of essential oils and their individual volatile constituents: A review. *Phytotherapy Research*, 21(3), 308–323. <https://doi.org/10.1002/ptr.2072>
- El-Houseiny, W., Khalil, A. A., Abd-Elhakim, Y. M., Arisha, A. H., Moselhy, A. A., Dahshan, H., & Ahmed, M. M. (2022). Alleviative effects of dietary *Silybum marianum* and co-enzyme Q10 on waterborne nickel-induced impaired growth, immunosuppression, tissue damage, immune-related genes dysregulation, and reduced resistance to *Pseudomonas aeruginosa* in *Oreochromis niloticus*. *Aquaculture Reports*, 26, 101308. <https://doi.org/10.1016/j.aqrep.2022.101308>
- Encyclopedia, 2024. Available from <https://www.encyclopedia.com/plants-and-animals/plants/plants/myrrh> (8 December 2024).
- Farag, M. R., Zheng, C., Gharib, H. S., El-Hady, E., Mahdy, E. A., Abo-Elmaaty, A., & Hassan, M. A. (2023). Physiological and neurobehavioral disturbances induced by Al₂O₃ nanoparticle intoxication in Nile tilapia fish: Benefits of dietary chamomile essential oil. *Aquaculture Nutrition*, 6700708. <https://doi.org/10.1155/2023/6700708>
- Fazio, F., Habib, S. S., Naz, S., Filiciotto, F., Cicero, N., Rehman, H. U., Saddozai, S., Rind, K. J., Rind, N.A., Shar, A. H. (2021). Effect of fortified feed with olive leaves extract on the haematological and biochemical parameters of *Oreochromis niloticus* (Nile tilapia). *Natural Product Research*, 36(6), 1575–1580. <https://doi.org/10.1080/14786419.2021.1883606>
- Fazio, F., Marafioti, S., Torre, A., Sanfilippo, M., Panzera, M., & Faggio, C. (2013). Haematological and serum protein profiles of *Mugil cephalus*: Effect of two different habitats. *Ichthyological Research*, 60(1), 36–42. <https://doi.org/10.1007/s10228-012-0303-1>
- Gabriel, N. N., Qiang, J., He, J., Ma, X. Y., Kpundeh, M. D., & Xu, P. (2015). Dietary Aloe vera supplementation on growth performance, some haemato-biochemical parameters and disease resistance against *Streptococcus iniae* in tilapia (GIFT). *Fish & Shellfish Immunology*, 44(2), 504–514. <https://doi.org/10.1016/j.fsi.2015.03.002>
- Gadir, S. A., Ahmed, I.M. (2014). *Commiphora myrrha* and commiphora Africana essential oils. *Journal of Chemical and Pharmaceutical Research*, 6(7):151-156
- Gaudet, M., Racicot, J. G., & Leray, C. (1975). Enzyme activities of plasma and selected tissues in rainbow trout *Salmo gairdneri* Richardson. *Journal of Fish Biology*, 7(4), 505–512. <https://doi.org/10.1111/j.1095-8649.1975.tb04625.x>
- Ghafarifarsani, H., Hoseinifar, S. H., Sheikhlari, A., Raissy, M., Chaharmahali, F. H., Maneepitaksanti, W., Faheem, M., & Doan, H. V. (2022). The effects of dietary thyme oil (*Thymus vulgaris*) essential oils for common carp (*Cyprinus carpio*): Growth performance, digestive enzyme activity, antioxidant defense, tissue and mucus immune parameters, and resistance against *Aeromonas hydrophila*. *Aquaculture Nutrition*, 7942506, 13 pages. <https://doi.org/10.1155/2022/7942506>
- Ghelichpour, M., Mirghaed, A. T., Dawood, M. A., Hoseinifar, S. H., & Van Doan, H. (2021). Alteration of haematological and antioxidant parameters in common carp (*Cyprinus carpio*) fed olive (*Olea europea*) leaf extract after exposure to Danitol®. *Aquaculture Research*, 52(3), 1088–1095. <https://doi.org/10.1111/are.14964>
- Habib, S. S., Saeed, M. Q., Masud, S., Kesbiç, O. S., Ujan, J. A., Cravana, C., Al-Rehaie, S.A., Mohany, M., & Fazio, F. (2024). Dietary Effect of Withania somnifera Root Powder on Growth, Hematobiochemical Parameters, Immunity, and Disease Resistance to *Aeromonas hydrophila* in *Cyprinus carpio*. *Aquaculture Nutrition*, 2024(1), 7321287. <https://doi.org/10.1155/2024/7321287>
- Haffor, Al-Said A. (2010): Effect of *Commiphora molmol* on leukocytes proliferation in relation to histological alterations before and during healing from injury. *Saudi Journal of Biological Sciences* 17, 139–146. <https://doi.org/10.1016/j.sjbs.2010.02.007>
- Hanus, L.O., Rezanka, T., Dembitsky V.M., Moussaieff, A. (2005). Myrrh--Commiphora chemistry. *Biomed Papers*, Jun;149(1):3-27. doi: 10.5507/bp.2005.001.
- Harikrishnan, R., Balasundaram, C., & Heo, M. S. (2012). Effect of Inonotus obliquus-enriched diet on hematology, immune response, and disease protection in kelp grouper, *Epinephelus bruneus*, against *Vibrio harveyi*. *Aquaculture*, 344, 48–53. <https://doi.org/10.1016/j.aquaculture.2012.03.010>
- Hashimoto, G. S. O., Neto, F. M., Ruiz, M. L., Achille, M., Chagas, E. C., Chaves, F. C. M., & Martins, M. L. (2016). Essential oils of *Lippia sidoides* and *Mentha piperita* against monogenean parasites and their influence on the hematology of Nile tilapia. *Aquaculture*, 450, 182–186. <https://doi.org/10.1016/j.aquaculture.2015.07.029>
- Hesser, E. F. (1960). Methods for routine fish hematology. *Progressive Fish-Culturist*, 22(4), 164–171. [https://doi.org/10.1577/1548-8659\(1960\)22\[164:MFRFH\]2.0.CO;2](https://doi.org/10.1577/1548-8659(1960)22[164:MFRFH]2.0.CO;2)

- Hoseini, S. M., & Yousefi, M. (2019). Beneficial effects of thyme (*Thymus vulgaris*) extract on oxytetracycline-induced stress response, immunosuppression, oxidative stress, and enzymatic changes in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Nutrition*, 25(2), 298–309. <https://doi.org/10.1111/anu.12853>
- Hoseinifar, S. H., Dadar, M., Khalili, M., Cerezuela, R., & Esteban, M. Á. (2017). Effect of dietary supplementation of palm fruit extracts on the transcriptomes of growth, antioxidant enzyme, and immune-related genes in common carp (*Cyprinus carpio*) fingerlings. *Aquaculture Research*, 48(7), 3684–3692. <https://doi.org/10.1111/are.13192>
- Hussain, A. I., Anwar, F., Sherazi, S. T. H., & Przybylski, R. (2008). Chemical composition, antioxidant and antimicrobial activities of basil (*Ocimum basilicum*) essential oils depends on seasonal variations. *Food Chemistry*, 108(3), 986–995. <https://doi.org/10.1016/j.foodchem.2007.12.010>
- Immanuel, G., Uma, R. P., Iyapparaj, P., Citarasu, T., Punitha Peter, S. M., Michael Babu, M., & Palavesam, A. (2009). Dietary medicinal plant extracts improve growth, immune activity and survival of tilapia *Oreochromis mossambicus*. *Journal of Fish Biology*, 74(7), 1462–1475. <https://doi.org/10.1111/j.1095-8649.2009.02212.x>
- Kesbiç, O. S., Parrino, V., Acar, Ü., Yilmaz, S., Paro, G. L., & Fazio, F. (2020). Effects of Monterey cypress (*Cupressus macrocarpa* Hartw) leaf essential oil as a dietary supplement on growth performance and haematological and biochemical parameters of common carp (*Cyprinus carpio* L.). *Annals of Animal Science*, 20(4), 1411–1426. <https://doi.org/10.2478/aoas-2020-0041>
- Khalil, A. A., Abd-Elhakim, Y. M., Said, E. N., Moselhy, A. A., Abu-Elsaoud, A. M., & El-Houseiny, W. (2022). Milk thistle and co-enzyme Q10 fortified diets lessen the nickel chloride-induced neurotoxic and neurobehavioral impairments in *Oreochromis niloticus* via regulating the oxidative stress response, acetylcholinesterase activity, and brain nickel content. *Aquaculture*, 553, 738102. <https://doi.org/10.1016/j.aquaculture.2022.738102>
- Madia, V.N., De Angelis, M., De Vita, D., Messore, A., De Leo, A., Ialongo, D., Tudino, V., Saccoliti, F., De Chiara, G., Garzoli, S., Scipione, L., Palamara, A.T., Di Santo, R., Nencioni, L., Costi, R. (2021) Investigation of *Commiphora myrrha* (Nees) Engl. Oil and Its Main Components for Antiviral Activity. *Pharmaceuticals (Basel)*. Mar 9;14(3):243. <https://doi.org/10.3390/ph14030243>.
- Magnadóttir, B. (2006). Innate immunity of fish (overview). *Fish & Shellfish Immunology*, 20(2), 137–151. <https://doi.org/10.1016/j.fsi.2004.09.006>
- Mansour, A. T., Arisha, A. H., Abdelaziz, R., Alwutayd, K. M., Van Doan, H., El-Murr, A. E., & El-Houseiny, W. (2024). Effects of extended dietary supplementation with Santalum album essential oil on hemato-biochemical changes, innate immune response, antioxidant status, and expression of related gene in Nile tilapia (*Oreochromis niloticus*). *Fish Physiology and Biochemistry*, 1–17. <https://doi.org/10.1007/s10695-024-01309-6>
- Marcotullio MC, Santi C, Mwankie GN, Curini M. (2009). Chemical composition of the essential oil of *Commiphora erythraea*. *Natural Product Communication* Dec;4(12). <https://doi.org/10.1177/1934578X0900401227>
- Martínez-Porchas, M., Martínez-Córdova, L. R., & Ramos-Enriquez, R. (2009). Cortisol and glucose: Reliable indicators of fish stress? *Pan-American Journal of Aquatic Sciences*, 4(2), 158–178.
- Mathew, R. T., Debnath, S., Kundu, P., Alkhamis, Y. A., Rahman, M. M., & Rahman, S. M. (2022). Growth and survival of goldfish (*Carassius auratus*) juveniles fed tubifex, custard meal and commercial feeds. *Scientific Journal of King Faisal University: Basic and Applied Sciences*, 23(1), 30–35. <https://doi.org/10.37575/b/vet/210080>
- Mitsumoto T, Ishii Y, Namiki M, Nakamura K, Takasu S, Ogawa K. A 90-day subchronic toxicity study of Myrrh in F344 rats. (2021). *Regul Toxicol Pharmacol*. 2021 Dec; 127:105076. <https://doi.org/10.1016/j.yrtph.2021.105076>.
- Naz, S., Majeed, S., Tasleem, S., Renuka Jyothi, S., Thakur, H., Anuradha, J., Ujan, J.A., Ullah, M., Zahid, M., Attaullah, S., Mohany, M. & Habib, S. S. (2024). Investigating the effects of wild olive *Olea europaea* leaf extract on growth, body composition, digestive enzyme activity, hematological parameters, and immune function in Nile Tilapia. *North American Journal of Aquaculture*, 86(4), 462–474. <https://doi.org/10.1002/naaq.10354>
- Ngugi, C. C., Oyoo-Okoth, E., & Muchiri, M. (2017). Effects of dietary levels of essential oil (EO) extract from bitter lemon (*Citrus limon*) fruit peels on growth, biochemical, haemato-immunological parameters and disease resistance in juvenile *Labeo victorinus* fingerlings challenged with *Aeromonas hydrophila*. *Aquaculture Research*, 48(5), 2253–2265. <https://doi.org/10.1111/are.13062>
- Nofal, M.Y., Alabodi, A.B., Alenazi, H.K., Alenazi, O.H., Yahia, A.M.B., Al-Shehri, N.Q. (2018). Effect of *Commiphora Molmol* (Myrrh) on Total Leukocyte Count and Histological Alterations in Mice. *International Journal of Pharmaceutical Research & Allied Sciences*, 7(4):60-66.
- Oberley, L. W., & Oberley, T. D. (1988). Role of antioxidant enzymes in cell immortalization and transformation. *Molecular and Cellular Biochemistry*, 84(2), 147–153. <https://doi.org/10.1007/BF00421049>
- Pourmozaffar, S., Hajimoradloo, A., Paknejad, H., & Rameshi, H. (2019). Effect of dietary supplementation with apple cider vinegar and propionic acid on hemolymph chemistry, intestinal microbiota and histological structure of hepatopancreas in white shrimp, *Litopenaeus vannamei*. *Fish & Shellfish Immunology*, 86, 900–905. <https://doi.org/10.1016/j.fsi.2018.12.019>
- Rashidian, G., Kajbaf, K., Prokić, M. D., & Faggio, C. (2020). Extract of common mallow (*Malva sylvestris*) enhances growth, immunity, and resistance of rainbow trout (*Oncorhynchus mykiss*) fingerlings against *Yersinia ruckeri* infection. *Fish & Shellfish Immunology*, 96, 254–261. <https://doi.org/10.1016/j.fsi.2019.12.018>
- Remmal, A., Tantaoui Elaraki, A., Bouchikhi, T., & Ettayebi, M. (1993). Inhibition of antibacterial activity of essential oils by Tween 80 and ethanol in liquid medium. *Journal of Pharmacy Belgium*, 48, 352–356.
- Rinchard, J., Lee, K. J., Dabrowski, K., Ciereszko, A., Blom, J. H., & Ottobre, J. S. (2003). Influence of gossypol from dietary cottonseed meal on haematology, reproductive steroids and tissue gossypol enantiomer concentrations in male rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Nutrition*, 9(4), 275–282. <https://doi.org/10.1046/j.1365-2095.2003.00253.x>
- Romero, J., Feijoo, C. G., & Navarrete, P. (2012). Antibiotics in

- aquaculture: Use, abuse and alternatives. In E. D. Carvalho, G. S. David, & R. J. Silva (Eds.), *Health and environment in aquaculture* (pp. 159–184). InTech Europe.
- Saikia, S. K., & Das, D. N. (2009). Feeding ecology of common carp (*Cyprinus carpio* L.) in a rice-fish culture system of the Apatani plateau (Arunachal Pradesh, India). *Aquatic Ecology*, 43(3), 559–568. <https://doi.org/10.1007/s10452-008-9174-y>
- Semwal, A., Kumar, A., & Kumar, N. (2023). A review on pathogenicity of *Aeromonas hydrophila* and their mitigation through medicinal herbs in aquaculture. *Heliyon*, 9, e14088. <https://doi.org/10.1016/j.heliyon.2023.e14088>
- Sheikhzadeh, N., Soltani, M., Mousavi, H. E., Khosravi, A., Bagheri, H., & Fathi, E. (2009). Effects of Eucalyptus globules labill essential oil on some immunological variables of common carp (*Cyprinus carpio*). *Journal of Veterinary Research*, 64, 47-4.
- Shen, T., Lou, HX. (2008). Bioactive constituents of myrrh and frankincense, two simultaneously prescribed gum resins in chinese traditional medicine. *Chem Biodivers*. 2008 Apr;5(4):540-53. <https://doi.org/10.1002/cbdv.200890051>.
- Shourbela, R. M., Khatib, S. A., Hassan, M. M., Van Doan, H., & Dawood, M. A. (2021). The effect of stocking density and carbon sources on the oxidative status and nonspecific immunity of Nile tilapia (*Oreochromis niloticus*) reared under biofloc conditions. *Animals*, 11(1), 184. <https://doi.org/10.3390/ani11010184>
- Simanjuntak, S. B. I., Indarmawan, I., & Wibowo, E. S. (2018). Impact of diets containing different levels of Spirulina platensis supplementation on growth, haematological, body composition, and biochemical parameters of Gurami (*Osphronemus gouramy*). *Turkish Journal of Fisheries and Aquatic Sciences*, 18(5), 681–690. https://doi.org/10.4194/1303-2712-v18_5_04
- Soltani, M., Sheikhzadeh, N., Ebrahimzadeh-Mousavi, H., & Zargar, A. (2010). Effects of Zataria multiflora essential oil on innate immune responses of common carp (*Cyprinus carpio*). *Journal of Fisheries & Aquatic Sciences*, 5(3), 191–199.
- Souza, E. M. D., Souza, R. C. D., Melo, J. F., Costa, M. M. D., Souza, S. A. D., Souza, A. M. D., & Copatti, C. E. (2020). *Cymbopogon flexuosus* essential oil as an additive improves growth, biochemical and physiological responses, and survival against *Aeromonas hydrophila* infection in Nile tilapia. *Anais da Academia Brasileira de Ciências*, 92(1), e20190140. <https://doi.org/10.1590/0001-3765202020190140>
- Su, S., Wang, T., Chen, T., Duan, J., Yu, L., & Yuping, T. (2011). Cytotoxicity activity of extracts and compounds from *Commiphora myrrha* resin against human gynecologic cancer cells. *Journal of Medicinal Plant Research*, 5(8), 1382–1389.
- Swamy, M. K., Akhtar, M. S., & Sinniah, U. R. (2016). Antimicrobial properties of plant essential oils against human pathogens and their mode of action: An updated review. *Evidence-Based Complementary and Alternative Medicine*, 21, 3012462. <https://doi.org/10.1155/2016/3012462>
- Tang, J., Cai, J., Liu, R., Wang, J., Lu, Y., Wu, Z., & Jian, J. (2014). Immunostimulatory effects of artificial feed supplemented with a Chinese herbal mixture on *Oreochromis niloticus* against *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, 39(2), 401–406. <https://doi.org/10.1016/j.fsi.2014.05.028>
- Ujan, J. A., Raza, A., Habib, S. S., Athar, R., Kesbiç, O. S., Batool, A. I., Habib, S. F., Mohany, M. & Fazio, F. (2025). Nettle (*Urtica dioica*) supplementation: impact on growth, hematology, immune response, and resilience to *Aeromonas hydrophila* in Labeo rohita fingerlings. *Veterinary Research Communications*, 49(1), 4. <https://doi.org/10.1007/s11259-024-10577-9>
- Wu, C. C., Hung Liu, C., Chang, Y. P., & Hsieh, S. L. (2010). Effects of hot-water extract of *Toona sinensis* on immune response and resistance to *Aeromonas hydrophila* in *Oreochromis mossambicus*. *Fish & Shellfish Immunology*, 29(2), 258–263. <https://doi.org/10.1016/j.fsi.2010.04.021>
- Wu, Y., Gong, O., Fang, H., Liang, W., Chen, M., & He, R. (2013). Effect of *Sophora flavescens* on non-specific immune response of tilapia (GIFT *Oreochromis niloticus*) and disease resistance against *Streptococcus agalactiae*. *Fish & Shellfish Immunology*, 34(2), 220–222. <https://doi.org/10.1016/j.fsi.2012.10.020>
- Yilmaz, S., & Ergün, S. (2018). Trans-cinnamic acid application for rainbow trout (*Oncorhynchus mykiss*): Effects on haematological, serum biochemical, non-specific immune, and head kidney gene expression responses. *Fish & Shellfish Immunology*, 78, 140–157. <https://doi.org/10.1016/j.fsi.2018.04.034>
- Yilmaz, S., Ergün, S., Yilmaz, E., Ahmadifar, E., Yousefi, M., & Abdel-Latif, H. M. (2024). Effects of a phyto-genic diet on growth, haemato-immunological parameters, expression of immune- and stress-related genes, and resistance of *Oncorhynchus mykiss* to *Lactococcus garvieae* infection. *Aquaculture*, 587, 740845. <https://doi.org/10.1016/j.aquaculture.2024.740845>
- Yousefi, M., Shabunin, S. V., Vatnikov, Y. A., Kulikov, E. V., Adineh, H., Hamidi, M. K., & Hoseini, S. M. (2020). Effects of lavender (*Lavandula angustifolia*) extract inclusion in diet on growth performance, innate immunity, immune-related gene expression, and stress response of common carp, *Cyprinus carpio*. *Aquaculture*, 515, 734588. <https://doi.org/10.1016/j.aquaculture.2019.734588>
- Zadmajid, V., & Mohammadi, C. (2017). Dietary thyme essential oil (*Thymus vulgaris*) changes serum stress markers, enzyme activity, and hematological parameters in gibel carp (*Carassius auratus gibelio*) exposed to silver nanoparticles. *Iranian Journal of Fisheries Sciences*, 16(3), 1063–1084. <https://doi.org/10.22092/IJFS.2018.114720>
- Zahran, E., Risha, E., AbdelHamid, F., Mahgoub, H. A., & Ibrahim, T. (2014). Effects of dietary Astragalus polysaccharides (APS) on growth performance, immunological parameters, digestive enzymes, and intestinal morphology of Nile tilapia (*Oreochromis niloticus*). *Fish & Shellfish Immunology*, 38(1), 149–157. <https://doi.org/10.1016/j.fsi.2014.03.002>
- Zhang, T., Zhang, Y., Jiang, N., Zhao, X., Sang, X., Yang, N., & Chen, Q. (2020). Dihydroartemisinin regulates the immune system by promotion of CD8+ T lymphocytes and suppression of B cell responses. *Science China Life Sciences*, 63(5), 737–749. <https://doi.org/10.1007/s11427-019-9550-4>