



Length Based Growth Estimation of Most Commercially Important Scombridae from Offshore Water of Pakistan Coast in the Arabian Sea

Quratulan Ahmed^{1,*}, Sabri Bilgin², Levent Bat²

¹ The Marine Reference Collection and Resource Centre, University of Karachi, Pakistan.

² Sinop University, Faculty of Fisheries, Sinop, Turkey.

Email: quratulanahmed_ku@yahoo.com

Abstract

Seasonal growth pattern of skipjack tuna (*Katsuwonus pelamis*), longtail tuna (*Thunnus tonggol*), kawakawa (*Euthynnus affinis*) and king mackerel (*Scomberomorus guttatus*) were studied during five years period (between August 2006 and December 2011). Seasonal samplings of the species were collected from offshore water of Pakistan coast in the Arabian Sea. The weight - length relationships (WLRs) among the species except *E. affinis* demonstrated that the growth is allometric type (Pauly's *t* test, $P < 0.005$). The seasonal von Bertalanffy growth model parameters computed from seasonal length - frequency distributions, were estimated as $L_{\infty} = 67.1$ cm TL, $K = 1.055$ year⁻¹, $C = 0.990$ for *E. affinis*, as $L_{\infty} = 66.8$ cm TL, $K = 1.378$ year⁻¹ and $C = 0.950$ for *K. pelamis*, as $L_{\infty} = 71.7$ cm TL, $K = 0.916$ year⁻¹ and $C = 0.849$ for *T. tonggol*, as $L_{\infty} = 55.7$ cm TL, $K = 1.049$ year⁻¹ and $C = 0.980$ for *S. guttatus*. The seasonal oscillation in growth rate was determined for the species ($C > 0.8$). The slowest period of growth corresponded to in September ($WP = 0.7$) for *E. affinis*, ($WP = 0.6$) in August for *K. pelamis*, ($WP = 0.4$) in May for *T. tonggol* and ($WP = 0.0$) in January for *S. guttatus*.

Keywords: Scombridae, seasonal growth, LFDA, ELEFAN, Arabian Sea, Pakistan

Introduction

Gillnet is the main fishing gear used for catching tuna and other large pelagic fishes in many countries of the world including Pakistan (Moazzam and Nawaz, 2015). There are four main harbors and nine primary landing sites along the coast of Pakistan. The largest is Karachi fish harbor, which is used by approximately 80 - 90% of the industrial fishing fleet, almost exclusively shrimp trawlers and larger gillnetters (Hornby et al., 2014). The gillnets are the gear employed for catching tuna by these wooden-hulled vessels and about 500 gillnet fishing vessels are engaged in the inshore and offshore fishery for large pelagic species with most vessels ranging between 10 and 20 m (Hornby et al., 2014; Khan, 2012). Only about 30 vessels are between 20 to 30 m and have on board freezing compartments and dual registration to fish in Pakistan and Iran's exclusive economic zone, EEZ (Khan, 2012). Large pelagic species such as skipjack tuna, *Katsuwonus pelamis*, longtail tuna, *Thunnus tonggol*, kawakawa, *Euthynnus affinis*, king mackerel, *Scomberomorus guttatus* and narrow-barred Spanish mackerel, *Scomberomorus commerson* support some of the more important fisheries in Pakistan (FAO, 2015). According to FAO statistics for last four years between 2010 and 2013, total catch of these fish species was reported as between 28463 and 30520 ton (mean: 29710.8±490.29 ton), while total marine fish catch was between 294814 and 308167 ton (mean: 302826.6±6060.19 ton) from the Pakistan coast (FAO, 2015) in the Arabian Sea.

This means that Scombrids fish catches constitutes about 10% of the total marine fish catch in Pakistan.



The distribution of *K. pelamis* includes the offshore waters with temperature ranging from 14.7 to 30°C (circumglobal in seas warmer than 15°C) while larvae are mostly restricted to waters with surface temperatures of at least 25°C (van der Land et al., 2001; Collette et al., 2011a; Froese and Pauly, 2015). *T. tonggol* is Indo-Pacific species inhabiting from the Red Sea and East Africa to Papua New Guinea, north to Japan, and south to Australia (Collette et al., 2011b; Froese and Pauly, 2015). The distribution of *E. affinis* includes Indian and Western Pacific Ocean and occurs in open waters with temperatures ranging from 18°C to 29°C but always remains close to the shoreline (Collette and Nauen, 1983; Collette et al., 2011c; Froese and Pauly, 2015). The distribution areas of *S. guttatus* include Indo-Pacific from the Persian Gulf, India and Sri Lanka to Southeast Asia, north to Hong Kong and Wakasa Bay, Sea of Japan (van der Land et al., 2001; Froese and Pauly, 2015). *S. commerson* is pelagic and oceanodromous and found in the Indo-West Pacific from the Red Sea and South Africa to Southeast Asia, north to China and Japan and south to southeast Australia, and to Fiji shrimps (Collette et al., 2011e; Froese and Pauly, 2015). These scomberids fish species feeds on a variety of fish, crustaceans, cephalopods, mollusks and spawns all year long in tropical waters and from spring to early fall in subtropical waters (Devaraj, 1998; van der Land et al., 2001; Collette et al., 2011a, b, c, d, e; Froese and Pauly, 2015).

Fish growth can be definite changes in fish length or weight over time and it is mostly rapid in early life most probably associated with the size at sexual maturity and slows with age (Von Bertalanffy, 1951). Growth is also considered to be the result of a counteraction of anabolism and catabolism of building materials and anabolic activities are in perpetuity greater than the catabolic activities for fish (Avşar, 2005). A number of environmental factors, such as temperature, oxygen concentration, salinity and photoperiod, influence the rate of growth. Nutrition, including the quality and quantity of food, also plays a significant role in growth regulation (Dutta, 1994). Fish exhibit a determinate type of growth in short-lived species of warmer regions and an indeterminate type in long-lived species of colder regions (Dutta, 1994). The most common type of animal growth is that growth rates continually decrease with time and finally the organism reaches a steady state in the adult (Von Bertalanffy, 1951). The growth of fish can also display seasonal oscillations, mainly due to fluctuations of temperature, food supply and investment of energy during spawning and maturity time.

Historically, different methods have been applied to determine of growth of Scombrids fish obtained different area. These studies based on length frequency analysis (LFA), hard parts such as otolith, dorsal spines and vertebrata and tagging studies. Most of these studies were conducted based on LFA and for this two techniques were used as popularly. Firstly, the Electronic Length Frequency Analysis (ELEFAN) program has been used to fit the von Bertalanffy growth (VBG) curve on seasonal length-frequency distributions without considering the age composition. Secondly, length frequencies have been transformed to ages either by Petersen method or by standard statistical procedures such as Bhattacharya method and then the VBG curve has been fitted to modal lengths (mean length at age) to estimate the growth parameters (Wild and Hampton, 1994; Yesaki, 1994a,b; Siddeek, 1995).

Reviews of the biology and fisheries for *K. pelamis* from Pacific Ocean, *T. tonggol* and *E. affinis* from the indo-pacific region were reported by Wild and Hampton (1994), Yesaki (1994a) and Yesaki (1994b), respectively. Moreover, the VBG curve parameters (L_{∞} , K , t_0) mostly estimated based on the ELEFAN procedure for *E. affinis* were reported from different areas by different authors (Yesaki, 1994b; Khan, 2004; Ahmed et al., 2015). In the previously studies growth estimation using direct and indirect methods for *K. pelamis* were also



reported from different geographical areas by different authors (Chi and Yang, 1973; Koya et al., 2012; Adams and Kerstetter, 2014). Studies concerning parameters estimation of VBG curve for *T. tonggol* from different geographical areas almost all of were based on otolith, modal progression analysis (MPA) and the ELEFAN were reported by Wilson (1981), Spongpan and Saikliang (1987), Yesaki, (1989), Prabhakar and Dubley (1989), Ghosh et al. (2010), Griffiths et al. (2010), Abdussamad et al. (2012) and Kaymaram et al. (2013). Length based growth estimation of *S. guttatus* were reported by Devaraj (1981) from the Palk Bay and the Mannar Gulf and by Rashid et al. (2010) from the Bay of Bengal off Bangladesh coast.

Various growth models have been proposed to explain the growth pattern of individuals for a particular species. Because organisms generally growth seasonally, a good description of the pattern of growth of an organism that lives for a number of years requires a seasonal adjustment to the growth rate (Henderson and Seaby, 2006). A modified von Bertalanffy growth model (five parameters model) has been developed to incorporate such a seasonal growth pattern (Somers, 1988). A seasonal growth pattern has recently been reported from different areas for different fish species. However, there is no detailed information on the seasonal growth rate of Scombrids fish from offshore water of Pakistan coast in the Arabian Sea. The aim of this study was to estimate the seasonal von Bertalanffy growth parameters using length frequency data for skipjack tuna (*K. pelamis*), longtail tuna (*T. tonggol*), kawakawa (*E. affinis*) and king mackerel (*S. guttatus*) during five years period (2006 - 2011) and also investigate the length weight relationships of these species from offshore water of pakistan coast in the Arabian Sea.

Materials and Methods

Study Area and Sampling

Scomberomorus commerson (n = 294), *Scomberomorus guttatus* (n = 278), *Euthynnus affinis* (n= 313), *Thunnus tonggol* (n = 300) and *Katsuwonus pelamis* (n = 325) were collected offshore water in exclusive economic zone (EEZ) of Pakistan coast in the Arabian Sea. Fishes were obtained from commercial fishermen engaged in fishing with gill nets on average 2500 m long, with a mesh size ranging between 13 cm to 17 cm (average 15 cm) and a depth of about 80 meshes during seasonal surveys. Seasons were grouped as winter (between December and February), spring (between March and May), summer (between June and August), and autumn (between September and November).

Weight - Length relationships (WLRs)

The length (*TL*) of the fish was measured from the tip of the anterior part of the mouth to the tip of caudal fin in (cm). Fish weight was measured after blot drying with a piece of clean towel. From the fresh samples were taken total length (*TL*) in (cm) and body weight (*W*) in (g) were measured to the nearest 0.1 cm and 0.01 g, respectively.

Least squares regression analysis with MS Excel software was used to calculate the weight length relationship parameters of all species. The weight length relationship was estimated as:

$$W = aTL^b ;$$

where *W* is the body weight (g), *TL* is the total length (cm), *a* is the intercept, and *b* is the slope of the regression line.



Comparison of the difference of slope value from $b = 3$ (isometric growth) for all species, Pauly's t -test was performed (Pauly 1984). Pauly's t -test statistic was calculated as:

$$t = \frac{Sd_{\log TL} |b - 3|}{Sd_{\log W} \sqrt{1 - r^2}} \sqrt{n - 2}$$

where $Sd_{\log TL}$ is the standard deviation of the log TL values, $Sd_{\log W}$ is the standard deviation of the log W values, n is the number of specimens used in the computation. The value of b is different from $b = 3$ if t value is greater than the tabled t values for $n-2$ degrees of freedom (Pauly, 1984).

Von Bertalanffy Growth Parameter Estimation

Growth in length has been described using the von Bertalanffy (1938) growth equation, based on either observed or back calculated length at ages. The length frequency distribution analysis (LFDA) package is also a PC based computer package for estimating growth parameters from fish length frequency distributions. Version 5.0 of LFDA includes methods for estimating the parameters of both non seasonal and seasonal versions of the von Bertalanffy growth curve (Kirkwood et al., 2003).

The standard or non-seasonal von Bertalanffy (1938) growth function (VBGF) is:

$$L_t = L_\infty (1 - e^{-k(t-t_0)})$$

Seasonal growth or five parameter von Bertalanffy growth model (5 P VBGF) was described using the Somers's (1988) version of the VBG equation:

$$L_t = L_\infty \left[1 - e^{\left[-K(t-t_0) - \left(C \frac{K}{2\pi} \right) \sin 2\pi(t-t_s) + \left(C \frac{K}{2\pi} \right) \sin 2\pi(t_0-t_s) \right]} \right]$$

where, L_t is length at age t , L_∞ is the asymptotic length to which the fish growth, K is the growth-rate parameter, t_0 is the nominal age at which the length is zero, C is the relative amplitude ($0 \leq C \leq 1$) of the seasonal oscillations, t_s is the phase of the seasonal oscillations ($-0.5 \leq t_s \leq 0.5$) denoting the time of year corresponding to the start of the convex segment of sinusoidal oscillation.

The time of the year when the growth rate is slowest, known as the winter point (WP), was calculated as:

$$WP = t_s + 0.5.$$

Seasonal VBG curves were fitted to the length distributions after first indicating a range of values of L_∞ and K and reducing iteratively the range to maximize the goodness of fit (Rn) of the curves to the data. Rn was calculated as:

$$Rn = \frac{10^{ASP}}{ESP}$$

where ASP is the available sum of peaks, computed by adding the best values of the available peaks, and ESP is the explained sum of peaks, computed by summing all the peaks and troughs hit by the VBG curve. Analysis of the seasonal length data were fitted to length frequency distributions grouped in 2 cm total length size classes using the ELEFAN procedure in the PC-based computer package Version 5.0 of LFDA (Kirkwood et al. 2003).



The ELEFAN procedure first restructures length frequencies and then fits a VBG curve to the restructured data. Both seasonal and non-seasonal VBG curves were fitted to the seasonal length distribution after providing a range of values for the parameters to be estimated and then iteratively reducing the range until the goodness of fit of the curve to the data is maximized. Attempts of growth estimation for *S. commerson* were not achieved at all due to length frequency distribution data and therefore, we performed LFDA procedure for *E. affinis*, *K. pelamis*, *S. guttatus* and *T. tonggol*. Because of the fact that significant differences of mean total length and size distribution were not determined between males and females, growth estimations were derived for combined sexes.

Growth performance comparisons were made using the growth performance index (Φ') which is preferred rather than using L_∞ and K individually (Pauly and Munro 1984) and is computed as:

$$\Phi' = \log(K) + 2 \log(L_\infty).$$

Results

A total of 1510 fish were sampled from 2006 to 2011 during fishing seasons. Mean total length of females and males for *E. affinis* (t -test: $P = 0.0575$), *K. pelamis* (t -test: $P = 0.6046$), *S. commerson* (t -test: $P = 0.0327$), *S. guttatus* (t -test: $P = 0.6267$) and *T. tonggol* (t -test: $P = 0.3687$) were not significant different.

Size-frequency distribution between females and males for *E. affinis* (Kolmogorov-Smirnov two sample test: $d = 0.1779$, $P = 0.4298$), *K. pelamis* (Kolmogorov-Smirnov two sample test: $d = 0.1015$, $P = 0.8981$), *S. commerson* (Kolmogorov-Smirnov two sample test: $d = 0.1241$, $P = 0.3113$), *S. guttatus* (Kolmogorov-Smirnov two sample test: $d = 0.0699$, $P = 0.9424$) and *T. tonggol* (Kolmogorov-Smirnov two sample test: $d = 0.1434$, $P = 0.5162$) were not significant different.

Weight - length relationships (WLRs)

The weight - length relationships of five species were showed in Table 1. WLRs was estimated as $W = 0.0076TL^{3.0882}$ ($r^2 = 0.8110$, $n = 313$, $sd_{\log TL} = 0.0218$, $sd_{\log W} = 0.0750$, Pauly's t -test = 0.9829, $P > 0.005$, isometric growth) for *E. affinis*, $W = 0.0016TL^{3.4571}$ ($r^2 = 0.8595$, $n = 325$, $sd_{\log TL} = 0.0237$, $sd_{\log W} = 0.0885$, Pauly's t -test = 5.8774, $P < 0.005$, positive allometric growth) for *K. pelamis*, $W = 0.1514TL^{2.1468}$ ($r^2 = 0.7746$, $n = 278$, $sd_{\log TL} = 0.0093$, $sd_{\log W} = 0.0228$, Pauly's t -test = 12.2398, $P < 0.005$, negative allometric growth) for *S. guttatus*, $W = 0.1558TL^{2.1626}$ ($r^2 = 0.8453$, $n = 294$, $sd_{\log TL} = 0.0113$, $sd_{\log W} = 0.0265$, Pauly's t -test = 15.4678, $P < 0.005$, negative allometric growth) for *S. commerson* and $W = 0.1011TL^{2.4606}$ ($r^2 = 0.8387$, $n = 300$, $sd_{\log TL} = 0.0228$, $sd_{\log W} = 0.0613$, Pauly's t -test = 8.6291, $P < 0.005$, negative allometric growth) for *T. tonggol*.

The slope of WLRs regression lines for 5 species except for *E. affinis* were significantly different from the isometric growth curve slope ($b = 3$) ($P < 0.005$). These results showed that *S. guttatus*, *S. commerson* and *T. tonggol* have negative allometric growth characteristics while *K. pelamis* exhibit positive allometric growth.

Von Bertalanffy Growth Parameters

The seasonal and non-seasonal von Bertalanffy growth parameters obtained from the LFDA for each species are summarized in Table 2. The R_n value of the non-seasonal growth curve for each species improved when a seasonal growth curve was fitted (Table 2), suggesting that, at least for our data, *E. affinis*, *K. pelamis*, *S.*



guttatus and *T. tonggol* exhibit a seasonal growth pattern. This was also apparent in Figures (1A, B - 4A, B) where sinusoidal pattern could be observed in the seasonal von Bertalanffy growth curve. On the other hand, the R_n value of the non-seasonal VBG curve improved by 14% for *K. pelamis* (Figure 1B), 27% for *T. tonggol* (Figure 2B), 30% for *E. affinis* (Figure 3B), 21% for *S. guttatus* (Figure 4B), and after fitting a seasonal VBG curve (Table 2).

Stronger seasonal variation of growth was determined for *E. affinis* ($C = 0.990$), *K. pelamis* ($C = 0.950$), *S. guttatus* ($C = 0.980$) and *T. tonggol* ($C = 0.849$). The start of slow growth period was in September ($WP = 0.7$) for *E. affinis*, in early August for *K. pelamis* ($WP = 0.6$), in early January for *S. guttatus* ($WP = 0.0$) and the slow growth period started in May ($WP = 0.4$) for *T. tonggol* (Table 2). Calculated growth performance indices (Φ' ; Table 2) of seasonal growth were a bit greater than it was for non-seasonal.

Discussion

The WLRs are important and have many applications in fish stock assessments, biomass estimations, ecological studies and modeling aquatic ecosystems. The WLRs are also provides valuable information on the habitat where the fish lives, condition, reproduction history, life cycle and the general health of fish species (Froese, 2006; Froese et al., 2011). Our results about the WLRs of five scomberids are not new, but provide current information on the species. In this study, four species showed allometric growth pattern, while one species showed isometric growth.

Jin et al. (2015) reported b values of WLRs for *K. pelamis* when fork length was > 60 cm were significantly less than 3 (negative allometric growth), but when fork length was < 60 cm they were significantly greater than 3 (positive allometric growth). In the present study, b value (3.4571) of *K. pelamis* was estimated as positive allometric growth (length ranged between 52 and 68 cm). In generally, reported b values of *K. pelamis* are similar (allometric growth) to the previous studies (Table 3). The WLRs was also showed that *T. tonggol* exhibited negative allometric growth ($b = 2.4606$, $P < 0.005$). Similar b value of 2.515 for *T. tonggol* (length ranged between 30.0 and 97.9 cm) was reported from Veraval waters (Ghosh et al., 2010). However, contrary to present study, b values for *T. tonggol* were reported by Kaymaram et al. (2013) as $b = 2.848$ (isometric) in the Persian Gulf, Oman Sea and by Abdussamad et al. (2012) as $b = 3.000$ (isometric) in the Indian coasts (Table 4). In the present study, isometric growth pattern was only determined for *E. affinis* ($b = 3.0882$, $P > 0.5$). Similar b value of 3.056 for *E. affinis* (length ranged between 26.0 and 69.9 cm) was reported from Veraval waters (Ghosh et al., 2010). However, contrary to the present study, b values for *E. affinis* were reported from other studies as isometric (Table 5). The WLRs showed that *S. guttatus* exhibited negative allometric growth ($b = 2.1468$, $P < 0.005$). Similarly, negative allometric growth for *S. guttatus* ($b = 2.862$; length ranged between 25 and 70 cm) was reported from Bay of Bengal off Bangladesh coasts (Rashid et al. 2010) and from Palk Bay and Mannar Gulf ($b = 2.860$; length ranged between 36.9 and 70.5 cm) (Devaraj, 1981) (Table 6). The WLRs of the other scombrids fish species, *S. commerson*, was also determined as negative allometric growth ($b = 2.1626$, length range = 45.8 and 51 cm) in the study area. These variations are probably due to factors related to ecosystem and biological condition like spawning time, sexual maturity, feeding behavior and competition for food (Ghosh et al., 2010). The variation in the exponent (b) of the WLRs of fish species could be also affected by geographic



locations, sampling area, seasons, size range and ecological factors such as temperature (Froese, 2006; Froese et al., 2011).

The VBG curve parameters (L_{∞} and K) of different *K. pelamis* stocks were showed great variability (Table 3). In the literature, values of L_{∞} were reported from 60 cm FL in Eastern Atlantic, Cap Vert (Cayré et al., 1986) to 127.8 cm FL in the Western Pacific (Chi and Yang, 1973). On the other hand, values of growth rate parameter (K) of *K. pelamis* stocks were reported between 0.14 year⁻¹ in the Eastern Atlantic South of 10° N latitude (Gaertner et al., 2008) and 2.08 year⁻¹ in the Eastern Atlantic, Senegal (Cayré et al., 1986). This variability may be due to particular characteristics of sampling region such as biotic and abiotic factors, state of the different exploited stocks, different length ranges and differences of growth estimation methods such as otolith, dorsal spine, vertebrae, tagging and length frequency analysis. In the present study, we estimated seasonal VBG curve parameters of *K. pelamis* first time. Estimated L_{∞} (66.8 cm) and K (1.378 year⁻¹) values were between the values specified in the early studies (Table 2) and pointed out that *K. pelamis* was rapid growth species in the study area. The relative amplitude of the seasonal oscillations ($C = 0.950$) was also implied that *K. pelamis* showed strong seasonality in growth and the lowest growth time was corresponding to early August ($WP = 0.6$) offshore water of Pakistan coast in the Arabian Sea. *K. pelamis* mature and spawn round the year in Indian waters with a peak from December to March and a minor one during June – August (Koya et al., 2012). It was also observed that this species spawn almost round the year with one major peak during late winter and a minor peak during the south-west monsoon (Koya et al., 2012). Moreover, *K. pelamis* known as spawn multiple times in areas (e.g. eastern Pacific Ocean) where the surface temperature is equal to or higher than 24°C (Buñag, 1956; Schaefer, 2001). The reproduction of *K. pelamis* in tropical areas is also reported as continuous over the year and a seasonal component in the spawning appears if the studied area is far from the equator (Andrade and Santos, 2004; Andrade and Kinas, 2004). Reproduction time of *K. pelamis* in the tropical western and central Pacific Ocean was reported as evidence of spawning activity throughout the year without a clear seasonal pattern (Ashida et al., 2010). Several biotic and abiotic factors, especially sea water temperature and more of the energy spent for gonadal growth could increase on reproduction activity in the spawning time (mentioned above) probably cause decrease of somatic growth (to early August) of *K. pelamis* in the study area. In addition to these, amount and quality of prey occurrence and seasonal variation food availability in stomach in *K. pelamis* could be attributed mainly to the seasonal variation in growth (Al-Zibdah and Odat, 2007).

In the previous studies (Table 4), L_{∞} values for *T. tonggol* were reported between 58.2 cm FL in Gulf of Thailand, South China Sea (Supongpan and Saikliang, 1987) and 133.7 cm FL in the Sea of Oman (Kaymaram et al., 2013). On the other hand, values of growth rate parameters (K) for *T. tonggol* stocks were reported from 0.18 year⁻¹ in Veraval waters (Ghosh et al., 2010) to 1.44 year⁻¹ in Gulf of Thailand, South China Sea (Supongpan and Saikliang, 1987). The VBG curve parameters recorded in the present study were different from other studies of *T. tonggol* in other geographical region. Namely, the L_{∞} value (71.7 cm) obtained in this study was relatively smaller than in other studies, but the K value (0.916 year⁻¹) obtained in this study was calculated to be higher than the other studies. This is probably due to range of length used to growth calculation for *T. tonggol*. Size range of *T. tonggol* individuals used for growth analyzes in this study was smaller and narrower size range (43 – 69 cm) than in other studies. For example, length ranges and L_{∞} values for *T. tonggol* were reported as length range: 30 – 97.9 cm, $L_{\infty} = 107.4$ cm by Ghosh et al. (2010), length range: 23.8 – 125 cm, $L_{\infty} =$



99.7 cm by Griffiths et al. (2010), length range: 26 – 125 cm, L_{∞} = 133.7 cm by Kaymaram et al. (2013), length range: 23 – 111 cm, L_{∞} = 123.5 cm by Abdussamad et al. (2012).

In the literature, values of L_{∞} for *E. affinis* stocks were reported from 55.1 cm FL in the Thailand Gulf, South China Sea (Supongpan and Saikliang, 1987) to 117.8 cm FL in the Northwestern Hawaiian Islands (Uchiyama, 1980). Similarly, values of growth rate parameter (K) of *E. affinis* stocks were reported between 0.42 year⁻¹ in the Northwestern Hawaiian Islands (Uchiyama, 1980) and 2.23 year⁻¹ in the Thailand Gulf, South China Sea (Supongpan and Saikliang, 1987). In the present study area *E. affinis* growth parameters (L_{∞} = 67.1 cm, K = 1.055 year⁻¹) were found similar to growth parameters values (L_{∞} = 64.6 cm, K = 1.00 year⁻¹) for *E. affinis* in the Indian Ocean (Sulistyaningsih et al., 2014). In addition to these results, some authors were reported different L_{∞} and K values for *E. affinis* from the other geographical areas (Table 5). This could be due to the differences in the environmental parameters, availability and competition for food. Different L_{∞} and K could be associated with variation in fishing intensity and differences in the environmental parameters and availability and competition for food. Moreover, these differences in growth parameters among the geographical area and different sampling years may be due to sampling error and the differences of the methods (such as length based and age based growth estimates) used to calculate the growth parameters. In the literature, there is no research concerning seasonal growth of *E. affinis*. This species showed seasonal variation in growth (C = 0.990) and Rn value of the non-seasonal VBG curve improved by 30% when seasonal VBG growth was applied. The start of slow growth period of *E. affinis* was corresponding to September (WP = 0.7). Al-Zibdah and Odat (2007) reported that food composition of *K. pelamis* and *E. affinis* suggests that the fish compete for the same food items. These are the fish *Atherinomorous lacunosus*, crustacean and molluscs. Prey occurrence in stomachs of two fish species is attributed mainly to the seasonal availability of food and this could be considered one of the causes of seasonal growth of *E. affinis* in the study area. Moreover, this seasonality in growth for *E. affinis* is possibly due to factors related to ecosystem and biological condition like reproductive activities such as gonad maturity stages (Al-Zibdah and Odat, 2007; Ghosh et al., 2010). Namely, the highest percentage of maturing females was observed during September - April off the west coast of Thailand (Yesaki, 1982). Moreover, Yesaki (1994b) reported that there are two spawning seasons for *E. affinis* in the equatorial regions. In the northern hemisphere, the principal spawning season generally occurs during the first half of the year (from October to May) and the secondary spawning season during the latter half. At higher latitudes of the tropical zone, there appears to be only one spawning season near the middle (during June - August) of the year. Besides the aforementioned reasons, another reason for the seasonal growth of *E. affinis* is probably thought to results from differences in different seasonal oceanography parameters such as temperature, current and salinity.

There are two previous studies based on length data have been completed to determine the growth of *S. guttatus* (Table 6). L_{∞} value (L_{∞} = 127.8 cm, K = 0.18 year⁻¹) reported by Devaraj (1981) was much higher than the value (L_{∞} = 73.5 cm, K = 0.6 year⁻¹) reported by Rashid et al. (2010) whereas the K value reported by Devaraj (1981) was much lower. The von Bertalanffy growth parameters obtained this study (L_{∞} = 55.7 cm, K = 1.049 year⁻¹) were lowest L_{∞} value and highest K value than the others (Devaraj, 1981; Rashid et al., 2010). We studied first time to describe of seasonal von Bertalanffy growth parameters for *S. guttatus*. Seasonal variation of growth (C = 0.980) was determined and the start of slow growth period corresponding to early January (WP = 0.0). Moreover, the Rn value of the non-seasonal VBG curves for *S. guttatus* improved by 21% after fitting a



seasonal VBG curve. Seasonal changes in fish growth mostly caused by differences in water temperature and reproductive activities affecting fish feeding mechanism (Bilgin et al., 2012; Bilgin et al., 2013). In the literature, reproduction time was reported during June and July in the Bay of Bengal (Rashid et al., 2010), from January to August, with a peak in April and May off the south coast of India (Devaraj, 1987), from April to July around Rameswaram Island between India and Sri Lanka (Collette and Nauen, 1983). As will be understood from these studies, start of reproduction activity time (between April and July) is not in agreement with the start of slow growth time (winter point =January). So, water temperature, and may be other biotic and abiotic parameters such as current, salinity, food and feeding habits could be considered mostly effecting the seasonal growth of *S. guttatus* in the study area. Moreover, Food and feeding habits of *S. guttatus* was studied by Deveraj (1998) from the northern Gulf of Mannar and Palk Bay along the southeast coast of India between July 1967 and July 1969. Deveraj (1998) reported that *S. guttatus* feeds on a limited number of about five species, of which the sardinella was the most important item of in the food in August and it was absent in the diet in from August through January in 1967 - 68 and in during January in 1968 – 69. The start of slow feeding period of *S. guttatus* was corresponding to more or less January and this period was in agreement with the start of slow growth time of *S. guttatus* ($WP = 0$, January).

References

- Abdussamad, E.M., Said Koya, K.P., Ghosh, S., Rohit, P., Joshi, K.K., Manojkumar, B., Prakasan, D., Kemparaju, S., Elayath, M.N.K., Dhokia, H.K., Sebastine, M. and Bineesh, K.K. 2012. Fishery, biology and population characteristics of longtail tuna, *Thunnus tonggol* (Bleeker, 1851) caught along the Indian coast. Indian Journal of Fisheries, 59 (2): 7-16.
- Adams, J.L. and Kerstetter, D.W. 2014. Age and growth of three coastal- pelagic tunas (Actinopterygii: Perciformes:Scombridae) in the Florida straits, USA: Blackfin tuna, *Thunnus atlanticus*, Little tunny, *Euthynnus alletteratus*, and Skipjack tuna, *Katsuwonus pelamis*. Acta Ichthyologica et Piscatoria, 44 (3): 201 - 211. doi:10.3750/AIP2014.44.3.04.
- Ahmed, Q., Yousuf, F., Sarfraz, M., Ali, Q.M., Balkhour, M., Safi, S.Z. and Ashraf, M.A. 2015. *Euthynnus affinis* (little tuna): fishery, bionomics, seasonal elemental variations, health risk assessment and conservational management. Frontiers Life Science, 8 (1): 71 – 96. doi:10.1080/21553769.2014.961617.
- Al-Zibdah, M. and Odat, N. 2007. Fishery status, growth, reproduction biology and feeding habit of two Scombrid fish from the Gulf of Aquaba, Red Sea. Lebanese Science Journal, 8 (2): 3-20.
- Andrade, H.A. and Campos, R.O. 2002. Allometry coefficient variations of the length - weight relationship of skipjack tuna (*Katsuwonus pelamis*) caught in the southwest South Atlantic. Fisheries Research, 55: 307- 312.
- Andrade, H.A. and Santos, J.A.T. 2004. Seasonal trends in the recruitment of skipjack tuna (*Katsuwonus pelamis*) to the fishing ground in the southwest Atlantic. Fisheries Research, 66: 185 – 194.
- Andrade, H.A. and Kinan, P.G. 2004. Estimation of birthdates and catch-at-age using length frequency analysis (LFA), with application for skipjack tuna (*Katsuwonus pelamis*) caught in the Southwest Atlantic. ICES Journal of Marine Science, 61: 798 – 811.
- Ashida, H., Tanabe, T., Satoh, K., Fukui, A., Tanaka, S. and Suzuki, N. 2010. Reproductive biology of male skipjack tuna *Katsuwonus pelamis* (Linnaeus) in the tropical western and central Pacific Ocean. Fisheries Science, 76: 785 – 793. doi: 10.1007/s12562-010-0282-y
- Avşar, D., 2005. Balıkçılık biyolojisi ve popülasyon dinamiği. Nobel Kitabevi, Adana.



- Bilgin, S., Bal, H. and Taşçı, B. 2012. Length based growth estimates and reproduction biology of whiting, *Merlangius merlangus euxinus* (Nordman, 1840) in the Southeast Black Sea. Turkish Journal of Fisheries and Aquatic Sciences, 12: 871 - 881. doi: 10.4194/1303-2712-v12_4_15.
- Bilgin, S., Taşçı, B. and Bal, H. 2013. Sexual seasonal growth of the European anchovy (*Engraulis encrasicolus*) caught by midwater trawl and purse seine in the southern Black Sea. Journal of the Marine Biological Association of the United Kingdom, 93(2): 333 - 339. doi:10.1017/S0025315412000732.
- Buñag, D.M. 1956. Spawning habits of some Philippine tuna based on diameter measurements of the ovarian ova. The Philipp J. Fish., 4: 145-177.
- Cayré, P., Diouf, T. and Fonteneau, A. 1986. Analyse des données de marquages et recaptures de listao (*Katsuwonus pelamis*) réalisés par le Sénégal et la République du Cap-Vert. In proceedings of the ICCAT Conference on the international skipjack year program, edited by PEK Symons, PM Miyake, GT Sakagawa. Madrid, Spain. ICCAT, 309–316.
- Chi, K.S. and Yang, R.T. 1973. Age and growth of skipjack tuna in the waters around the southern part of Taiwan. [Summ. in Chin.] Nat Taiwan Univ Sci Rep Acta Oceanogr Taiwanica, 3: 199 - 221.
- Chur, V.N. and Zharov, V.L. 1983. Determination of age and growth of the skipjack tuna, *Katsuwonus pelamis* (Scombridae) from the southern part of the Gulf of Guinea. Journal of Ichthyology, 23 (3): 53–67.
- Collette, B.B. and Nauen, C.E. 1983. FAO Species Catalogue. Vol. 2. Scombrids of the world. An annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date. Rome: FAO. FAO Fish Synop., 125 (2):137 p.
- Collette, B., Acero, A., Amorim, A.F., Boustany, A., Canales Ramirez, C., Cardenas, G., Carpenter, K.E., de Oliveira Leite, Jr., N., Di Natale, A., Fox, W., Fredou, F.L., Graves, J., Guzman-Mora, A., Viera Hazin, F.H., Juan Jorda, M., Kada, O., Minte Vera, C., Miyabe, N., Montano Cruz, R., Nelson, R., Oxenford, H., Salas, E., Schaefer, K., Serra, R., Sun, C., Teixeira Lessa, R.P., Pires Ferreira Travassos, P.E., Uozumi, Y. and Yanez, E. 2011a. *Katsuwonus pelamis*. The IUCN Red List of Threatened Species. Version 2014.3. <www.iucnredlist.org>. Downloaded on 13 April 2015.
- Collette, B., Di Natale, A., Fox, W., Juan Jorda, M., Miyabe, N., Nelson, R., Sun, C. and Uozumi, Y. 2011b. *Thunnus tonggol*. The IUCN Red List of Threatened Species. Version 2014.3. <www.iucnredlist.org>. Downloaded on 13 April 2015.
- Collette, B., Chang, S.K., Fox, W., Juan Jorda, M., Miyabe, N., Nelson, R. and Uozumi, Y. 2011c. *Euthynnus affinis*. The IUCN Red List of Threatened Species. Version 2014.3. <www.iucnredlist.org>. Downloaded on 13 April 2015.
- Collette, B., Di Natale, A., Fox, W., Juan Jorda, M. and Nelson, R. 2011d. *Scomberomorus guttatus*. The IUCN Red List of Threatened Species. Version 2014.3. <www.iucnredlist.org>. Downloaded on 13 April 2015.
- Collette, B., Chang, S.K., Di Natale, A., Fox, W., Juan Jorda, M., Miyabe, N. and Nelson, R. 2011e. *Scomberomorus commerson*. The IUCN Red List of Threatened Species. Version 2014.3. <www.iucnredlist.org>. Downloaded on 13 April 2015.
- Devaraj, M. 1981. Age and growth of three species of seer fishes *Scomberomorus commerson*, *S. guttatus* and *S. lineolatus*. Indian Journal of Fisheries, 28(1/2): 104-127.
- Devaraj, M. 1987. Maturity, spawning and fecundity of the spotted seer, *Scomberomorus guttatus*, in the Gulf of Mannar and Palk Bay. Indian Journal of Fisheries, 34: 48 – 77.
- Devaraj, M. 1998. Food and feeding habits of the spotted seer, *Scomberomorus guttatus* (Bloch and Schneider), in the Gulf of Mannar and Palk Bay Journal of the Marine Biological Association of India, 40: 105-124.
- Dutta, H., 1994. Growth in fishes. Gerontology, 40: 97-112.
- FAO, 2015. <http://www.fao.org/fishery/statistics/global-capture-production/en>. Download November 20, 2015.



- Froese R. 2006. Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22: 241–253.
- Froese, R., Tsikliras A.C. and Stergiou, K.I. 2011. Editorial note on weight-length relations of fishes. *Acta Ichthyologica et Piscatoria*, 41: 261-263.
- Froese, R. and Pauly, D. 2015. Editor FishBase. World Wide Web electronic publication. www.fishbase.org, version (02/2015).
- Gaertner, D, de Molina, A.D., Ariz, J., Pianet, R. and Hallier, J.P. 2008. Variability of the growth parameters of the skipjack tuna (*Katsuwonus pelamis*) among areas in the eastern Atlantic: analysis from tagging data within a meta-analysis approach. *Aquat. Living Resour.*, 21: 349 – 356. doi:10.1051/alr:2008049.
- Garbin, T. and Castello, J.P. 2014. Changes in population structure and growth of skipjack tuna, *Katsuwonus pelamis* during 30 years of exploitation in the southwestern Atlantic. *Latin American Journal of Aquatic Research*, 42(3): 434-546. doi: 103856/vol42-issue3-fulltext-13.
- Ghosh, S., Pillai, N.G.K. and Dhokia, H.K. 2010. Fishery, population characteristics and yield estimates of coastal tunas at Veraval. *Indian Journal of Fisheries*, 57 (2): 7-13.
- Griffiths, S.P., Fry, G.C., Manson, F.J. and Lou, D.C., 2010. Age and growth of longtail tuna (*Thunnus tonggol*) in tropical and temperate waters of the central Indo-Pacific. *ICES Journal of Marine Science*, 67 (1): 125-134.
- Hallier, J.P. and Gaertner, D. 2006. Estimated growth rate of the skipjack tuna (*Katsuwonus pelamis*) from tagging surveys conducted in the Senegalese area (1996-1999) within a meta-analysis framework. *Col Vol Sci Pap ICCAT.*, 59 (2): 411-420.
- Henderson, P.A. and Seaby, R.M. 2006. Growth II. Pisces Conservation Ltd., Lymington, England.
- Hornby, C., Khan, M.M., Zyllich, K. and Zeller, D., 2014. Reconstruction of Pakistan's marine fisheries catches 1950-2010. FAO report, September 6th, 2013.
- Jin, S., Yan, X., Zhang, H. and Fan, W. 2015. Weight-length relationships and Fulton's condition factors of skipjack tuna (*Katsuwonus pelamis*) in the western and central Pacific Ocean. *Peer J.*, 12:3: e758 <https://dx.doi.org/10.7717/peerj.758>.
- Kaymaram, F. and Darvishi, M. 2012. Growth and mortality parameters of *Euthynnus affinis* in the northern part of the Persian Gulf and Oman Sea. *Indian Ocean Tuna Commission (IOTC), Working Party on Neritic Tunas (WPNT), IOTC-2012-WPNT02-14: 1-14.*
- Kaymaram, F., Darvishi, M., Behzadi, S. and Ghasemi, S. 2013. Population dynamic parameters of *Thunnus tonggol* (Bleeker, 1851) in the Persian Gulf and Oman Sea. *Iranian Journal of Fisheries Sciences*, 12 (4): 855 – 863.
- Khan, M.Z. 2004. Age and growth, mortality and stock assessment of *Euthynnus affinis* (Cantor) from Maharashtra waters. *Indian Journal of Fisheries*, 51 (2): 209 – 213.
- Khan, M.M., 2012. Tuna situation analysis-Pakistan. World Wildlife Fund (WWF), Karachi. 43 p.
- Kirkwood, G.P, Aukland, R. and Zara, S.J. 2003. Software: Length Frequency Distribution Analysis (LFDA) Version 5.0. Marine Resources Assessment Group (MRAG) Ltd., London, UK.
- Koya, K.P.S., Joshi, K.K., Abdussamad, E.M., Rohit, P., Sivadas, M., Kuriakose, S., Ghosh, S., Koya, M., Dhokia, H.K., Prakasan, D., Kunhi Koya, V.A. and Sebastine, M. 2012. Fishery, biology and stock structure of skipjack tuna, *Katsuwonus pelamis* (Linnaeus, 1758) exploited from Indian waters. *Indian Journal of Fisheries*, 59 (2): 39- 47.
- Moazzam, M. and Nawaz, R., 2015. By-catch of tuna gillnet fisheries of Pakistan: A serious threat to non-target, endangered and threatened species. *The Marine Biological Association of India*, 56 (1), 85-90, doi: 10.6024/jmbai.2014.56.1.01750s-13.
- Pauly, D. 1984. Fish population dynamics in tropical water: a manual for use with programmable calculators. *ICLARM Studies and Reviews* 8.



- Pauly, D. and Munro, J.L. 1984. Once more on the comparison of growth in fish and invertebrates. ICLARM Fishbyte, 2: 21.
- Prabhakar, A. and Dudley, R.G. 1989. Age, growth and mortality rates of longtail tuna *Thunnus tonggol* (Bleeker) in Omani waters based on length data. Indo-Pac. Tuna Dev. Mgt. Programme, IPTP/89/GEN, 16: 90 - 96.
- Rashid, H., Mustafa, M.G. and Dewan, S. 2010. Population dynamics and the management of the Indo-Pacific king mackerel *Scomberomorus guttatus* from the upper Bay of Bengal off Bangladesh coast. The Bangladesh Veterinarian, 27 (2): 82 – 90.
- Rohit, P., Chellappan, A., Abdussamad, E.M., Joshi, K.K., Koya, K.P.S., Sivadas, M., Ghosh, S., Rathinam, A.M.M., Kemparaju, S., Dhokia, H.K., Prakasan, D. and Beni, N. 2012. Fishery and bionomics of the little tuna, *Euthynnus affinis* (Cantor, 1849) exploited from Indian waters. Indian Journal of Fisheries, 59 (3): 33 - 42.
- Schaefer, K.M. 2001. Assessment of skipjack tuna (*Katsuwonus pelamis*) spawning activity in the eastern Pacific Ocean. Fishery Bulletin, 99: 343 – 350.
- Sibert, J.R., Kearney, R.E. and Lawson, T.A. 1983. Variation in growth increments of tagged skipjack *Katsuwonus pelamis*. South Pacific Commission, Nouméa, New Caledonia, Tuna and Billfish Assesst Program Tech Report No: 10, 43 pp.
- Siddeek, M.S.M. 1995. Review of fisheries biology of *Scomberomorus* and *Acanthocybium* species in the western Indian Ocean (FAO Area 51). Working Group on Pelagics, Gulfs Committee for Fisheries Management and Development. Working Paper WGP 95/2. 32 p.
- Somers, I.F. 1988. On a seasonally oscillating growth function. Fishbyte, 6(1): 8-11.
- Sulistyaningsih, R.K., Jatmiko, I. and Wujdi, A. 2014. Length frequency distribution and population parameters of Kawakawa (*Euthynnus affinis* - Cantor, 1849) caught by purse seine in the Indian Ocean (a case study in Northwest Sumatra IFMA 572). Indian Ocean Tuna Commission (IOTC), Working Party on Neritic Tunas (WPNT), IOTC-2014-WPNT04-20, 1-14 p. (available: http://www.iotc.org/sites/default/files/documents/2014/06/IOTC-2014-WPNT04-20_-_Kawakawa_length_freq_Indonesia.pdf).
- Supongpan, S. and Saikliang, P. 1987. Fisheries status of tuna purse seiners (using sonar) in the Gulf of Thailand. Rep Mar Fish Div Dep Fish Bangkok., (3): 78 p.
- Taghavi, M.S.A., Hashemi, S.A. and Kochanian, P. 2010. Population biology and assessment of Kawakawa (*Euthynnus affinis*) in Coastal Waters of the Persian Gulf and Sea of Oman (Hormozgan Province). Iranian Journal of Fisheries Sciences, 9 (2): 315 – 326.
- Uchiyama, J.H. 1980. Survey of the pelagic fishes of the northwestern Hawaiian Islands. In Status of resource investigations in the northwestern Hawaiian Islands, edited by R.W. Grigg and R.T. Pfund. Univ. Hawaii Sea Grant College Prog., Honolulu, Hawaii, pp. 251-63.
- Uchiyama, J.H. and Struhsaker, P. 1981. Age and growth of skipjack tuna, *Katsuwonus pelamis*, and yellowfin tuna, *Thunnus albacares*, as indicated by daily growth increments of sagittae. Fishery Bulletin, 79 (1): 151-162.
- van der Land, J., Costello, M.J., Zavodnik, D., Santos, R.S., Porteiro, F.M., Bailly, N., Eschmeyer, W.N. and Froese, R. 2001. Pisces, in: Costello, M.J. et al. (Ed.) (2001). European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification. Collec Patrimoines Nat., 50: 357-374.
- von Bertalanffy, L. 1938. A quantitative theory of organic growth (Inquires on growth laws II). Human Biology, 10: 181 - 213.
- von Bertalanffy, L. 1951. Metabolic types and growth types. The American Naturalist LXXXV 821, 111-117.
- Wankowski, J.W.J. 1981. Estimated growth of surface – schooling skipjack tuna, *Katsuwonus pelamis*, and yellowfin tuna, *Thunnus albacares*, from the Papua New Guinea region. Fishery Bulletin, 79 (3): 517- 532.
- Wild, A. and Hampton, J. 1994. A review of the biology and fisheries for skipjack tuna, *Katsuwonus pelamis*, in the Pacific Ocean. FAO Fish Techn Pap., 336 (2): 1-51.



- Wilson, M.A. 1981. The biology, ecology and exploitation of longtail tuna, *Thunnus tonggol* (Bleeker) in Oceania. MSc. thesis. School of Biological Sciences, Macquarie University, Sydney, 195 p.
- Yesaki, M. 1982. Thailand. Biological and environmental observations. A report prepared for the Pole – and - Line Tuna Fishing in Southern Thailand Project. FAO. FI:DP/THA/77/008:46 p.
- Yesaki, M. 1989. Estimates of age and growth of kawakawa (*Euthynnus affinis*), longtail tuna (*Thunnus tonggol*) and frigate tuna (*Auxis thazard*) from the Gulf of Thailand based on length data. Indo-Pac.Tuna Dev.Mgt.Programme, IPTP/89/GEN/17:94-108.
- Yesaki, M. 1994a. A review of the biology and fisheries for longtail tuna (*Thunnus tonggol*), in the Indo-Pacific region. FAO Fish Techn Pap., 336 (2): 370 -387.
- Yesaki, M. 1994b. A review of the biology and fisheries for kawakawa (*Euthynnus affinis*), in the Indo-Pacific region. FAO Fish Techn Pap., 336 (2): 388 -408.

Accepted Manuscript

**Table 1.** Weight – length relationship parameters and growth types of five species. ns: non-significant, a is the intercept, and b is the slope of the regression line.

Species	a	b	r^2	n	Pauly's t -test	P	Growth type
<i>K. pelamis</i>	0.0016	3.4571	0.8595	325	5.8774	$P < 0.005$	positive allometric
<i>T. tonggol</i>	0.1011	2.4606	0.8387	300	8.6291	$P < 0.005$	negative allometric
<i>E. affinis</i>	0.0076	3.0882	0.8110	313	0.9829	ns	isometric
<i>S. guttatus</i>	0.1514	2.1468	0.7746	278	12.2398	$P < 0.005$	negative allometric
<i>S. commerson</i>	0.1558	2.1626	0.8453	294	15.4678	$P < 0.005$	negative allometric

Table 2. Seasonal and non-seasonal von Bertalanffy growth function parameters. L_∞ is the asymptotic length to which the fish growth, K is the growth-rate parameter, t_0 is the nominal age at which the length is zero, C is the relative amplitude of the seasonal oscillations, t_s is the phase of the seasonal oscillations, WP is winter point, Rn is the goodness of fit, ϕ' is growth performance index.

Growth curve	Species	Parameters					WP	Rn	ϕ'
		L_∞ (cm)	K (year ⁻¹)	t_0 (year)	t_s	C			
Hoening seasonal	<i>K. pelamis</i>	66.8	1.378	-0.05	0.11	0.950	0.6	0.503	3.789
	<i>T. tonggol</i>	71.7	0.916	-0.99	-0.06	0.849	0.4	0.679	3.673
	<i>E. affinis</i>	67.1	1.055	-0.84	0.16	0.990	0.7	0.633	3.677
	<i>S. guttatus</i>	55.7	1.049	-0.42	-0.50	0.980	0.0	0.779	3.512
Non seasonal	<i>K. pelamis</i>	66.3	1.223	-0.45				0.443	3.730
	<i>T. tonggol</i>	69.9	0.934	-0.09				0.536	3.659
	<i>E. affinis</i>	66.3	0.927	-0.23				0.486	3.610
	<i>S. guttatus</i>	55.8	1.081	-0.27				0.644	3.527



Table 3. von Bertalanffy growth parameters, length range, b values and growth performance index (ϕ') of *Katsuwonus pelamis*. FL: fork length (cm), TL: total length (cm), MPA: Modal progression analyzes, ELEFAN: Electronic length frequency analyzes,* cited by Wild and Hampton (1994). ϕ' calculated by $2*\log(L_{\infty})+\log(K)$ equation.

Species	L_{∞} (cm)	K (year ⁻¹)	t_0 (year)	ϕ'	b	$L_{\min-max}$ (cm)	Methods	Area	References	
<i>K. pelamis</i>	TL				2.972	56-75		Aqaba Gulf, Red Sea	Al-Zibdah and Odat 2007	
<i>K. pelamis</i>	FL				3.293	35-85		South Atlantic	Andrade and Campos 2002	
<i>K. pelamis</i>	FL	66.9	0.24	-3.80	3.03	3.430	47-74	Otolith	Western Atlantic, S. Brazil, in 2004	Garbin and Castello 2014
<i>K. pelamis</i>	FL	85.4	0.15	-3.90	3.04			Otolith	Western Atlantic, S. Brazil, in 2007	Garbin and Castello 2014
<i>K. pelamis</i>	FL	72.5	0.33	-1.20	3.24	3.440	49-62	Otolith	Western Atlantic, S. Brazil, in 2008	Garbin and Castello 2014
<i>K. pelamis</i>	FL	112.8	0.24	-1.70	3.48		29-78	Otolith	South Florida waters (USA)	Adams and Kerstetter 2014
<i>K. pelamis</i>	FL	102.0	0.55	-0.02	3.76		30-82	Otolith	Central Pacific	Uchiyama and Struhsaker 1981
<i>K. pelamis</i>	FL	92.5	0.16	-2.90	3.14			Bhattacharya	Western Atlantic, S. Brazil,	Garbin and Castello 2014
<i>K. pelamis</i>	FL	74.8	0.51		3.46		30-74	MPA	Central Pacific, Papua New Guine	Wankowski 1981
<i>K. pelamis</i>	FL	92.0	0.50		3.63	3.147	12-82	ELEFAN	Indian waters	Koya et al. 2012
<i>K. pelamis</i>	FL	97.3	0.25		3.38			Tagging	Senegalese, north of Mauritania	Hallier and Gaertner 2006
<i>K. pelamis</i>	FL	61.3	1.25		3.67			Tagging	West Pacific	Sibert et al. 1983
<i>K. pelamis</i>	FL	75.5	0.77		3.64			Tagging	East Pacific	Sibert et al. 1983
<i>K. pelamis</i>	FL	112.3	0.14		3.25		42-60	Tagging	Eastern Atlantic Sout of 10°N lat.	Gaertner et al., 2008
<i>K. pelamis</i>	FL	89.4	0.38		3.48		36-62	Tagging	Eastern Atlantic North of 10°N lat.	Gaertner et al. 2008
<i>K. pelamis</i>	FL	62.0	2.08		3.90			Tagging	Eastern Atlantic, Senegal	Cayré et al. 1986*
<i>K. pelamis</i>	FL	60.0	1.54		3.74			Tagging	Eastern Atlantic, Cap Vert	Cayré et al. 1986*
<i>K. pelamis</i>	FL	86.7	0.31	-0.32	3.36		35-74	Dorsal spine	Atlantic, Guine Gulf	Chur and Zharov 1983*
<i>K. pelamis</i>	FL	103.6	0.30	-0.02	3.51		27-65	Vertebrae	Western Pacific	Chi and Yang 1973*



Table 4. von Bertalanffy growth parameters, length range and b values and growth performance index (ϕ') of *Thunnus tonggol*. FL: fork length, TL: total length, LF: length frequency, ELEFAN: Electronic length frequency analyzes, MPA: Modal progression analyzes. *: cited by Yesaki (1994a). ϕ' calculated by $2*\log(L_{\infty})+\log(K)$ equation.

Species		L_{∞} (cm)	K (year ⁻¹)	t_0 (year)	ϕ'	b	$L_{\min-\max}$ (cm)	Methods	Area	References
<i>T. tonggol</i>	FL	107.4	0.18	-0.07	3.32	2.515	30-97.9	ELEFAN	Veraval, Arabian Sea	Ghosh et al. 2010
<i>T. tonggol</i>	FL	99.7	0.23	-1.50	3.36	2.826	23.8-125	Otolith	Australia, Carpentaria Gulf - Iluka	Griffiths et al. 2010
<i>T. tonggol</i>	FL	133.7	0.35		3.80	2.840	26-125	ELEFAN	Persian Gulf, Oman Sea	Kaymaram et al. 2013
<i>T. tonggol</i>	FL	123.5	0.51	-0.03	3.89	3.000	23-111	ELEFAN	Indian coast	Abdussamad et al. 2012
<i>T. tonggol</i>	FL	108.0	0.55		3.81			MPA	Thailand Gulf, South China Sea	Yesaki 1989*
<i>T. tonggol</i>	FL	58.2	1.44		3.69			MPA	Thailand Gulf, South China Sea	Supongpan and Saikliang 1987*
<i>T. tonggol</i>	FL	133.6	0.23		3.61			ELEFAN	Oman	Prabhakar and Dubley 1989*
<i>T. tonggol</i>	FL	131.8	0.40	-0.04	3.84		45-111	Otolith	Papua Gulf	Wilson 1981*

**Table 5.** von Bertalanffy growth parameters, length range and b values and growth performance index (ϕ') of *Euthynnus affinis*. FL: fork length (cm), TL: total length (cm), LF: length frequency, ELEFAN: Electronic length frequency analyzes, MPA: Modal progression analyzes. * cited by Yesaki (1994b). ϕ' calculated by $2*\log(L_{\infty})+\log(K)$ equation.

Species	L_{∞} (cm)	K (year ⁻¹)	t_0 (year)	ϕ'	b	$L_{\min-\max}$ (cm)	Methods	Area	References
<i>E. affinis</i>	FL 81.7	0.79		3.72	2.786	26-73	ELEFAN	Maharashtra	Khan 2004
<i>E. affinis</i>	FL 87.7	0.51	-0.23	3.59		41-73	ELEFAN	Persian Gulf and Sea of Oman	Taghavi et al. 2010
<i>E. affinis</i>	FL 81.9	0.56	0.03	3.57	2.889	48-72	ELEFAN	Karachi, Arabian Sea	Ahmed et al. 2015
<i>E. affinis</i>	FL 64.6	1.00	-0.13	3.62		23.5-61.5	ELEFAN	Indian Ocean, Northwest Sumatera	Sulistyaningsih et al. 2014
<i>E. affinis</i>	FL 81.9	0.56	0.03	3.57	2.889	14-80	ELEFAN	Indian waters	Rohit et al. 2012
<i>E. affinis</i>	FL 72.5	0.56	-0.03	3.47	3.056	26-69.9	ELEFAN	Veraval, Arabian Sea	Ghosh et al. 2010
<i>E. affinis</i>	FL 95.1	0.67		3.78	2.870	28-88	ELEFAN	Persian Gulf and Sea of Oman	Kaymaram and Darvishi 2012
<i>E. affinis</i>	FL 55.1	2.23	-0.02	3.83			MPA	Thailand Gulf, South China Sea	Supongpan and Saikliang 1987*
<i>E. affinis</i>	FL 76.0	0.96		3.74			MPA	Thailand Gulf, South China Sea	Yesaki 1989*
<i>E. affinis</i>	FL 117.8	0.42	-0.03	3.77			Otolith	Northwestern Hawaiian Islands	Uchiyama 1980*
<i>E. affinis</i>	TL				3.140	46-64		Aqaba Gulf, Red Sea	Al-Zibdah and Odat 2007

Table 6. von Bertalanffy growth parameters, length range and b values and growth performance index (ϕ') of *Scomberomorus guttatus*. TL: total length, ELEFAN: Electronic length frequency analyzes. ϕ' calculated by $2*\log(L_{\infty})+\log(K)$ equation.

Species	L_{∞} (cm)	K (year ⁻¹)	t_0 (year)	ϕ'	b	$L_{\min-\max}$ (cm)	Methods	Area	References
<i>S. guttatus</i>	TL 127.8	0.18	-0.47	3.47	2.860	36.9-70.5	Petersen	Palk Bay, Mannar Gulf	Devaraj 1981
<i>S. guttatus</i>	TL 73.5	0.60		3.51	2.862	25-70	ELEFAN	Bay of Bengal off Bangladesh coast	Rashid et al. 2010




Figure 1. Non-seasonal (A) and Hoenig seasonal (B) von Bertalanffy Growth Function (VBGF) curves of *Katsuwonus pelamis*.




Figure 2. Non-seasonal (A) and seasonal (B) von Bertalanffy Growth Function (VBGF) curves of *Thunnus tonggol*.




Figure 3. Non-seasonal (A) and Hoenig seasonal (B) von Bertalanffy Growth Function (VBGF) curves of *Euthynus affinis*.




Figure 4. Non-seasonal (A) and Hoenig seasonal (B) von Bertalanffy Growth Function (VBGF) curves of *Scomberomorus guttatus*.