



## Prevalence of *Marteilia* spp. in Thirteen Shellfish Species Collected from China Coast

Liji Xie<sup>1</sup>, Zhixun Xie<sup>1,\*</sup>

<sup>1</sup>Guangxi Veterinary Research Institute, Department of Biotechnology, Guangxi Key Laboratory of Veterinary Biotechnology, 51 You Ai North Road, Nanning, Guangxi 530001, China.

\* Corresponding Author: Tel.: +867713120371;  
E-mail: xiezhixun@126.com

Received 04 February 2017  
Accepted 19 September 2017

### Abstract

*Marteilia* spp. is listed as a notifiable parasite by the Office International des Epizooties (OIE), and cause physiological disorders and eventually death of the shellfish. Little is known about the prevalence of *Marteilia* spp. in shellfish of China. Therefore we investigated the prevalence of *Marteilia* spp. in 13 species of shellfish collected in the coastal areas of China between 2006 to 2012. A total of 11,581 individual shellfish comprising 13 different species was collected from seven bay areas in China. The shellfish was tested for the prevalence of *Marteilia* spp. following a PCR protocol recommended by OIE. *Marteilia* spp. was found in the shellfish samples (*Crassostrea rivularis*, *Crassostrea gigas* and *Crassostrea angulata*), *Meretrix meretrix*, *Chlamys nobilis*, *Ruditapes philippinarum*, *Perna viridis*, *Tegillarca granosa*, and *Sinonovacula constricta*. The detection rates of *Marteilia* spp. by PCR method ranged from 0% to 6.67%, with *R. philippinarum* (6.67%) showing the highest detection rate. *Marteilia* spp. sequence of the individual positive shellfish had a high sequence homology (97.4% to 99.6%) with the sequences of other *Marteilia* type "M". Mixed positive of *Marteilia* spp. with other protozoan species (*Haplosporidium nelsoni* and *Perkinsus* spp.) was a common phenomenon (26.28%).

**Keywords:** Shellfish, *Marteilia* spp., PCR detection, Chinese coast.

### Introduction

In China, the total area of 1,409,000 hectares was used for an annual shellfish production of 12,666,500 tons in the year of 2011, which is up to 60% of the total global output (Ministry of Commerce of the People's Republic of China, <http://policy.mofcom.gov.cn/export/shellfish/c1.action>). Oysters are the main shellfish product generated in China. A breeding area of more than 116,000 hectares is employed, and more than 3,600,000 tons are produced annually in China, representing approximately 80% of the global oyster output. In recent years, the annual production of shellfish has been declining due to environmental changes and the occurrence of diseases. *Marteilia* spp. infection poses a serious threat to the coastal ecosystem and the supply of shellfish worldwide (Berthe, Le Roux, Adlard, & Figueras, 2004; Villalba, Mourelle, Carballal, & Carmen Lopez, 1993).

*Marteilia* spp. is listed as a notifiable parasite by the Office International des Epizooties (OIE). *Marteilia* spp. is a protozoan parasite resulting in physiological disorders and eventually death of the animal (Feist, Hine, Bateman, Stentiford, &

Longshaw, 2009; Berthe *et al.*, 2000). This protozoan parasite is distributed over a wide range of areas, including Oceania and Europe around Albania, Croatia, France, Greece, Italy, Morocco, Portugal, Spain, Sweden, Tunisia, China and the United Kingdom (Wang, Lu, & Liang, 2012; Lopez-Flores *et al.*, 2008a; Audemard *et al.*, 2004; Kleeman, Le, Berthe, & Adlard, 2002). *Marteilia* spp. infection has been identified in many types of mollusks throughout the world, including oyster species (Villalba *et al.*, 1993), mussel species (Wang, Lu, & Liang, 2012; Le Roux *et al.*, 2001) and clam species (Lopez-Flores *et al.*, 2008b).

However, little has been known about the prevalence of *Marteilia* spp. in shellfish in the coastal areas of China. In this present study, the prevalence of *Marteilia* spp. in 13 shellfish species (*Crassostrea rivularis*, *Crassostrea gigas*, *Crassostrea angulata*, *Meretrix meretrix*, *Chlamys nobilis*, *Ruditapes philippinarum*, *Perna viridis*, *Tegillarca granosa*, *Sinonovacula constricta*, *Haliotis discus hannai*, *Paphia undulate*, *Tapes dorsatus* and *Coelomacra antiquate*) cultured in seven bay areas in China were investigated using a PCR protocol described by Le Roux *et al.* (2001) and this PCR method is also

recommended by OIE.

*C. rivularis* is the major oyster species in Zhanjiang, Haikou and Qin Zhou, and *C. gigas* is the major oyster species in Dalian and Qingdao, while *C. angulata* is the major oyster species in Xiamen. We therefore analyzed the prevalence of *Marteilia* spp. in these three different oyster species (*C. rivularis*, *C. gigas* and *C. angulata*) sampled from various locations (Dalian, Qingdao, Wenzhou, Xiamen, Zhanjiang, Haikou and Qin Zhou) and in different seasons (spring, summer, autumn and winter).

## Materials and Methods

### Sample Collection

Samples of the 13 shellfish species were collected between 2006 and 2012 from seven bay areas in China: Dalian (Bohai Sea), Qingdao (Yellow Sea), Wenzhou (East China Sea), Xiamen (East China Sea), Zhanjiang (South China Sea), Haikou (South China Sea) and Qin Zhou (South China Sea) bays (Figure 1). A total of 11,581 shellfish samples were collected for diagnosis, and their specific details are shown in Table 1. A total of 5,609 oyster samples were collected. To find out the differentiation of the prevalence of *Marteilia* spp. among different oyster species, at different locations and in different seasons, these 5,609 oyster samples were collected from different oyster species at different locations within four seasons. In terms of oyster species, these 5,609 oyster samples include 3,397 *C. rivularis*, 1,351 *C. gigas* and 861 *C. angulata*. Regarding to the locations, these 5,609 oyster samples include 694 oyster (*C. gigas*) of Dalian, 657 oyster (*C. gigas*) of Qingdao, 597 oyster (122 *C. angulata* + 475 *C. rivularis*) of Wenzhou, 739 oyster (*C. angulata*) of Xiamen, 540 oyster (*C. rivularis*) of Zhanjiang, 454 oyster (*C. rivularis*) of Haikou and 1,928 oyster (*C. rivularis*) of Qin Zhou. And when counted by different seasons, these 5,609 oyster samples include 1,432 in spring, 1,326 in summer, 1,384 in autumn and 1,467 in winter.

### DNA Extraction

Genomic DNA was extracted from the excised tissue samples (gill and digestive gland together) using EasyPure® Marine Animal Genomic DNA Kit as described by Xie *et al.*, (2013). The tissue samples were frozen in liquid nitrogen for 3 min and then ground into powder. The powder was then digested overnight with proteinase K at 56°C, and the genomic DNA was extracted.

### PCR, Cloning, and Sequencing

PCR is performed with primers targeting the ITS1 (internal transcribed spacer) region (Pr4: 5'-CCG-CAC-ACG-TTC-TTC-ACT-CC-3' and Pr5: 5'-CTC-GCG-AGT-TTC-GAC-AGA-CG-3'), which

were also the OIE recommended primers and expected to amplify a 412 base pairs (bp) DNA fragment. Deionized, nuclease-free water was used as a negative control. The PCR reaction solution consisted of 15 µl of 2×PCR mix<sup>a</sup>, 0.2 µM of each primer, and 3 µl of the purified genomic template DNA. A denaturation step at 94°C for 10 min was followed by 35 cycles consisting of a denaturation step at 94°C for 1 min, an annealing step at 55°C for 1 min and an extension step at 72 °C for 1 min. After the final extension at 72°C for 10 min, PCR products were analyzed by 1% agarose gel electrophoresis followed by ethidium bromide staining. The results were visualized under UV light.

Individual positive samples from *C. rivularis*, *C. nobilis*, *R. philippinarum*, *P. viridis*, *T. granosa* and *S. constricta* were selected and again amplified under the conditions similar to those of the screening PCR, except for the composition of the PCR mixture. The total volume of each reaction was increased to 50µl, including 25µl of 2×PCR mix, 0.3 µM of each primer, and 5 µl of the purified genomic DNA as a template. The amplified PCR products were gel-extracted using a Gel Extraction Kit, and were ligated into pMD18-T cloning vector, respectively. Plasmids with the inserted DNA fragment were then transformed into competent *E. coli* DH5α cells. After an overnight incubation, single colonies were inoculated in liquid culture and incubated at 37°C overnight with shaking. The recombinant plasmids for the individual positive sample were analyzed and verified by the PCR method described above.

### Phylogenetic Analysis

The recombinant plasmid DNA for the individual positive sample was sequenced, and the sequence data were analyzed using DNASTar software. The sequences were subjected to phylogenetic analysis against the known *Marteilia* spp. sequences deposited in GenBank.

### Mixed Positive of *Marteilia* spp. with other Protozoans

To investigate if *Marteilia* spp.-positive shellfish were also mixed positive with *Haplosporidium nelsoni*, *Perkinsus* spp., and *Bonamia ostreae*, 156 *Marteilia* spp.-positive shellfish samples were also tested for these three parasites by PCR as previously described (Casas, Villalba, & Reece, 2002; Carnegie, Barber, Culloty, Figueras, & Distel, 2000; Day, Franklin, & Brown, 2000). The primers used in the PCR amplifications are listed in Table 2.

## Results

### Prevalence of *Marteilia* spp. in 13 Shellfish Species


A total of 11,581 shellfish was analyzed in this

**Table 1.** Prevalence of *Marteilia* spp. in 13 shellfish species in China

Shellfish species	No. of shellfish tested	No. of positive sample	Detection rate	No. of mixed positive	Mixed positive rate*
Oyster ( <i>C. rivulari</i> )	3,397	14	0.42%	5	35.71%
Oyster ( <i>C. gigas</i> )	1,351	2	0.15%	0	0
Oyster ( <i>C. angulata</i> )	861	17	1.97%	5	29.41%
Oyster (Total)	5,609	33	0.59%	10	30.30%
<i>M. meretrix</i>	1,026	2	0.19%	0	0
<i>C. nobilis</i>	1,012	9	0.89%	0	0
<i>R. philippinarum</i>	1,005	67	6.67%	23	34.33%
<i>P. viridis</i>	513	3	0.58%	0	0
<i>T. granosa</i>	506	11	2.17%	2	18.18%
<i>S. constricta</i>	603	31	5.14%	6	19.35%
<i>Haliotis discus hannai</i>	521	0	0	0	0
<i>P. undulata</i>	569	0	0	0	0
<i>T. dorsatus</i>	107	0	0	0	0
<i>C. antiquate</i>	110	0	0	0	0
Total	11,581	156	1.35%	41	26.28%

**Table 2.** Primers used for PCR detection of pathogens: *Haplosporidium nelsoni*, *Perkinsus* spp. and *Bonamia ostreae*

Parasites	Primers	Sequences (5'—3')	Length of PCR products (bp)	Reference
<i>Haplosporidium nelsoni</i>	MSX-A	CGACTTTGGCATT AGGTTT CAGACC	573	11
	MSX-B	ATGTGTTGGT GACGCT AACCG		
<i>Perkinsus</i> spp.	PerkITS-85	CCGCTTTGTTT GGATCCC	703	10
	PerkITS-750	ACATCAGGCCTTCTAATGATG		
<i>Bonamia ostreae</i>	CF	CGGGGGCATAATT CAGGAAC	760	7
	CR	CCATCTGCTGGAGACA CAG		

**Figure 1.** Sites for shellfish collection in coastal bay areas in China.

study, of which 156 (1.35%) were *Marteilia* spp. positive by PCR detection. The detection rate in the 13 shellfish species tested ranged from 0% to 6.67% (Table 1 and Figure 2). *R. philippinarum* (6.67%) and *S. constricta* (5.14%) showed the highest detection rate, whereas no *Marteilia* spp. positive was detected in *Haliotis discus hannai*, *P. undulate*, *T. dorsatus* or *C. antiquate* samples. The presence of *Marteilia* spp. in the other five shellfish species was lower than 2.17%.

### Prevalence of *Marteilia* spp. in Three Different Oyster Species

A proportion of *C. rivularis*, *C. gigas* and *C. angulata* was *Marteilia* spp. positive. The detection rate of these three oyster species ranged from 0.15% to 1.97% (Table 1). *C. angulata* samples showed the highest detection rate (1.97%), whereas *C. gigas* was the least positive oyster species.

**Prevalence of *Marteilia* spp. in the Oysters from Different Locations and Season**


The detection rate of oysters from different locations ranged from 0% to 2.30% (Table 3). *Marteilia* spp. positive samples were not present in the seawaters of Dalian, Wenzhou, Zhanjiang and Haikou bay areas in China. The oysters (*C. angulata*) in Xiamen bay (2.30%) had the highest detection rate of *Marteilia* spp. among the seven tested bay areas. There is no significant difference in the detection rate of *Marteilia* spp. in different seasons (data not shown).

**Phylogenetic Analysis**

The *Marteilia* spp. sequence of the individual positive sample from *C. rivulari*, *C. nobilis*, *R. philippinarum*, *P. viridis*, *T. granosa* and *S. constricta* had a high sequence homology (97.4% to 99.6%) with the sequences of other *Marteilia* type “M” from GenBank (Figure 3).

**Mixed Positive of *Marteilia* spp. with other Protozoan Parasites**


Among all 11,581 shellfish samples tested, 156



**Figure 2.** Prevalence of *Marteilia* spp. in 13 shellfish species.

**Table 3.** Prevalence of *Marteilia* spp. in the oysters form different locations

Location (species)	No. of oysters tested	No. of positive samples	Detection rate
Dalian ( <i>C. gigas</i> )	694	0	0
Qingdao ( <i>C. gigas</i> )	657	2	0.30%
Wenzhou (122 <i>C. angulata</i> + 475 <i>C. rivularis</i> )	597	0	0
Xiamen ( <i>C. angulata</i> )	739	17	2.30%
Zhanjiang ( <i>C. rivularis</i> )	540	0	0
Haikou ( <i>C. rivularis</i> )	454	0	0
Qinzhou ( <i>C. rivularis</i> )	1,928	14	0.73%
Total	5,609	33	0.59%



**Figure 3.** Phylogenetic sequences analysis of *Marteilia* spp. isolates.

shellfish were *Marteilia* spp. positive, and 41 samples (10 Oysters, 23 *R. philippinarum*s, 6 *S. constrictas* and 2 *T. granosa*) were mixed positive, resulting in a mixed positive rate of 26.28% (41/156, Table 1). As shown in Table 4, 30 samples were mixed positive for *Marteilia* spp. with *Perkinsus* spp.(mixed positive rate: 19.23%, 30/156). Three samples were mixed positive for *Marteilia* spp. with *Haplosporidium nelsoni* (mixed positive rate: 1.92%, 3/156). Eight samples were mixed positive for *Marteilia* spp. with *Perkinsus* spp. and *Haplosporidium nelsoni* (mixed positive rate: 5.13%, 8/156). None of the samples were mixed positive for *Marteilia* spp. with *Bonamia ostreae*. Mixed positive rate for *Marteilia* spp. with *Perkinsus* spp. (19.23%) was the highest among all tested samples.

## Discussion

*R. philippinarum* (6.67%) and *S. constricta* (5.14%) are the highly positive shellfish species among *Marteilia* spp. positive species (*C. rivularis*, *C. gigas* and *C. angulata*, *M. meretrix*, *C. nobilis*, *R. philippinarum*, *P. viridis*, *T. granosa* and *S. constricta*) in the tested seven bay areas in China. To our knowledge, this study is the first report on the prevalence of *Marteilia* spp. in oyster (*C. rivularis*, *C. gigas* and *C. angulata*), *M. meretrix*, *C. nobilis*, *R. philippinarum*, *T. granosa* and *S. constricta* in China. Previous studies have reported the detection rate of *Marteilia* spp. in Mussel in China to be 2.8% (5/180) (Wang, Lu, & Liang, 2012). Little is known about mortalities or other pathological effects of *Marteilia* spp. infection on oysters, *M. meretrix*, *C. nobilis*, *R. philippinarum*, *P. viridis*, *T. granosa* and *S. constricta* in the seven bay areas in China, as have been documented for infection of *Ostrea edulis* and *Mytilus edulis* and *Mytilus galloprovincialis* in other parts of the world (Itoh, Momoyama, & Ogawa, 2005; Lopez-Flores *et al.*, 2004; Fuentes *et al.*, 2002; Villalba, 1993). Therefore, these need to be studied in the future.

*Marteilia* spp. prevalence is highly variable – up to 98% in *O. edulis*, and higher prevalence is expected depending on farming practices and in the areas that have had more than one year of exposure to infection

(Feist, *et al.*, 2009; Berthe, *et al.*, 2004). Stocking at low density has been shown to be effective on the control and prevention of *Marteilia* spp. infection (Feist *et al.*, 2009). Copepod *Paracartia grani* (Copepoda, Calanoida) could contribute to the transmission of *Marteilia refringens* (Arzul *et al.*, 2013; Carrasco *et al.*, 2008; Audemard *et al.*, 2002). *Mecocyclops*, *Euchaeta concinna*, *Calanus sinicus* (Copepoda, Calanoida), *Paracalanus parvus* (Copepoda, Calanoida), *Calocalanus pavo* (Copepoda, Calanoida), *Phyllodiaptomus tunguidus* (Copepoda, Calanoida), etc., have been reported in China. However, to our knowledge, no study has been reported on the *Paracartia grani* in China. The general husbandry practices are usually stocking at high density in China, but *Marteilia* spp. has the low prevalence in Chinese coastal bays, which may due to lack of intermediate host *Paracartia grani* (or *Paracartia grani* is not the dominant species) in these seven bay areas of China, so it needs to be further studied.

The detection rates of *Ostrea edulis*, *Mytilus edulis* and *Mytilus galloprovincialis* are up to 98%, 40% and 5% (Feist, *et al.*, 2009; Berthe, Le Roux *et al.*, 2004; Le Roux *et al.*, 2001), respectively. But the detection rates among three oyster species in China are very low, ranging from 0.15% (*C. gigas*) to 1.97% (*C. angulata*), which suggests that the oyster species *Ostrea edulis* and *Mytilus edulis* may be more suitable for *Marteilia* spp. than other oyster species. The detection rates of oyster also different among the seven tested locations, ranging from 0% (Dalian, Wenzhou, Zhanjiang and Haikou) to 2.30% (Xiamen). *Marteilia* spp. positive was detected in Qingdao (0.30%), Xiamen (2.30%) and Qinzhou (0.73%), and Xiamen (2.30%) had the highest prevalence in the seven bay areas tested. These results suggest that the aquatic conditions in Dalian, Wenzhou, Zhanjiang and Haikou may not be optimal for the survival of *Marteilia* spp., and the aquatic conditions in Xiamen might provide a suitable environment for *Marteilia* spp. to infect oysters. The biological and environmental factors affecting the prevalence of *Marteilia* spp. in oysters should be further investigated.

The *Marteilia* spp. has three types “M”, “O” and

**Table 4.** Mixed positive of *Marteilia* spp. with other protozoan parasites

Shellfish species	Number of case			
	<i>Marteilia</i> spp. + <i>Perkinsus</i> spp.	<i>Marteilia</i> spp.+ <i>Haplosporidium</i> <i>nelsoni</i>	<i>Marteilia</i> spp.+ <i>Haplosporidium nelsoni</i> + <i>Perkinsus</i> spp.	<i>Marteilia</i> spp.+ <i>Bonamia</i> <i>ostreae</i>
<i>C. rivularis</i>	4	1	0	0
<i>C. angulata</i>	5	0	0	0
<i>R. philippinarum</i>	14	1	8	0
<i>S. constrictas</i>	5	1	0	0
<i>T. granosa</i>	2	0	0	0
Total	30	3	8	0
Mixed positive rate	19.23%	1.92%	5.13%	0

“C” (Carrasco *et al.*, 2012). The *Marteilia* spp. sequence of the individual positive sample from *C. rivulari*, *C. nobilis*, *R. philippinarum*, *P. viridis*, *T. granosa* and *S. constricta* detected in this study, had a high sequence homology with the sequences of the *Marteilia* type “M”. Because the number of the shellfish samples we sequenced was limited, there might be that also *Marteilia* type “O” and “C” in the shellfish of China, indicating further investigation is needed.

In this study, it is the first time to reveal that mixed positive of *Marteilia* spp. with *Perkinsus* spp. or *Haplosporidium nelsoni* was a common phenomenon, and that mixed positive of *Marteilia* spp. with *Perkinsus* spp. was the most common event (19.23%, 30/156). To our knowledge, this is the first report about the mixed positive of *Marteilia* spp. with other protozoan. It is still unknown whether the existence of *Perkinsus* spp. would bring a suitable living condition for *Marteilia* spp.. And more researches are required to figure out whether these mixed positive observations affect the mortality or survival rate of shellfish.

The prevalence of *Marteilia* spp. in 13 species of shellfish in China is based on the PCR detection. In order to confirm the infection of *Marteilia* spp. inside the tissue of the shellfish in China, histological analyses or *in situ* hybridization will be performed in our further study.

## Acknowledgments

This work was supported by the Guangxi Science and Technology Bureau (0630001-3M) and by the The National Ten-Thousand Talents Program of China (W02060083). We would like to thank Liqiang Zhao (Dalian Ocean University, Dalian), Bin Lin (Yellow Sea Fisheries Research Institute, Chinese Academy of Fishery Science, Qingdao), Maocang Yan (Zhejiang Marine Culture Research Institute, Wenzhou), Lei Wang (Xiamen University, Xiamen), Weibing Lan (Guangdong Ocean University, Zhanjiang) and Li Fang (Hainan Diagnosis Center of Animal Diseases, Haikou) for their assistance with the samples collection.

## References

- Arzul, I., Chollet, B., Boyer, S., Bonnet, D., Gaillard, J., Baldi, ... Bouchoucha, M. (2013). Contribution to the understanding of the cycle of the protozoan parasite *Marteilia refringens*, *Parasitology*, *141*(2), 227-240. <http://dx.doi.org/10.1017/S0031182013001418>.
- Audemard, C., Le Roux, F., Barnaud, A., Collins, C., Sautour, B., Sauria, P.G., Berthe, F. (2002). Needle in a haystack: involvement of the copepod *Paracartia grani* in the life-cycle of the oyster pathogen *Marteilia refringens*, *Parasitology*, *124*(03), 315-323. <http://dx.doi.org/10.1017/S0031182001001111>.
- Audemard, C., Sajus, M.C., Barnaud, A., Sautour, B., Sauriau, P.G., & Berthe, F.J. (2004). Infection dynamics of *Marteilia refringens* in flat oyster *Ostrea edulis* and copepod *Paracartia grani* in a claire pond of Marennes-Oleron Bay, *Diseases of Aquatic Organisms*, *61*(1-2), 103-111. <http://dx.doi.org/10.3354/dao061103>.
- Berthe, F.C.J., Le Roux, F., Adlard, R.D., & Figueras, A. (2004). Marteiliosis in molluscs: A review, *Aquatic Living Resources*, *17*(4), 433-448. <http://dx.doi.org/10.1051/alr:2004051>
- Berthe, F.C.J., Le Roux, F., Peyretailade, E., Peyret, P., Rodriguez, D., & Gouy, M. (2000). Phylogenetic Analysis of the Small Subunit Ribosomal RNA of *Marteilia refringens* Validates the Existence of Phylum Paramyxia (Desportes and Perkins, 1990) 1, *Journal of Eukaryotic Microbiology*, *47*(3), 288-293. <http://dx.doi.org/10.1111/j.1550-7408.2000.tb00049.x>
- Carnegie, R.B., Barber, B.J., Culloty, S.C., Figueras, A.J., & Distel, D.L. (2000). Development of a PCR assay for detection of the oyster pathogen *Bonamia ostreae* and support for its inclusion in the Haplosporidia, *Diseases of Aquatic Organisms*, *42*(3), 199-206. <http://dx.doi.org/10.3354/dao042199>.
- Carrasco, N., Andree, K.B., Lacuesta, B., Roque, A., Rodgers, C., & Furones, M.D. (2012). Molecular characterization of the *Marteilia* parasite infecting the common edible cockle *Cerastoderma edule* in the Spanish Mediterranean coast: A new *Marteilia* species affecting bivalves in Europe?, *Aquaculture*, *324*(325), 20-26. <http://dx.doi.org/10.1016/j.aquaculture.2011.10.017>.
- Carrasco, N., Arzul, I., Chollet, B., Robert, M., Joly, J.P., Furones, M.D., & Berthe, F.C.J. (2008). Comparative experimental infection of the copepod *Paracartia grani* with *Marteilia refringens* and *Marteilia maurini*, *Journal of Fish Diseases*, *31*(7), 497-504. <http://dx.doi.org/10.1111/j.1365-2761.2008.00910.x>
- Casas, S.M., Villalba, A., & Reece, K.S. (2002). Study of perkinsosis in the carpet shell clam *Tapes decussatus* in Galicia (NW Spain). I. Identification of the aetiological agent and *in vitro* modulation of zoosporulation by temperature and salinity, *Diseases of Aquatic Organisms*, *50*, 51-65. <http://dx.doi.org/10.3354/dao050051>.
- Day, J.M., Franklin, D.E., & Brown, B.L. (2000). Use of Competitive PCR to Detect and Quantify *Haplosporidium nelsoni* Infection (MSX disease) in the Eastern Oyster (*Crassostrea virginica*), *Marine Biotechnology*, *2*(5), 456-465.
- Feist, S.W., Hine, P.M., Bateman, K.S., Stentiford, G.D., & Longshaw, M. (2009). *Paramarteilia canceri* sp. n. (Cercozoa) in the European edible crab (*Cancer pagurus*) with a proposal for the revision of the order Paramyxida Chatton, 1911, *Folia Parasitologica*, *56*(2), 73-85. <http://dx.doi.org/10.14411/fp.2009.012>
- Fuentes, J., Lopez, J.L., Mosquera, E., Vázquez, J., Villalba, A., & Álvarez, G. (2002). Growth, mortality, pathological conditions and protein expression of *Mytilus edulis* and *M. galloprovincialis* crosses cultured in the Ria de Arousa (NW of Spain), *Aquaculture*, *213*(1), 233-251. [http://dx.doi.org/10.1016/S0044-8486\(02\)00046-7](http://dx.doi.org/10.1016/S0044-8486(02)00046-7).
- Kleeman, S.N., Le, R.F., Berthe, F., & Adlard, R.D. (2002). Specificity of PCR and *in situ* hybridization assays designed for detection of *Marteilia sydneyi* and *M. refringens*, *Parasitology*, *125*(2), 131-141. <http://dx.doi.org/10.1017/S0031182002001919>.

- Itoh, N., Momoyama, K., & Ogawa, K. (2005). First report of three protozoan parasites (a haplosporidian, *Marteilia* sp. and *Marteilioides* sp.) from the Manila clam, *Venerupis* (= *Ruditapes*) *philippinarum* in Japan, *Journal of Invertebrate Pathology*, 88(3), 201-206. <http://dx.doi.org/10.1016/j.jip.2004.12.001>.
- Le Roux, F., Lorenzo, G., Peyret, P., Audemard, C., Figueras, A., Vivares, C., Berthe, F. (2001). Molecular evidence for the existence of two species of *Marteilia* in Europe, *Journal of Eukaryotic Microbiology*, 48(4), 449-454. <http://dx.doi.org/10.1111/j.15507408.2001.tb00178.x>.
- Lopez-Flores, I., de la Herrán, R., Garrido-ramos, M. A., Navas, J. I., Ruiz-rejón, C., & Ruiz-rejón, M. (2004). The molecular diagnosis of *Marteilia refringens* and differentiation between *Marteilia* strains infecting oysters and mussels based on the rDNA IGS sequence, *Parasitology*, 129(4), 411-419. <http://dx.doi.org/10.1017/S0031182004005827>.
- Lopez-Flores, I., Garrido-Ramos, M.A., de la Herrán, R., Ruiz-Rejón, C., Ruiz-Rejón, M., & Navas, J.I. (2008a). Identification of *Marteilia refringens* infecting the razor clam *Solen marginatus* by PCR and in situ hybridization, *Molecular & Cellular Probes*, 22(3), 151-155. <http://dx.doi.org/10.1016/j.mcp.2008.01.002>.
- Lopez-Flores, I., Robles, F., Valencia, J.M., Grau, A., Villalba, A., de la Herrán, R., Navas, J.I. (2008b). Detection of *Marteilia refringens* using nested PCR and in situ hybridisation in *Chamelea gallina* from the Balearic Islands (Spain), *Diseases of Aquatic Organisms*, 82(1), 79-87. <http://dx.doi.org/10.3354/dao01966>.
- Villalba, A., Mourelle, S.G., Carmen, L.M., Carballal, M.J., & Azevode, C. (1993). Marteiliasis affecting cultured mussels *Mytilus galloprovincialis* of Galicia (NW Spain). I. Etiology, phases of the infection, and temporal and spatial variability in prevalence, *Diseases of Aquatic Organisms*, 16, 61-72. <http://dx.doi.org/10.3354/dao016061>.
- Villalba, A., Mourelle, S.G., Carballal, M.J., & Carmen Lopez, M. (1993). Effects on infection by the protistan parasite *Marteilia refringens* on the reproduction of cultured mussels *Mytilus galloprovincialis* in Galicia (NW Spain), *Diseases of Aquatic Organisms*, 17(3), 205-213. <http://dx.doi.org/10.3354/dao017205>.
- Wang, Z., Lu, X., & Liang, Y. (2012). A *Marteilia*-like parasite in blue mussels *Mytilus edulis* in China, *Journal of Aquatic Animal Health*, 24(3), 161-164. <http://dx.doi.org/10.1080/08997659.2012.675926>.
- Xie, Z., Xie, L., Fan, Q., Pang, Y., Deng, X., Xie, Z.Q., Khan, M. I. (2013). A duplex quantitative real-time PCR assay for the detection of *Haplosporidium* and *Perkinsus* species in shellfish, *Parasitology Research*, 112(4), 1597-1606. <http://dx.doi.org/10.1007/s00436-013-3315-5>.