

DNA Barcoding Revealed a High Percentage of Mislabeling in Commercial Fish Products: The First Empirical Survey in South Texas

Rita B. Tommasi , Sanjana M. Lamia, Dysocheata Van, Isaiah Estrada, Zoen L. Kuchler, Dean Ramsey, Jyoti Tamang, Gen Kaneko^{1,*} , Hashimul Ehsan¹ 

¹College of Natural and Applied Science, University of Houston-Victoria, 3007, N. Ben Wilson, Victoria, TX 77901, USA

How to Cite

Tommasi, R.B., Lamia, S.M., Van, D., Estrada, I, Kuchler, Z.L., Ramsey, D., Tamang, J., Kaneko, G., Ehsan H. (2024). DNA Barcoding Revealed a High Percentage of Mislabeling in Commercial Fish Products: The First Empirical Survey in South Texas. *Turkish Journal of Fisheries and Aquatic Sciences*, 24(7), TRJFAS25069. <https://doi.org/10.4194/TRJFAS25069>

Article History

Received 24 January 2024

Accepted 08 May 2024

First Online 23 May 2024

Corresponding Author

E-mail: ehsanh@uhv.edu

Keywords

FDA

Cytochrome

Imported fish

Fraudulent

Abstract

Seafood mislabeling is a widespread problem that have produced a growing distrust of seafood industry. In this study, we examined the prevalence of mislabeling in fish samples from regional markets in the greater Houston area and close-by coastal communities. A total of 63 fish fillet samples were purchased, labeled, and stored at -20°C in individual packages until DNA extraction. DNA fragments of roughly 700 base pairs encoding cytochrome oxidase I (COI) were PCR-amplified from each DNA sample. With 99.6–100% nucleotide identity in the GenBank and BOLD databases, all samples were correctly identified at the species level. The scientific names identified by DNA barcoding were matched with legally acceptable market names using the US Food and Drug Administration (FDA) Seafood List. Out of the 63 samples examined, 13 samples (20.6%) were mislabeled. Moreover, 24 samples (38.1%) did not use the acceptable market names, indicating that the FDA policy is poorly implemented in seafood industry. The first DNA barcoding survey this area warranted the need of continuous monitoring and the dissemination of the regulation combined with taxonomic knowledge.

Introduction

Fish mislabeling is a typical form of economic deception for the consumer. Intentional mislabeling is outright fraudulent, in which cheap fish is sold with a name of expensive fish such as halibut, swordfish, and snapper (Rasmussen & Morrissey, 2009). The confusion between the common name, acceptable market name, and scientific name is another source of fish mislabeling. For example, “flounder” includes 49 and 68 different species in FishBase and the US Food and Drug Administration (FDA) Seafood List, respectively (Food and Drug Administration, 2018; Froese & Pauly, 2023). To avoid misleading consumers, the European Union requires declaration of the scientific name for

unprocessed products to ensure robust seafood labeling (Paolacci et al., 2021). However, the US remains lenient, originating about 40% of the world’s mislabeled seafood (Kroetz et al., 2020). The regular assessment of fish product authenticity is of a particular importance to ensure the quality and food safety in the US fish market.

DNA barcoding has been a useful tool in identifying species. Aquatic species can be identified by various methods such as isozyme markers, restriction fragment length polymorphism (RFLP), and nuclear magnetic resonance (NMR) spectroscopy (Cermakova et al., 2023; Kaneko et al., 2019), but DNA barcoding has advantage over these methods in terms of accuracy and efficiency. Mitochondrial DNA (mtDNA) is a major target in DNA barcoding because: (1) it generally lacks large noncoding

regions; (2) it is abundant and relatively easy to analyze; (3) it does not undergo genetic rearrangements such as recombination; (4) sequence ambiguities resulting from heterozygous genotypes can be avoided, and (5) regions with low intraspecies and high interspecies variation are already identified. Although several mtDNA regions such as 16S rRNA (Itoi et al., 2005; Kochzius et al., 2008) and cytochrome b (Ha et al., 2018; Itoi et al., 2020; Pepe et al., 2005) have been used for DNA barcoding, the 5' region of cytochrome c oxidase subunit I (COI) has become a dominant region for identifying fish mislabeling (Di Pinto et al., 2013; Panprommin et al., 2020).

With a coastal extension of 450 miles, Texas is home to a large portion of fish commerce. Near-shore wetlands provide efficient natural water filters through plants and soils, serving as important nurseries for fish, crab, and shellfish in The Gulf of Mexico. A total economic value of commercial fisheries landed in Texas exceeded \$168 million in 2022 (NOAA, 2022), and correct labeling is an important obligation to retain ecological and financial responsibility of the state. However, we did not find any DNA barcoding studies on seafood mislabeling specifically targeting local markets or restaurants in Texas — only two previous studies analyzed fish samples from this region to our knowledge. A previous DNA barcoding study included a sushi restaurant in Austin along with those in New York and San Francisco as cities where consumers likely demand high-quality food (Khaksar et al., 2015). Forty-three fish samples from Austin and Houston were analyzed in 2012 as a part of a nation-wide fish mislabeling survey (Warner et al., 2019). While the public inspection should be in effect, considering the scale and period of these previous studies, a new DNA barcoding survey, even preliminary, will bring new insights into how fish products are handled in this large economic hub of fish commerce.

In the present study, we tested fish samples from local markets in the greater Houston area and nearby coastal cities. Several samples were mislabeled, but we also found a high number of unaccepted market names. Based on the results, possible future strategies to reduce fish mislabeling were discussed.

Methods

DNA Barcoding

Fish fillet samples (n=63) were purchased from local markets in South Texas coastal cities (Table 1t, mainly from the Houston area). All samples were labeled and stored at -20°C in individual packages until DNA extraction. A 10 - 50 mg piece of fast muscle tissue was cut from each sample, being sure to avoid contamination. Negative controls without DNA template were employed. A part of the samples was then analyzed by a BioRad Fish Barcoding Kit (Bio-Rad Laboratories Inc., Hercules, CA) according to the

manufacturer's instructions. PCR condition was 2 min at 94°C, followed by 35 cycles of 30 sec at 94°C, 2 min at 55°C, 1 min at 72°C. The final extension step was 10 min at 72°C. The PCR products were purified by a NucleoSpin Gel and PCR Clean-up kit (Macherey-Nagel, Düren, Germany). COI gene fragments were amplified from some samples according to the USDA recommended protocol (https://www.fsis.usda.gov/sites/default/files/media_file/2020-11/CLG-FPCR.pdf) using two primers: FishCO1LBC_m13F (5'-CACGACGTTGTAACGACTCAACYAATCAYAAAGATATYGGCAC-3') and FishCO1HBC_m13R (5'-GATAACAATTTACACAGGACTTCYGGGTGRCCRAARAATCA-3'). Advantage-HF 2 PCR kit (Clontech Laboratories, Santa Clara, CA) was used with the USDA primers. PCR was conducted at 95°C for 3 min, followed by 34 cycles of 95°C for 30 s, 55°C for 30 s, and 72°C for 1 min. The final extension step was at 72°C for 5 min. DNA sequences were determined by Eurofins Genomics (Louisville, KY). The obtained sequences were analyzed by 4Peaks (Nucleobytes) and submitted to the Barcode of Life Data (BOLD) system (<http://www.boldsystems.org/>) to determine species. The BLASTN program at the NCBI BLAST (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) was used with default parameters when the determined COI sequence was shorter than 500 bp. US Food and Drug Administration Seafood List (Food and Drug Administration, 2018) and FishBase (Froese & Pauly, 2023) were used to retrieve a list of possible species from labels (Supplementary 0 S1).

Results and Discussion

A DNA barcoding survey on fish mislabeling can choose one of the two strategies: testing a wide variety of species (Lakra et al., 2016; Pardo et al., 2018; Smith et al., 2008) or focusing on a specific group of fish (Bektas et al., 2019; Di Pinto et al., 2013; Lowenstein et al., 2009). We employed the former strategy since the present study was the first DNA barcoding survey specific to this geographical area to our knowledge. However, we included relatively higher numbers of expensive, white-fleshed species such as cod, flounder, and monkfish, which tend to be mislabeled (Hu et al., 2018).

The COI gene sequences solely containing the coding region were successfully determined for the 63 samples using the conventional PCR sequencing approach (Table 1). The average length of COI region determined was 595.5 bp. All samples were identified at the species level with 99.62-100% nucleotide identity. Out of the 63 samples, 13 samples (20.6%) were determined to be mislabeled, which was generally comparable to the lower end of mislabeling frequencies reported in US and Mexico: 15-17% for grocery in a nation-wide survey (Warner et al., 2019), 16.3% in restaurants in Austin, New York, and San Francisco (Khaksar et al., 2015). Higher mislabeling rates have also been reported such as 30.8% in markets and restaurants in Mazatlan, Mexico City, and Cancun (Munguia-Vega et

Table 1. Fish samples analyzed in the present study

Label	Possible species	BOLD/BLAST top hit	BOLD/BLAST identity (%)	Mislabeled	Accession No.	Length
Pollock	See Table S1	<i>Gadus chalcogrammus</i>	99.83	No	MH119970	609
Pacific cod*	<i>Gadus macrocephalus</i>	<i>G. chalcogrammus</i>	100	Yes	MK283727	508
Pacific cod*	<i>Gadus macrocephalus</i>	<i>G. macrocephalus</i>	100	No	MK283728	585
Pacific cod*	<i>Gadus macrocephalus</i>	<i>G. macrocephalus</i>	100	No	MK283729	587
Pacific cod*	<i>Gadus macrocephalus</i>	<i>G. macrocephalus</i>	100	No	MK283730	608
Pacific cod*	<i>Gadus macrocephalus</i>	<i>G. macrocephalus</i>	100	No	MK283731	585
Alaskan cod*	Considered as Alaska cod <i>Gadus macrocephalus</i>	<i>G. macrocephalus</i>	100	No	MK283715	593
Alaskan cod*	Considered as Alaska cod <i>Gadus macrocephalus</i>	<i>G. macrocephalus</i>	100	No	MK283718	627
Atlantic cod*	<i>Gadus morhua</i>	<i>G. morhua</i>	99.83	No	OR461757	581
Pacific whiting	<i>Merluccius productus</i>	<i>M. productus</i>	100	No	MK283720	627
Pacific whiting	<i>Merluccius productus</i>	<i>M. productus/angustimanus</i>	100	No	MK283732	617
Pacific whiting	<i>Merluccius productus</i>	<i>M. productus/angustimanus</i>	100	No	MK283733	596
Pacific whiting	<i>Merluccius productus</i>	<i>M. productus/angustimanus</i>	100	No	MK283734	610
Pacific whiting	<i>Merluccius productus</i>	<i>M. productus/angustimanus</i>	100	No	MK283735	610
Pacific whiting	<i>Merluccius productus</i>	<i>M. productus/angustimanus</i>	100	No	MK283736	597
Atlantic croaker*	<i>Micropogonias undulatus</i>	<i>M. undulatus</i>	100	No	KX163997	670
Yellow croaker*	<i>Larimichthys polyactis</i>	<i>Barbonymus altus</i>	100	Yes	MH119965	651
Flounder	See Table S1	<i>Lepidopsetta polyxystra</i>	100	No	MK283711	589
Flounder	See Table S1	<i>Lepidopsetta polyxystra</i>	100	No	MK283716	618
Flounder	See Table S1	<i>Lepidopsetta polyxystra</i>	100	No	MK283717	610
Flounder	See Table S1	<i>Limanda aspera</i>	100	No	MK283721	587
Flounder	See Table S1	<i>Limanda aspera</i>	100	No	MK283722	598
Flounder	See Table S1	<i>Limanda aspera</i>	100	No	MK283723	610
Flounder	See Table S1	<i>Limanda aspera</i>	100	No	MK283724	588
Flounder	See Table S1	<i>Limanda aspera</i>	100	No	MK283725	632
Flounder	See Table S1	<i>Limanda aspera</i>	100	No	MK283726	593
Summer flounder*	<i>Paralichthys dentatus</i>	<i>P. dentatus</i>	100	No	KX164002	629
Atlantic halibut*	<i>Hippoglossus hippoglossus</i>	<i>H. stenolepis</i>	100	Yes	KX164003	653
Halibut	<i>Hippoglossus hippoglossus</i> or <i>H. stenolepis</i>	<i>H. stenolepis</i>	100	No	MK283719	649
Monkfish	See Table S1	<i>L. americanus</i>	100	No	MH760553	563
Monkfish	See Table S1	<i>L. americanus</i>	100	No	MH760554	579
Monkfish	See Table S1	<i>L. americanus</i>	100	No	MH760555	589
Monkfish	See Table S1	<i>L. americanus</i>	100	No	MH760556	496
Monkfish	See Table S1	<i>L. americanus</i>	100	No	MH760557	432
Yellowfin tuna*	<i>Thunnus albacares</i>	<i>Xiphias gladius</i>	100	Yes	MH119957	597
Yellowfin tuna*	<i>Thunnus albacares</i>	<i>T. albacares</i>	100	No	MH119959	334
Amberjack	See Table S1	<i>Seriola dumerili</i>	100	No	MH119972	669
Blue runner	<i>Caranx crysos</i>	<i>C. crysos</i>	100	No	MH119976	633
Mackerel	See Table S1	<i>Atule mate</i>	99.62	Yes	MH119973	588
King mackerel	<i>Scomberomorus cavalla</i>	<i>S. cavalla</i>	100	No	MH119964	663
Norway mackerel*	Considered as mackerel; see Table S1	<i>Scomber scombrus</i>	100	No	MH119971	669
Atlantic pomfret*	<i>Brama brama</i>	<i>Peprilus medius</i>	100	Yes	MH119960	669
Big grouper*	Considered as giant grouper <i>Epinephelus fucoglottus</i> x <i>E. lanceolatus</i>	<i>Hyporthodus niphobles</i>	100	Yes	MH119963	462
Black drum*	<i>Pogonias cromis</i>	<i>Pogonias cromis</i>	100	No	KX164000	605
Seabass	See Table S1	<i>Dissostichus eleginoides</i>	100	Yes	MH119956	480
Snapper	See Table S1	<i>Pristipomoides multidens</i>	100	No	MH119958	437
Round scad*	<i>Decapterus punctatus</i>	<i>D. macarellus</i>	99.85	Yes	MH119969	675
Round scad*	<i>Decapterus punctatus</i>	<i>D. macarellus</i>	100	Yes	MH119978	663
Starry butterfish*	Considered as butterfish; see Table S1	<i>Peprilus medius</i>	100	Yes	MH119975	666
Shark	See Table S1	<i>Carcharhinus brevipinna</i>	100	No	MH119961	675
Striped Pangasius (Swai)	<i>Pangasianodon hypophthalmus</i>	<i>Pristipomoides multidens</i>	100	Yes	MK283712	640
Swai	<i>Pangasinodon hypophthalmus</i>	<i>P. hypophthalmus</i>	100	No	MH119968	222
Farmed Swai	<i>Pangasinodon hypophthalmus</i>	<i>P. hypophthalmus</i>	100	No	MH119967	639
Natural Tilapia*	Considered as tilapia; see Table S1	<i>Oreochromis niloticus</i>	100	No	MK283714	560
Catfish	See Table S1	<i>Ictalurus punctatus/dugesii</i>	100	No	MK283713	624
Catla	<i>Gibelion catla (Labeo catla)</i>	<i>G. catla</i>	100	No	KX163998	629
Grass carp	<i>Ctenopharyngodon idella</i>	<i>C. idella</i>	100	No	MH119977	615
Rohu	See Table S1	<i>Labeo rohita</i>	100	No	KX164001	669
White perch	<i>Morone americana</i>	<i>M. americana</i>	100	No	MH119962	675
White perch	<i>Morone americana</i>	<i>M. americana</i>	100	No	MH119974	606
Ganges river sprat*	<i>Corica soborna</i>	<i>C. soborna</i>	100	No	KX164004	647
Gangetic ailia	<i>Ailia coila</i>	<i>A. coila</i>	100	No	KX163999	658
River barb*	Considered as barb; see Table S1	<i>Micropogonias undulatus</i>	100	Yes	MH119966	660

al., 2022) and 47% in sushi restaurants in Los Angeles (Willette et al., 2017). Our previous survey on the genetically modified (GM) plants have found no false labeling in organic products, which should not contain any GM plants (Tegeler et al., 2020). These results may indicate the overall honest attitude of food industries toward food labeling in this area. In parallel, the low mislabeling rate is attributed to the generous regulation in US labeling guideline (Food and Drug Administration, 2018). When FishBase was used to match the scientific name and common name, as commonly done in previous studies (Lamendin et al., 2015; Panprommin & Manosri, 2022), the mislabeling rate was 36.5% (23 out of 63, [Supplementary Tables S2 and S3](#)). It should be noted that we analyzed more than 80 samples in the present study, but PCR products were obtained only from the 63 samples. This may be attributed to the use of degenerate universal primers or PCR amplification issues.

As shown in Table 1, 2 samples were mislabeled in the 17 white-fleshed marine fish (pollock, cod, Pacific whiting, and croaker). Pacific cod is a common name exclusively used for *Gadus macrocephalus* (FishBase, [Supplementary Table S3](#)), but this species was substituted with Alaska pollock *G. chalcogrammus*. These species share a similar morphology, and this substitution has also been reported in other studies (Feldmann et al., 2021; Helgoe et al., 2020). Moreover, many fish products did not have acceptable market names. For example, only “cod” and “Alaska cod” are acceptable market names of *G. macrocephalus* and *G. morhua* (FDA Seafood List, [Supplementary Table S1](#)), and thus “Pacific cod” and “Alaskan cod” samples have legal problems. Pacific whiting samples were identified as a correct species *Merluccius productus*, although the COI sequence also showed 100% nucleotide identity with Panama hake *M. angustimanus*. While this represents a limitation of DNA barcoding based on the COI gene, these two species are genetically close to each other. Both Atlantic croaker and yellow croaker must be labeled just as “croaker” ([Supplementary Table S1](#)). Yellow croaker *Larimichthys polyactis* was substituted with red tailed tinfoil *Barbonymus altus*, which belongs to the family Cyprinidae (freshwater fish such as carp and minnow). This substitution is likely an intentional fraudulent mislabeling.

We analyzed 12 flatfish samples including flounder and halibut. The flounder samples were identified as Northern rock sole *Lepidopsetta polyxystra* and yellowfin sole *Limanda aspera* (Table 1). These species are acceptable as “flounder” in the FDA Seafood list, but not in the FishBase. The difference accounted for the discrepancy in the mislabeling rates, 20.6% based on the FDA Seafood list and 36.5% based on FishBase. Summer flounder and Atlantic halibut were unacceptable market names. Flatfish mislabeling has been commonly reported from various regions of the world, including Italy (Pappalardo and Ferrito, 2015) and Brazil (Staffen

et al., 2017), possibly due to their characteristic morphology.

Three species in the genus *Lophius* can be sold as monkfish in US ([Supplementary Table S1](#)). Our DNA barcoding showed that all monkfish samples were *L. americanus*, identifying no mislabeling. The angel shark *Squatina squatina* used to be sold as monkfish, exposing the species to the risk of extinction (Porcher & Darvell, 2022), but this was not the case in the present study.

Other marine fish products showed relatively high frequencies of mislabeling and unaccepted market names (Table 1, from yellowfin tuna to shark). Among them, the substitution of yellowfin tuna *Thunnus albacares* with swordfish *Xiphias gladius* and that of seabass with Patagonian toothfish *Dissostichus eleginoides* could be intentional. The substituted species are relatively expensive and differ from the substituting species in morphology and distribution. Others may be attributed to the confusion. In this study, we analyzed a relatively high number of white-fleshed species that were commonly mislabeled. However, since the mislabeling frequency was higher than expected in other marine fish species, further studies could focus on these species with increased number of samples.

Two mislabeling cases were found in inexpensive freshwater fish (Table 1, swai to river barb). Swai *Pangasianodon hypophthalmus* was substituted with a marine fish, Goldbanded jobfish *Pristipomoides multidens*. This fish is also called Goldband snapper and is generally considered as a palatable fish species. River barb turned out to be Atlantic croaker *Micropogonias undulatus*, which is also a high-value species. While more samples are needed to reach a conclusion, these mislabeling cases may be attributed to confusion and/or poor traceability.

In summary, the first fish DNA barcoding study in the greater Houston area demonstrated that 20.6% of fish fillet samples were mislabeled, and 38.1% of the samples used unacceptable market names. Seafood mislabeling not only exploits consumers financially but also presents a potential hazard to individuals with allergies. Several approaches that have been proposed in previous studies should facilitate the solution of this problem (Cawthorn et al., 2018; Hu et al., 2018). Namely, mandatory inclusion of scientific names is the first policy to be implemented, which would eliminate the ambiguity between the scientific name, common name, and acceptable market name. Inclusion of geographical origin and harvest methods also benefit consumers. A regular DNA barcoding survey by individuals well acquainted with local community will help consumers to make educated decisions and encourage the seafood industry to use more accurate labeling

Ethical Statement

Not applicable

Funding Information

This work was supported by the maintenance and operation fund from the Dean's Office, College of Natural and Applied Science, the University of Houston-Victoria. GK was supported by M.G. and Lillie A. Johnson Foundation, Victoria, Texas

Author Contribution

Sanjana Lamia, Dysocheata Van, Isaiah Estrada was responsible for writing and editing the manuscript and contributed to its intellectual conception. Rita Tommasi, Zoen Kuchler, Dean Ramsey and Jyoti Tamang did the investigation, data curation, and analysis. Gen Kaneko and Hashimul Ehsan guided the paper's tone and concept, identified areas in need of revision and approved the final form.

Conflict of Interest

The authors declare that they have no competing interests.

Acknowledgements

The authors are grateful to the M.G. and Lillie A. Johnson Foundation, Victoria, Texas, for supporting students involved in this research.

References

- Bektas, Y., Aksu, I., Kaya, C., & Turan, D. (2019). DNA barcoding of the genus *Capoeta* (Actinopterygii: Cyprinidae) from Anatolia. *Turkish Journal of Fisheries and Aquatic Sciences*, 19(9), 739-752. https://doi.org/10.4194/1303-2712-v19_9_03
- Cawthorn, D. M., Baillie, C., & Mariani, S. (2018). Generic names and mislabeling conceal high species diversity in global fisheries markets. *Conservation Letters*, 11(5), e12573. <https://doi.org/10.1111/conl.12573>
- Cermakova, E., Lencova, S., Mukherjee, S., Horka, P., Vobruba, S., Demnerova, K., & Zdenkova, K. (2023) Identification of fish species and targeted genetic modifications based on DNA analysis: state of the art. *Foods*, 12(1), 228. <https://doi.org/10.3390/foods12010228>
- Di Pinto, A., Di Pinto, P., Terio, V., Bozzo, G., Bonerba, E., Ceci, E., & Tantillo, G. (2013). DNA barcoding for detecting market substitution in salted cod fillets and battered cod chunks. *Food Chemistry*, 141(3), 1757-1762. <https://doi.org/10.1016/j.foodchem.2013.05.093>
- Feldmann, F., Ardura, A., Blanco-Fernandez, C., & Garcia-Vazquez, E. (2021). DNA Analysis detects different mislabeling trend by country in European cod fillets. *Foods*, 10(7), 1515. <https://doi.org/10.3390/foods10071515>
- Food and Drug Administration. (2018). *The Seafood List*. <https://www.accessdata.fda.gov/scripts/fdcc/index.cfm?set=seafoodlist>
- Froese, R., & Pauly, D. (2023). *FishBase*. <https://www.fishbase.org> Accessed August 24th, 2023.
- Ha, T. T. T., Huong, N. T., Hung, N. P., & Guiguen, Y. (2018). Species identification using DNA barcoding on processed panga catfish products in Viet Nam revealed important mislabeling. *Turkish Journal of Fisheries and Aquatic Sciences*, 18(3), 457-462. https://doi.org/10.4194/1303-2712-v18_3_11
- Helgoe, J., Oswald, K. J., & Quattro, J. M. (2020). A comprehensive analysis of the mislabeling of Atlantic cod (*Gadus morhua*) products in Spain. *Fisheries Research*, 222, 105400. <https://doi.org/10.1016/j.fishres.2019.105400>
- Hu, Y., Huang, S. Y., Hanner, R., Levin, J., & Lu, X. (2018). Study of fish products in Metro Vancouver using DNA barcoding methods reveals fraudulent labeling. *Food Control*, 94, 38-47. <https://doi.org/10.1016/j.foodcont.2018.06.023>
- Itoi, S., Nakaya, M., Kaneko, G., Kondo, H., Sezaki, K., & Watabe, S. (2005). Rapid identification of eels *Anguilla japonica* and *Anguilla anguilla* by polymerase chain reaction with single nucleotide polymorphism-based specific probes. *Fisheries Science*, 71(6), 1356-1364. <https://doi.org/10.1111/j.1444-2906.2005.01102.x>
- Itoi, S., Tanaka, M., Mochizuki, Y., Oyama, H., Togawa, S., Yamada, R., Ito, T., Shishido, H., Masuda, Y., & Abe, K. (2020). Tandem PCR-RFLP analysis helps distinguish among three Japanese gnomefish (Teleostei: Scombroidea: Scombridae). *Ichthyological Research*, 67, 197-202. <https://doi.org/10.1007/s10228-019-00705-w>
- Kaneko, G., Ushio, H., & Ji, H. (2021) Application of magnetic resonance technologies in aquatic biology and seafood science. *Fisheries Science*, 85, 1-17. <https://doi.org/10.1007/s12562-018-1266-6>
- Khaksar, R., Carlson, T., Schaffner, D. W., Ghorashi, M., Best, D., Jandhyala, S., Traverso, J., & Amini, S. (2015). Unmasking seafood mislabeling in US markets: DNA barcoding as a unique technology for food authentication and quality control. *Food Control*, 56, 71-76. <https://doi.org/10.1016/j.foodcont.2015.03.007>
- Kochzius, M., Nölte, M., Weber, H., Silkenbeumer, N., Hjörleifsdóttir, S., Hreggvidsson, G., Marteinson, V., Kappel, K., Planes, S., & Tinti, F. (2008). DNA microarrays for identifying fishes. *Marine Biotechnology*, 10, 207-217. <https://doi.org/10.1007/s10126-007-9068-3>
- Kroetz, K., Luque, G. M., Gephart, J. A., Jardine, S. L., Lee, P., Chicojay Moore, K., Cole, C., Steinkruger, A., & Donlan, C. J. (2020). Consequences of seafood mislabeling for marine populations and fisheries management. *Proceedings of the National Academy of Sciences*, 117(48), 30318-30323. <https://doi.org/10.1073/pnas.2003741117>
- Lakra, W. S., Singh, M., Goswami, M., Gopalakrishnan, A., Lal, K., Mohindra, V., Sarkar, U., Punia, P., Singh, K., & Bhatt, J. (2016). DNA barcoding Indian freshwater fishes. *Mitochondrial DNA Part A*, 27(6), 4510-4517. <https://doi.org/10.3109/19401736.2015.1101540>
- Lamendin, R., Miller, K., & Ward, R. D. (2015). Labelling accuracy in Tasmanian seafood: an investigation using DNA barcoding. *Food Control*, 47, 436-443. <https://doi.org/10.1016/j.foodcont.2014.07.039>
- Lowenstein, J. H., Amato, G., & Kolokotronis, S.-O. (2009). The real maccyiii: identifying tuna sushi with DNA barcodes—contrasting characteristic attributes and genetic distances. *PLoS One*, 4(11), e7866. <https://doi.org/10.1371/journal.pone.0007866>
- Munguia-Vega, A., Terrazas-Tapia, R., Dominguez-Contreras, J. F., Reyna-Fabian, M., & Zapata-Morales, P. (2022). DNA barcoding reveals global and local influences on patterns

- of mislabeling and substitution in the trade of fish in Mexico. *PLoS One*, 17(4), e0265960
<https://doi.org/10.1371/journal.pone.0265960>
- NOAA. (2022). *NOAA Commercial Fisheries Landings*. Retrieved Aug 8 from [fisheries.noaa.gov/national/sustainable-fisheries/commercial-fisheries-landings](https://www.fisheries.noaa.gov/national/sustainable-fisheries/commercial-fisheries-landings) (Accessed Aug 8, 2023)
- Panprommin, D., & Manosri, R. (2022). DNA barcoding as an approach for species traceability and labeling accuracy of fish fillet products in Thailand. *Food Control*, 136, 108895. <https://doi.org/10.1016/j.foodcont.2022.108895>
- Panprommin, D., Soontornprasit, K., Tuncharoen, S., & Iamchuen, N. (2020). The utility of DNA barcoding for the species identification of larval fish in the lower ing river, Thailand. *Turkish Journal of Fisheries and Aquatic Sciences*, 20(9), 671-679 http://doi.org/10.4194/1303-2712-v20_9_02
- Paolacci, S., Mendes, R., Klapper, R., Velasco, A., Ramilo-Fernandez, G., Munoz-Colmenero, M., Potts, T., Martins, S., Avignon, S., & Maguire, J. (2021). Labels on seafood products in different European countries and their compliance to EU legislation. *Marine Policy*, 134, 104810 <https://doi.org/10.1016/j.marpol.2021.104810>
- Pappalardo, A.M. & Ferrito, V. (2015) DNA barcoding species identification unveils mislabeling of processed flatfish products in southern Italy markets. *Fisheries Research*, 164, 153-158.
<https://doi.org/10.1016/j.fishres.2014.11.004>
- Pardo, M. Á., Jiménez, E., Viðarsson, J. R., Ólafsson, K., Ólafsdóttir, G., Daniélsdóttir, A. K., & Pérez-Villareal, B. (2018). DNA barcoding revealing mislabeling of seafood in European mass caterings. *Food Control*, 92, 7-16 <https://doi.org/10.1016/j.foodcont.2018.04.044>
- Pepe, T., Trotta, M., Di Marco, I., Cennamo, P., Anastasio, A., & Cortesi, M. L. (2005). Mitochondrial cytochrome b DNA sequence variations: an approach to fish species identification in processed fish products. *Journal of Food Protection*, 68(2), 421-425.
<https://doi.org/10.4315/0362-028x-68.2.421>
- Porcher, I. F., & Darvell, B. W. (2022). Shark fishing vs. conservation: Analysis and synthesis. *Sustainability*, 14(15), 9548. <https://doi.org/10.3390/su14159548>
- Rasmussen, R. S., & Morrissey, M. T. (2009). Application of DNA-based methods to identify fish and seafood substitution on the commercial market. *Comprehensive Reviews in Food Science and Food Safety*, 8(2), 118-154. <https://doi.org/10.1111/j.1541-4337.2009.00073.x>
- Smith, P., McVeagh, S., & Steinke, D. (2008). DNA barcoding for the identification of smoked fish products. *Journal of Fish Biology*, 72(2), 464-471.
<https://doi.org/10.1111/j.1095-8649.2007.01745.x>
- Staffen, C.F., Staffen, M.D., Becker, M.L., Löfgren, S.E., Muniz, Y.C.N., de Freitas, R.H.A., & Marrero, A.R. (2017) DNA barcoding reveals the mislabeling of fish in a popular tourist destination in Brazil. *PeerJ* 5:e4006
<https://doi.org/10.7717/peerj.4006>
- Tegeler, Z., Kaneko, G., & Ehsan, H. (2020). Polymerase chain reaction detection of genetically modified organisms: a preliminary survey that predicts honest labeling under the mandatory labeling law. *Texas Public Health Journal*, 72, 8-12.
- Warner, K. A., Lowell, B., Timme, W., Shaftel, E., & Hanner, R. H. (2019). Seafood sleuthing: how citizen science contributed to the largest market study of seafood mislabeling in the US and informed policy. *Marine Policy*, 99, 304-311.
- Willette, D. A., Simmonds, S. E., Cheng, S. H., Esteves, S., Kane, T. L., Nuetzel, H., Pilaud, N., Rachmawati, R., & Barber, P. H. (2017). Using DNA barcoding to track seafood mislabeling in Los Angeles restaurants. *Conservation Biology*, 31(5), 1076-1085.