



Purification of Carbonic Anhydrase from *Capoeta umbla* (Heckel, 1843) Gills and Toxicological Effects of Some Metals on Enzyme Activity

Muammer Kırıcı^{1*}, Mahinur Kırıcı², Şükrü Beydemir³, Muhammed Atamanalp⁴

¹ Bingöl University, Faculty of Agriculture, Department of Fisheries, 12000-Bingöl, Turkey.

² Bingöl University, Faculty of Arts and Science, Department of Chemistry, 12000-Bingöl, Turkey.

³ Bursa Technical University, Department of Chemistry, Faculty of Natural Sciences, Architecture and Engineering, 16190-Bursa, Turkey.

⁴ Atatürk University, Faculty of Fisheries, 25240-Erzurum, Turkey.

* Corresponding Author: Tel.: +90.536 8917150; Fax: +90.426 2160029;
E-mail: muammerkirci@hotmail.com

Received 14 April 2015
Accepted 18 March 2016

Abstract

In this study, *in vitro* effects of some metal ions (Fe^{3+} , Cd^{2+} , Pb^{2+} and Ni^{2+}) on cytoplasmic carbonic anhydrase (CA, EC 4.2.1.1) from *Capoeta umbla* gill was investigated. CA was purified from the gills of *C. umbla* for the first time. It was purified with the Sepharose-4B-L-Tyrosine Sulphanilamide affinity chromatography method. The overall purification was approx. 31.69-fold with a yield of 53.33%, and a specific activity of 326.73 EU/mg proteins. Sodium dodecyl sulphate–polyacrylamide gel electrophoresis (SDS-PAGE) showed a single band corresponding to a molecular weight of approx. 29 kDa. The constants of the enzyme inhibitor complex (K_i) and 50% inhibitory values (IC_{50}) for metal ions were determined by Lineweaver–Burk graphs and plotting activity % vs. $[I]$, respectively. The K_i constants and IC_{50} values were 0.012 ± 0.0135 and 0.136 mM for Fe^{3+} , 0.019 ± 0.0113 and 0.191 mM for Cd^{2+} , 0.041 ± 0.0075 and 0.289 mM for Pb^{2+} , and 0.120 ± 0.034 and 0.924 mM for Ni^{2+} . It was determined that Fe^{3+} , Cd^{2+} and Pb^{2+} inhibited the enzyme competitively while Ni^{2+} inhibited the enzyme noncompetitively. The potential inhibitor for *C. umbla* gill CA was found as Fe^{3+} from these results.

Keywords: *Capoeta umbla*, carbonic anhydrase, gills, metal toxicity.

Capoeta umbla (Heckel, 1843) Solungaç Dokusundan Karbonik Anhidraz Enziminin Saflaştırılması ve Bazı Metallerin Enzim Aktivitesi Üzerine Toksikolojik Etkilerinin İncelenmesi

Özet

Bu çalışmada, *Capoeta umbla* solungaç sitoplazmik karbonik anhidraz enzimi üzerine bazı metallerin (Fe^{3+} , Cd^{2+} , Pb^{2+} ve Ni^{2+}) *in vitro* etkileri incelendi. Karbonik anhidraz enzimi *C. umbla* solungaç dokusundan ilk kez saflaştırıldı. Enzim, Sepharose 4B–L–tirozin–sulfanilamid afinite kromatografisi yöntemi ile saflaştırıldı. Bu yöntem sonucunda enzim, spesifik aktivitesi 326,73 EÜ/mg protein ve %53,3 verimle yaklaşık 31,69 kat olarak saflaştırıldı. Sodyum dodesil sülfat–poliakrilamid jel elektroforezinde (SDS-PAGE) yaklaşık 29 kDa’da tek bant olarak görüldü. Metal iyonları için enzim inhibitör kompleks sabitleri (K_i) ve %50 inhibitör değerleri (IC_{50}), sırasıyla Lineweaver–Burk grafikleri ve % aktivite– $[I]$ grafikleri çizilerek hesaplandı. K_i sabitleri ve IC_{50} değerleri, Fe^{3+} için $0,012 \pm 0,0135$ ve $0,136$ mM, Cd^{2+} için $0,019 \pm 0,0113$ ve $0,191$ mM, Pb^{2+} için $0,041 \pm 0,0075$ ve $0,289$ mM ve Ni^{2+} için $0,120 \pm 0,034$ ve $0,924$ mM olarak belirlendi. Fe^{3+} , Cd^{2+} ve Pb^{2+} iyonları enzimi, yarışmalı olarak inhibe ederken, Ni^{2+} iyonu yarışmasız olarak inhibe etmiştir. Sonuç olarak, Fe^{3+} iyonunun *C. umbla* solungaç karbonik anhidraz enzimi için güçlü bir inhibitör olduğu belirlendi.

Anahtar Kelimeler: *Capoeta umbla*, karbonik anhidraz, solungaç, metal toksisitesi.

Introduction

Metals are natural trace components of the aquatic environment, but their levels have increased due to industrial, agricultural and mining activities (Kalay and Canlı, 2000). This situation may be hazardous for living systems, especially aquatic organisms, including specific enzymes. It is well-known that enzymes catalyse almost all chemical reactions in the metabolism of the living systems.

These chemical substances including pollutants, pesticides, drugs and metal ions influence metabolism at low concentrations by decreasing or increasing enzyme activities (Ekinci *et al.*, 2007). Specifically, some enzymes including carbonic anhydrase (CA) are considered drug- and chemical-targeted enzymes. For example, CA is expressed in almost all the tissues of living things. Because of these, the enzyme, which has great importance in the balance of pH and respiration in various tissues, is a known target

enzyme for different substances (Coban *et al.*, 2008; Söyüt and Beydemir, 2011).

CA is a zinc metalloenzyme catalysing the reversible hydration of CO₂ to produce H⁺ and HCO₃⁻ (Supuran, 2010). CA plays key roles in a wide variety of physiological processes involving CO₂ and HCO₃⁻. In animals the various CA isozymes are found in many different tissues and are involved in a number of different physiological processes, including bone resorption, calcification, ion transport, acid–base transport, and a number of different metabolic processes such as biosynthetic reactions (gluconeogenesis, lipogenesis, and ureagenesis). In algae and plants they play an important role in photosynthesis (Ivanov *et al.*, 2007; Cannon *et al.*, 2010; Zhang *et al.*, 2010; Lionetto *et al.*, 2012).

The vertebrate gas exchange organ, the gills is essentially composed of a highly complex vasculature, surrounded by a high surface area epithelium that provides a thin barrier between a fish's blood and aquatic environment. Fish gills are the first point of contact between metals and impurities in water. Metals can bind to gill of fish and corrupt the ion-regulatory and respiratory functions of the gills (Playle, 1998). CA, abundantly present in gill epithelial cells, is assumed to play a role in these processes (Perry and Laurent, 1990; Evans *et al.*, 2005). The erythrocyte CA is the primary and possibly the only site of HCO₃⁻ dehydration in fish, forming CO₂ which diffuses across the respiratory gill epithelium into the ventilatory water. A small portion of the CO₂ is hydrated by cytoplasmic branchial CA, supplying HCO₃⁻ and H⁺ for Na⁺/H⁺ (NH₄⁺) and Cl⁻/HCO₃⁻ apical ionic exchangers. In view of these processes regulating acid–base and NaCl balance, the cell-specific localization of CA in fish gills has been investigated (Henry *et al.*, 1988; Sender *et al.*, 1999) and a large interspecies variability has been observed (Conley and Mallatt, 1988).

In the present study, CA, which is very important and ubiquitous in metabolism, was purified from *C. umbla* gill by a fast and simple technique. Additionally, *in vitro* inhibition effects of some metals (Fe³⁺, Cd²⁺, Pb²⁺ and Ni²⁺) *C. umbla* gill CA were investigated.

Materials and Methods

Chemicals

Sepharose-4B, protein assay reagents and chemicals for electrophoresis were purchased from Sigma–Aldrich. All other chemicals were of analytical grade and obtained from Merck.

Preparation of the Homogenate

Fish samples were caught from Murat River (Bingöl, Genç). Gill samples were taken from each

fish. Gills samples were washed three times with 0.9% NaCl, an isotonic saline solution. The gill cells were lysed by immersion in liquid nitrogen (approximately -163 °C). The lysed sample was transferred to a buffer solution (50mM Tris–HCl + 0.1 M Na₂SO₄, pH 8.0) and centrifuged at 4°C, 15000 g for 60 min (Le Trang *et al.*, 1983). The plasma and precipitate were removed. Supernatant was used in further studies.

Purification of CA from *C. umbla* Gills by Affinity Chromatography

CNBr activated Sepharose-4B was washed with ddH₂O. After that, tyrosine was attached to the activated gel as a spacer arm and finally diazotized sulphanilamide clamped with tyrosine molecule as ligand. The homogenate was applied to the prepared Sepharose-4B-L-Tyrosine Sulphanilamide affinity column equilibrated with 25 mM Tris–HCl/0.1M Na₂SO₄ (pH 8.7). The affinity gel was washed with 25 mM Tris–HCl/22mM Na₂SO₄ (pH 8.7). *C. umbla* gill CA was eluted with 1 M NaCl/25 mM Na₂HPO₄ (pH 6.3). All procedures were performed at 4°C (Bayram *et al.*, 2008). The enzyme used this *in vitro* experiment section was dialyzed.

Measurement of CA Activity

CA activity was assayed by following the hydration of CO₂ according to the method described by Wilbur and Anderson (Wilbur and Anderson, 1948). CO₂-hydratase activity as an enzyme unit (EU) was calculated by using the equation $(t_0 - t_c) / t_c$ where t_0 and t_c are the times for pH change of the non-enzymatic and enzymatic reactions, respectively.

Protein Determination

Protein levels were spectrophotometrically measured at 595 nm according to Bradford's method (Bradford, 1976), with bovine serum albumin being used as a standard.

SDS–Polyacrylamide Gel Electrophoresis (SDS–PAGE)

The control of enzyme purity was carried out using Laemmli's procedure (Laemmli, 1970) in 3% and 8% acrylamide concentrations for running and stacking gel, respectively. SDS (10%) was added to the gel solution. The gel was stabilized in a solution containing 50% propanol + 10% TCA + 40% distilled water for 30 min. Staining was performed for about 2 h in a solution of 0.1% Coomassie Brilliant Blue R-250 + 50% methanol + 10% acetic acid. Finally, washing was carried out in a solution of 50% methanol + 10% acetic acid + 40% distilled water until the protein bands were cleared.

In vitro Inhibition Assays

The effects of increasing concentrations of Fe^{3+} , Cd^{2+} , Pb^{2+} and Ni^{2+} on *C. umbla* gill CA activities were determined colorimetrically using CO_2 -hydratase assay. The metals were also tested in the hydratase activity assay in triplicate at each concentration used. Different concentrations of metals were examined in preliminary assays and as described in the literature (Vitale *et al.*, 1999). Enzyme activities were measured in the presence of different concentrations of Fe^{3+} (0.06, 0.10, 0.12, 0.13 and 0.26 mM), Cd^{2+} (0.07, 0.09, 0.14, 0.21 and 0.35 mM), Pb^{2+} (0.20, 0.29, 0.32, 0.36 and 0.43 mM) and Ni^{2+} (0.32, 0.65, 0.97, 1.13 and 1.29 mM). Control enzyme activity in the absence of a metal was taken as 100%. For each metal, an activity % vs. inhibitor concentration tube was drawn using conventional polynomial regression software (Microsoft Office 2010, Excel). Metal concentrations that produced 50% inhibition (IC_{50}) calculated from graphs (Söyüt and Beydemir, 2011).

To determine K_i constants, three different inhibitor concentrations were tested: 0.04, 0.12 and 0.26 mM for Fe^{3+} ; 0.07, 0.14 and 0.21 mM for Cd^{2+} ; 0.20, 0.32 and 0.43 mM for Pb^{2+} and 0.45, 0.65 and 1.27 mM for Ni^{2+} . *P*-nitrophenylacetate was used as a

substrate. The substrate concentrations were 0.15, 0.3, 0.45, 0.6 and 0.75 mM in these experiments. The Lineweaver-Burk curves obtained were used for the determination of K_i and inhibitor type (Lineweaver and Burk, 1934).

Results

The *C. umbla* gill CA was purified by Sepharose 4B-L-tyrosine-sulfanylamine affinity chromatography and purity was confirmed by SDS-PAGE (Figure 1). The enzyme was purified 31.69-fold with a yield of 53.33% with a specific activity of 326.73 EU/mg protein from *C. umbla* gill (Table 1).

In addition to purification of the enzyme, we investigated the *in vitro* effects of Fe^{3+} , Cd^{2+} , Pb^{2+} and Ni^{2+} on fish gill CA activity. All metal showed inhibitory effects on the enzyme. IC_{50} values of Fe^{3+} , Cd^{2+} , Pb^{2+} and Ni^{2+} were 0.136, 0.191, 0.289 and 0.924 mM, and K_i constants were 0.012 ± 0.0135 , 0.019 ± 0.0113 , 0.041 ± 0.0075 and 0.120 ± 0.034 mM, respectively (Table 2). While Fe^{3+} , Cd^{2+} and Pb^{2+} showed competitive inhibition, Ni^{2+} showed noncompetitive inhibition (Figure 2, Figure 3). According to these results, Fe^{3+} has higher inhibition effects than Cd^{2+} , Pb^{2+} and Ni^{2+} .






Figure 1. SDS-Polyacrylamide Gel Electrophoresis of *C. umbla* CA Purified by Sepharose 4B-L-Tyrosine-Sulphanilamide Affinity Gel Chromatography (1: Standard proteins. 2: *C. umbla* gill CA).

Table 1. Summary of purification procedure for *C. umbla* gill CA enzyme by a Sepharose-4B-tyrosine-sulfanylamine affinity column chromatography

Purification steps	Activity (EU/ml)	Total volume (ml)	Protein (mg/ml)	Total protein (mg)	Total activity (EU)	Specific activity (EU/mg)	Yield (%)	Purification Factor
Homogenate	300	60	29.10	1746	18000	10.309	100	1
Sepharose-4B-L tyrosine-sulphanilamide affinity chromatography	2141.67	6	4.897	29.382	9600	326.73	53.33	31.69

Table 2. The results of the activity of CA; K_i , IC_{50} values and inhibition types

Inhibitor	IC_{50} (mM)	K_i (mM)	Inhibition type
Fe^{3+}	0.136	0.012 ± 0.0135	Competitive
Cd^{2+}	0.191	0.019 ± 0.0113	Competitive
Pb^{2+}	0.289	0.041 ± 0.0075	Competitive
Ni^{2+}	0.924	0.120 ± 0.034	Noncompetitive

**Figure 2.** Activity-%-[Metal] regression analysis graphs for *C. umbla* gill CA in the presence of four different metal concentrations.

Discussions

Several ecological changes occur in water because of human activities, including agricultural,

industrial and municipal wastes. The pollution of the aquatic environment with heavy metals has become a serious health concern because of their toxicity and accumulation by organisms (Mendil *et al.*, 2010).

Some metals, such as Cd, Hg and Pb, have no known essential role in living organisms, and are toxic at even low concentrations. But at low levels, some metals such as Cu, Co, Zn, Fe and Mn are essential for enzymatic activity and several biological processes. The essential metals also become toxic at high concentrations (Bryan, 1976). For example, many enzymes contain Zn^{2+} within their structures, such as CA (Lionetto et al., 2012).

CAs, the zinc-binding enzyme family, has important roles in a number of tissues for living physiology, for example, pH and CO_2 homeostasis (Soyut et al., 2008). CO_2 , produced in fish tissues, is hydrated rapidly by CA enzyme, converted into HCO_3^- , and transported in the blood (Geers and Gros, 2000; Esbaugh et al., 2005; Esbaugh and Tufts, 2006; Supuran, 2008). Approximately 98% of the transported and stored CO_2 is in HCO_3^- form. At the respiratory epithelium, erythrocytic CA catalyses the rapid dehydration of HCO_3^- to molecular CO_2 , which then diffuses passively into the ventilatory water stream. Moreover, the CO_2/HCO_3^- system constitutes one of the most important physiological buffers for acid-base regulation (Doğan, 2006; Alp et al., 2010; Ekinçi and Beydemir, 2010).

Up to now, CA has been purified from many different tissues including human erythrocytes (Bayram et al., 2008; Şentürk et al., 2009), fish gills (Bone et al., 1995; Skaggs and Henry, 2002; Hisar et al., 2004; Ceyhun et al., 2010), fish erythrocytes (Bülbül et al., 2003; Hisar et al., 2005; Doğan, 2006), rainbow trout brain and liver (Söyüt and Beydemir, 2008; Söyüt et al., 2008). Although, CA and the inhibitory effects of many metals on CA have been studied in most fish tissues, no study has been done on *C. umbla* gill CA, yet.

For this reason, we investigated *in vitro* effects of some metals, Fe^{3+} , Cd^{2+} , Pb^{2+} and Ni^{2+} , on CA enzyme from *C. umbla* gill. It was purified with the Sepharose-4B-L-Tyrosine Sulphanilamide affinity chromatography method. The overall purification was approx. 31.69-fold with a yield of 53.33%, and a specific activity of 326.73 EU/mg proteins. The enzyme purity and subunit molecular weight (29 kDa) were determined by SDS-PAGE electrophoresis method (Figure 1). Additionally, the four metals (Fe^{3+} , Cd^{2+} , Pb^{2+} and Ni^{2+}) dose-dependently inhibited *in vitro* CA activity. IC_{50} values for Fe^{3+} , Cd^{2+} , Pb^{2+} and Ni^{2+} were 0.136, 0.191, 0.289 and 0.924 mM, and K_i constants were 0.012 ± 0.0135 , 0.019 ± 0.0113 , 0.041 ± 0.0075 and 0.120 ± 0.034 mM, respectively (Table 2). While Fe^{3+} , Cd^{2+} and Pb^{2+} showed competitive inhibition, Ni^{2+} showed noncompetitive inhibition (Figure 2, Figure 3). According to these results, Fe^{3+} has higher inhibition effects than Cd^{2+} , Pb^{2+} and Ni^{2+} .

In vitro effects of some heavy metals on enzymes, such as intestinal and branchial CA and $Na^+ - K^+ - ATPase$, which play a key role in salt and osmoregulation and acid-base balance in the teleost

fish, was studied. The early work of Christensen and Tucker (1976) demonstrated CA inhibition by heavy metals for the first time in fish. The study was carried out on red blood cells CA of the teleost *Oncorhynchus mykiss*. Erythrocyte CA, which represents the most abundant pool of the enzyme in fish, appeared significantly *in vitro* inhibited by several heavy metals, such as, Cd^{2+} , Cu^{2+} , Ag^+ , and Zn^{2+} (Lionetto et al., 2012). In another study, CA activities in gill and intestinal of European eel (*Anguilla Anguilla*) homogenates were significantly inhibited by Cd^{2+} . The gill CA was much more sensitive to heavy metal as compared to the enzyme activity in the intestine, as observed by comparing the IC_{50} values (gill IC_{50} : 0.01 mM and intestine IC_{50} : 0.036 mM) (Lionetto et al., 2000). The effects of ammonia and urea on branchial CA enzyme which plays a key role in ion regulation, osmoregulation and acid-base balance of rainbow trout (*Oncorhynchus mykiss*) were investigated. CA activities of the groups exposed to ammonia and urea were measured at 1, 2 and 3 h. The inhibitory effects of ammonia and urea on gill CA activities started immediately after the exposures. The differences between the initial CA activities for the controls were not significant ($P > 0.01$). The CA activities were significantly ($P < 0.01$) inhibited both in ammonia and urea group. However, the ammonia was inhibited more than urea since there were significant differences between the final values of gill CA activities (Hisar et al., 2004). Another study demonstrated *in vitro* and *in vivo* effects of the pesticides, deltamethrin, diazinon, propoxur and cypermethrin on the activity of rainbow trout gill CA. The four pesticides dose-dependently inhibited *in vitro* CA activity. IC_{50} values for deltamethrin, diazinon, propoxur and cypermethrin were 0.137, 0.267, 0.420 and 0.460 μM , respectively. *In vitro* results showed that pesticides inhibit fish gill CA activity in the rank order of deltamethrin > diazinon > propoxur > cypermethrin. Besides, *in vivo* studies of deltamethrin were performed on CA activity of rainbow trout gill. Fish gill CA was significantly inhibited at three concentrations (0.25, 1.0 and 2.5 $\mu g/L$) at 24 and 48 h (Ceyhun et al., 2010).

Söyüt and Beydemir (2011) investigated the *in vitro* effects of Co^{2+} , Zn^{2+} , Cu^{2+} , Cd^{2+} and Ag^+ on chemical-targeted CA enzyme from rainbow trout kidney. The enzyme was purified with a specific activity of 17.285 EU/mg and 31.7% yield and approximately 1800-fold using simple affinity purification method. Molecular weights of the subunit and native enzyme were estimated as 28.7 kDa and 26.9 kDa via SDS-PAGE and Sephadex-G 200 column, respectively. Inhibitory effects of Co^{2+} , Zn^{2+} , Cu^{2+} , Cd^{2+} and Ag^+ on CA activity were determined using the esterase method under *in vitro* conditions. IC_{50} and K_i values were calculated for metals. K_i values for Co^{2+} , Zn^{2+} , Cu^{2+} , Cd^{2+} and Ag^+ were 0.035, 1.2, 34.8, 103 and 257 mM from Lineweaver-Burk graphs, respectively. Consequently, *in vitro* inhibition rank order was determined as

$\text{Co}^{2+} > \text{Zn}^{2+} > \text{Cu}^{2+} > \text{Cd}^{2+} > \text{Ag}^+$. Alım et al. (2014) investigated the inhibition effects of some metal concentrations (Ag^+ , Cu^{2+} , Pb^{2+} , Zn^{2+} , Cd^{2+} , Co^{2+}) on the activity of CA from tuna (*Thunnus thynnus* Linnaeus, 1758) gill, *in vitro*. The tuna gill CA was obtained by affinity chromatography with a yield of 31%, a specific activity of 1062 EU/mg protein and a purification of 44.06-fold. The purity of the enzyme was confirmed by SDS-PAGE. Inhibitory effects of metals on CA activity were determined at different concentrations using the hydratase method under *in vitro* conditions. Consequently, *in vitro* inhibition rank order was determined as $\text{Ag}^+ > \text{Cu}^{2+} > \text{Pb}^{2+} > \text{Zn}^{2+} > \text{Cd}^{2+} > \text{Co}^{2+}$. From these results, they showed that Ag^+ is the most potent inhibitor of CA enzyme.

Consequently, we investigated the *in vitro* effects of the metals, Fe^{3+} , Cd^{2+} , Pb^{2+} and Ni^{2+} , on the activity of *C. umbla* gill CA. Carbonic anhydrase was purified from the gills of *C. umbla* for the first time. K_i and IC_{50} parameters of these metals were determined. It is clear that Fe^{3+} is the most potent inhibitor for CA enzyme. K_i and IC_{50} graphs show that Fe^{3+} , Cd^{2+} , Pb^{2+} inhibit CA in a competitive manner and Ni^{2+} inhibit CA in a noncompetitive manner (Figure 2, Figure 3, Table 2). According to K_i constants, the best inhibitor for *C. umbla* gill CA is Fe^{3+} . Our results are in agreement with others reported about teleost fish in the literature. *C. umbla*, living in rivers and lakes, is consumed highly by the local residents. The rising industrial and settlement areas near the rivers cause a great danger for the creatures living in watery areas. The required precautions should be taken immediately.

References

- Alım, Z., Çamur, B., Beydemir, Ş. and Küfrevioğlu, Ö.İ. 2014. The correlation between some metal concentrations and carbonic anhydrase activity in Tuna (*Thunnus thynnus* Linnaeus, 1758) gill. Hacettepe Journal of Biology and Chemistry, 42 (2): 219-224.
- Alp, C., Ekinci, D., Gültekin, M.S., Şentürk, M., Şahin, E. and Küfrevioğlu, Ö.İ. 2010. A novel one-pot synthesis of new 1-tosyl pyrrol-2-one derivatives and analysis for carbonic anhydrase inhibitory potencies. Bioorganic and Medicinal Chemistry, 18: 4468-4474. doi:10.1016/j.bmc.2010.04.072.
- Bayram, E., Şentürk, M., Küfrevioğlu, Ö.İ. and Supuran, C.T. 2008. *In vitro* inhibition of salicylic acid derivatives on human cytosolic carbonic anhydrase isozymes I and II. Bioorganic and Medicinal Chemistry, 16: 9101-9105. doi: 10.1016/j.bmc.2008.09.028.
- Bone, Q., Marshall, N.B. and Blaxter, J.H.S. 1995. In Sensory systems and communication. In: Q. Bone, N.B. Marshall and J.H.S. Blaxter (Ed.), Biology of Fishes Chapman, Hall, New York: 219-261.
- Bradford, M.M. 1976. A rapid and sensitive method for the quantization of microgram quantities of protein utilizing the principle of protein-dye binding. Analytical Biochemistry, 72: 248-251.
- Bryan, G.W. 1976. Some effects of heavy metal tolerance in aquatic organisms. In: A.P.M. Lockwood (Ed.) Effects of pollutants on aquatic organisms, Cambridge University Press. Cambridge, England. pp.7.
- Bülbul, M., Hisar, O., Beydemir, Ş., Çiftci, M. and Küfrevioğlu, Ö.İ. 2003. The *in vitro* and *in vivo* inhibitory effects of some sulfonamide derivatives on rainbow trout (*Oncorhynchus mykiss*) erythrocyte carbonic anhydrase activity. Journal of Enzyme Inhibition and Medicinal Chemistry, 18 (4): 371-375.
- Cannon, G.C., Heinhorst, S. and Kerfeld C.A. 2010. Carboxysomal carbonic anhydrases: Structure and role in microbial CO(2) fixation. Biochimica et Biophysica Acta-Proteins and Proteomics, 1804: 382-392. doi:10.1016/j.bbapap.2009.09.026.
- Ceyhun, S.B., Şentürk, M., Erdoğan, O. and Küfrevioğlu, Ö.İ. 2010. *In vitro* and *in vivo* effects of some pesticides on carbonic anhydrase enzyme from rainbow trout (*Oncorhynchus mykiss*) gills. Pesticide Biochemistry and Physiology, 97: 177-181. doi:10.1016/j.pestbp.2010.01.003.
- Ceyhun, S.B., Şentürk, M., Yerlikaya, E., Erdoğan, O., Küfrevioğlu, Ö.İ. and Ekinci, D., 2011. Purification and characterization of carbonic anhydrase from the teleost fish *Dicentrarchus labrax* (European Seabass) liver and toxicological effects of metals on enzyme activity. Environmental Toxicology and Pharmacology, 32 (1): 69-74. doi: 10.1016/j.etap.2011.03.013.
- Christensen, G.M. and Tucker, J.H. 1976. Effects of selected water toxicants on the *in vitro* activity of fish carbonic anhydrase. Chemico Biological Interactions, 13: 181-192. doi:10.1016/0009-2797(76)90007-7.
- Coban T.A., Beydemir, S., Gulcin, I. and Ekinci, D. 2008. The effect of ethanol on erythrocyte carbonic anhydrase isoenzymes activity: an *in vitro* and *in vivo* study. Journal of Enzyme Inhibition and Medicinal Chemistry, 23: 266-270. doi:10.1080/14756360701474780.
- Conley, D.M. and Mallatt, J. 1988. Histochemical localization of Na^+ - K^+ ATPase and carbonic anhydrase activity in gills of 17 fish species. Canadian Journal of Zoology, 66: 2398-2405.
- Doğan, S. 2006. The *in vitro* effects of some pesticides on carbonic anhydrase activity of *Oncorhynchus mykiss* and *Cyprinus carpio carpio* fish. Journal of Hazardous Materials, 132: 171-176. doi:10.1016/j.jhazmat.2005.10.006
- Ekinci, D. and Beydemir, Ş. 2010. Risk assessment of pesticides and fungicides for acid-base regulation and salt transport in rainbow trout tissues. Pesticide Biochemistry and Physiology, 97: 66-70. doi:10.1016/j.pestbp.2009.12.006.
- Ekinci, D., Beydemir, Ş. and Küfrevioğlu, Ö.İ. 2007. *In vitro* inhibitory effects of some heavy metals on human erythrocyte carbonic anhydrases. Journal of Enzyme Inhibition and Medicinal Chemistry, 22: 745-750. doi: 10.1080/14756360601176048.
- Esbaugh, A.J. and Tufts, B.L. 2006. The structure and function of carbonic anhydrase isozymes in the respiratory system of vertebrates. Respiratory Physiology and Neurobiology, 154: 185-198. doi:10.1016/j.resp.2006.03.007.
- Esbaugh, A.J., Perry, S.F., Bayaa, M., Georgalis, T., Nickerson, J., Tufts, B.L. and Gilmour, K.M., 2005. Cytoplasmic carbonic anhydrase isozymes in rainbow trout *Oncorhynchus mykiss*: comparative physiology

- and molecular evolution. *Journal of Experimental Biology*, 208: 1951–1961. doi: 10.1242/jeb.01551.
- Evans, D.H., Piermarini, P.M. and Choe, K.P. 2005. The multifunctional fish gill: Dominant site of gas exchange, osmoregulation, acid-base regulation, and excretion of nitrogenous waste. *Physiological Reviews*, 85: 97-177. doi:10.1152/physrev.00050.2003.
- Geers, C. and Gros, G. 2000. Carbon Dioxide Transport And Carbonic Anhydrase In Blood And Muscle. *Physiological Reviews*, 80: 681-715.
- Henry, R.P., Smatresk, N.J. and Cameron, J.N. 1988. The distribution of branchial carbonic anhydrase and the effects of gill and erythrocyte carbonic anhydrase inhibition in the channel catfish *Ictalurus punctatus*. *Journal of Experimental Biology*, 134: 201-218.
- Hisar, O., Beydemir, Ş., Gülçin, İ., Hisar, Ş.A., Yanık, T. and Küfrevioğlu, Ö.İ. 2005. The effects of melatonin hormone on carbonic anhydrase enzyme activity in rainbow trout (*Oncorhynchus mykiss*) erythrocytes *in vitro* and *in vivo*. *Turkish Journal of Veterinary and Animal Sciences*, 29: 841–845.
- Hisar, Ş.A., Hisar, O., Yanık, T. and Aras, S.M. 2004. Inhibitory effects of ammonia and urea on gill carbonic anhydrase enzyme activity of rainbow trout (*Oncorhynchus mykiss*). *Environmental Toxicology and Pharmacology*, 17: 125–128. doi:10.1016/j.etap.2004.03.009.
- Ivanov, B.N., Ignatova, L.K. and Romanova, A.K. 2007. Diversity in forms and functions of carbonic anhydrase in terrestrial higher plants. *Russian Journal of Plant Physiology*, 54: 143–162. doi:10.1134/S102144370702001X.
- Kalay, M. and Canlı, M. 2000. Elimination of essential (Cu, Zn) and nonessential (Cd, Pb) metals from tissues of a freshwater fish *Tilapia zillii* following an uptake protocol. *Turkish Journal of Zoology*, 24: 429–436.
- Laemmli, U.K. 1970. Cleavage of structural proteins during assembly of the head of bacteriophage T4. *Nature*, 227: 680–683. doi:10.1038/227680a0.
- Le Trang, N., Bhargava, K.K. and Cerami, A. 1983. Purification of glutathione reductase from gerbil liver in two steps. *Analytical Biochemistry*, 133: 94–99. doi:10.1016/0003-2697(83)90226-9.
- Lineweaver, H. and Burk, D. 1934. The determination of enzyme dissociation constants. *Journal of American Chemical Society* 56: 658–666. doi: 10.1021/ja01318a036.
- Lionetto, M.G., Caricato, R., Giordano, M.E., Erroi, E. and Schettino, T. 2012. Carbonic anhydrase and heavy metals, In: D. Ekinci (Ed.), *Biochemistry*, InTech Open Press, Croatia: 205-224.
- Lionetto, M.G., Giordano, M.E., Vilella, S. and Schettino, T. 2000. Inhibition of eel enzymatic activities by cadmium. *Aquatic Toxicology*, 48: 561–571. doi:10.1016/S0166-445X(99)00056-9.
- Mendil, D., Ünal, Ö.F., Tüzen, M. and Soylak, M. 2010. Determination of trace metals in different fish species and sediments from the River Yeşilirmak in Tokat, Turkey. *Food and Chemical Toxicology*, 48: 1383–1392. doi:10.1016/j.fct.2010.03.006.
- Perry, S.F. and Laurent, P. 1990. The role of carbonic anhydrase in carbon dioxide excretion, acid–base balance and ionic regulation in aquatic gill breathers. In: J.P. Truchot and B. Lahlou (Ed.), *Animal Nutrition and Transport Processes. 2. Transport, Respiration and Excretion: Comparative and Environmental Aspects*, Karger, Basel: 39–57.
- Playle, R.C. 1998. Modelling metal interactions at fish gills. *The Science of the Total Environment*, 219: 147–163. doi:10.1016/S0048-9697(98)00232-0.
- Sender, S., Bottcher, K., Cetin, Y. and Gros, G. 1999. Carbonic anhydrase in the gills of seawater and freshwater-acclimated flounders *Platichthys flesus*: purification, characterization, and immunohistochemical localization. *Journal Of Histochemistry and Cytochemistry*, 47: 43–50. doi: 10.1177/002215549904700105.
- Şentürk, M., Gülçin, İ., Daştan, A., Küfrevioğlu, Ö.İ. and Supuran, C.T. 2009. Carbonic anhydrase inhibitors: Inhibitor of human erythrocyte I and II isoenzymes with antioxidant phenolic compounds. *Bioorganic and Medicinal Chemistry*, 17: 3207–3211.
- Skaggs, S.H., and Henry P.R. 2002. Inhibition of carbonic anhydrase in the gills of two euryhaline crabs, *Callinectes sapidus* and *Carcinus maenas*, by heavy metals. *Comparative Biochemistry and Physiology Part C*, 133: 605–612. doi:10.1016/S1532-0456(02)00175-8.
- Söyüt, H. and Beydemir Ş. 2008. Purification and some kinetic properties of carbonic anhydrase from rainbow trout (*Oncorhynchus mykiss*) liver and metal inhibition. *Protein and Peptide Letters*, 15: 528–535. doi: 10.2174/092986608784567627.
- Söyüt, H. and Beydemir Ş. 2011. The impact of heavy metals on the activity of carbonic anhydrase from rainbow trout (*Oncorhynchus mykiss*) kidney. *Toxicology and Industrial Health*, 28: 296-305. doi: 10.1177/0748233711410914.
- Söyüt, H., Beydemir, Ş. and Hisar, O. 2008. Effects of some metals on carbonic anhydrase from brains of rainbow trout. *Biological Trace Element Research*, 123: 179–190. doi:10.1007/s12011-008-8108-9.
- Supuran, C.T. 2008. Carbonic anhydrases: novel therapeutic applications for inhibitors and activators. *Nature Reviews Drug Discovery*, 7: 81–168. doi: 10.1038/nrd2467.
- Supuran, C.T. 2010. Carbonic anhydrase inhibitor. *Bioorganic and Medicinal Chemistry Letters*, 20: 3467-3474. doi:10.1016/j.bmcl.2010.05.009.
- Vitale, A.M., Monserrat, J.M., Castilho, P. and Rodriguez, E.M. 1999. Inhibitory effects of cadmium on carbonic anhydrase activity and ionic regulation of the estuarine crab *Chasmagnathus granulata* (Decapoda, Grapsidae). *Comparative Biochemistry Physiology C*, 122: 121–129. doi:10.1016/S0742-8413(98)10094-4.
- Wilbur, K.M. and Anderson, N.G. 1948. Electrometric and colorimetric determination of carbonic anhydrase. *Journal of Biological Chemistry*, 176: 147–154.
- Zhang, B.Y., Yang, F., Wang, G.C. and Peng, G. 2010. Cloning and quantitative analysis of the carbonic anhydrase gene from *Porphyra yezoensis*. *Journal of Phycology*, 46:290-296. doi: 10.1111/j.1529-8817.2009.00801.x.